

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2003 (15.05.2003)

PCT

(10) International Publication Number
WO 03/040108 A1

(51) International Patent Classification⁷: **C07D 239/94**,
401/14, 401/12, 409/12, 403/12, 417/14, 413/14, 409/14,
A61K 31/505, A61P 35/00

Robert, Hugh [GB/GB]; Alderley Park, Macclesfield,
Cheshire SK10 4TG (GB).

(21) International Application Number: PCT/GB02/04931

(74) Agent: **ASTRAZENECA**; Global Intellectual Property,
Mersey, Alderley Park, Macclesfield, Cheshire SK10
4TG (GB).

(22) International Filing Date: 31 October 2002 (31.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0126433.2 3 November 2001 (03.11.2001) GB
0129059.2 5 December 2001 (05.12.2001) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except MG, US*): **AS-
TRAZENECA AB** [SE/SE]; S-151 85 Sodertalje (SE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for MG only*): **ASTRAZENECA UK LIM-
ITED** [GB/GB]; 15 Stanhope Gate, London, Greater Lon-
don W1K 1LN (GB).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **HENNEQUIN,
Laurent, Francois, Andre** [FR/GB]; Alderley Park, Mac-
clesfield, Cheshire SK10 4TG (GB). **KETTLE, Jason,
Grant** [GB/GB]; Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). **PASS, Martin** [GB/GB]; Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). **BRADBURY,**

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: QUINAZOLINE DERIVATIVES AS ANTITUMOR AGENTS



(I)

(57) Abstract: The invention concerns quinazoline derivatives of Formula (I); wherein each of Q¹, Z, R¹ and Q² have any of the meanings defined in the description; processes for their preparation, pharmaceutical compositions containing them and their use in the manufacture of a medicament for use in the prevention or treatment of tumours which are sensitive to inhibition of erbB receptor tyrosine kinases.

QUINAZOLINE DERIVATIVES AS ANTITUMOR AGENTS

The invention concerns certain novel quinazoline derivatives, or
5 pharmaceutically-acceptable salts thereof, which possess anti-tumour activity and are
accordingly useful in methods of treatment of the human or animal body. The invention also
concerns processes for the manufacture of said quinazoline derivatives, to pharmaceutical
compositions containing them and to their use in therapeutic methods, for example in the
manufacture of medicaments for use in the prevention or treatment of solid tumour disease in
10 a warm-blooded animal such as man.

Many of the current treatment regimes for diseases resulting from the abnormal
regulation of cellular proliferation such as psoriasis and cancer, utilise compounds that inhibit
DNA synthesis and cellular proliferation. To date, compounds used in such treatments are
generally toxic to cells however their enhanced effects on rapidly dividing cells such as
15 tumour cells can be beneficial. Alternative approaches to these cytotoxic anti-tumour agents
are currently being developed, for example selective inhibitors of cell signalling pathways.
These types of inhibitors are likely to have the potential to display an enhanced selectivity of
action against tumour cells and so are likely to reduce the probability of the therapy possessing
unwanted side effects.

20 Eukaryotic cells are continually responding to many diverse extracellular signals that
enable communication between cells within an organism. These signals regulate a wide
variety of physical responses in the cell including proliferation, differentiation, apoptosis and
motility. The extracellular signals take the form of a diverse variety of soluble factors
including growth factors as well as paracrine and endocrine factors. By binding to specific
25 transmembrane receptors, these ligands integrate the extracellular signal to the intracellular
signalling pathways, therefore transducing the signal across the plasma membrane and
allowing the individual cell to respond to its extracellular signals. Many of these signal
transduction processes utilise the reversible process of the phosphorylation of proteins that are
involved in the promotion of these diverse cellular responses. The phosphorylation status of
30 target proteins is regulated by specific kinases and phosphatases that are responsible for the
regulation of about one third of all proteins encoded by the mammalian genome. As
phosphorylation is such an important regulatory mechanism in the signal transduction process,
it is therefore not surprising that aberrations in these intracellular pathways result in abnormal

cell growth and differentiation and so promote cellular transformation (reviewed in Cohen *et al*, Curr Opin Chem Biol, 1999, 3, 459-465).

It has been widely shown that a number of these tyrosine kinases are mutated to constitutively active forms and/or when over-expressed result in the transformation of a variety of human cells. These mutated and over-expressed forms of the kinase are present in a large proportion of human tumours (reviewed in Kolibaba *et al*, Biochimica et Biophysica Acta, 1997, 133, F217-F248). As tyrosine kinases play fundamental roles in the proliferation and differentiation of a variety of tissues, much focus has centred on these enzymes in the development of novel anti-cancer therapies. This family of enzymes is divided into two groups - receptor and non-receptor tyrosine kinases e.g. EGF Receptors and the SRC family respectively. From the results of a large number of studies including the Human Genome Project, about 90 tyrosine kinase have been identified in the human genome, of this 58 are of the receptor type and 32 are of the non-receptor type. These can be compartmentalised in to 20 receptor tyrosine kinase and 10 non-receptor tyrosine kinase sub-families (Robinson *et al*, Oncogene, 2000, 19, 5548-5557).

The receptor tyrosine kinases are of particular importance in the transmission of mitogenic signals that initiate cellular replication. These large glycoproteins, which span the plasma membrane of the cell possess an extracellular binding domain for their specific ligands (such as Epidermal Growth Factor (EGF) for the EGF Receptor). Binding of ligand results in the activation of the receptor's kinase enzymatic activity that is encoded by the intracellular portion of the receptor. This activity phosphorylates key tyrosine amino acids in target proteins, resulting in the transduction of proliferative signals across the plasma membrane of the cell.

It is known that the erbB family of receptor tyrosine kinases, which include EGFR, erbB2, erbB3 and erbB4, are frequently involved in driving the proliferation and survival of tumour cells (reviewed in Olayioye *et al*, EMBO J, 2000, 19, 3159). One mechanism in which this can be accomplished is by overexpression of the receptor at the protein level, generally as a result of gene amplification. This has been observed in many common human cancers (reviewed in Klapper *et al*, Adv. Cancer Res., 2000, 77, 25) such as breast cancer (Sainsbury *et al*, Brit. J. Cancer, 1988, 58, 458; Guerin *et al*, Oncogene Res., 1988, 3, 21; Slamon *et al*, Science, 1989, 244, 707; Klijn *et al*, Breast Cancer Res. Treat., 1994, 29, 73 and reviewed in Salomon *et al*, Crit. Rev. Oncol. Hematol., 1995, 19, 183), non-small cell lung cancers (NSCLCs) including adenocarcinomas (Cerny *et al*, Brit. J. Cancer, 1986, 54,

265; Reubi *et al.*, Int. J. Cancer, 1990, 45, 269; Rusch *et al.*, Cancer Research, 1993, 53, 2379; Brabender *et al.*, Clin. Cancer Res., 2001, 7, 1850) as well as other cancers of the lung (Hendler *et al.*, Cancer Cells, 1989, 7, 347; Ohsaki *et al.*, Oncol. Rep., 2000, 7, 603), bladder cancer (Neal *et al.*, Lancet, 1985, 366; Chow *et al.*, Clin. Cancer Res., 2001, 7, 1957, Zhau *et al.*, Mol. Carcinog., 3, 254), oesophageal cancer (Mukaida *et al.*, Cancer, 1991, 68, 142), gastrointestinal cancer such as colon, rectal or stomach cancer (Bolen *et al.*, Oncogene Res., 1987, 1, 149; Kapitanovic *et al.*, Gastroenterology, 2000, 112, 1103; Ross *et al.*, Cancer Invest., 2001, 19, 554), cancer of the prostate (Visakorpi *et al.*, Histochem. J., 1992, 24, 481; Kumar *et al.*, 2000, 32, 73; Scher *et al.*, J. Natl. Cancer Inst., 2000, 92, 1866), leukaemia (Konaka *et al.*, Cell, 1984, 37, 1035, Martin-Subero *et al.*, Cancer Genet Cytogenet., 2001, 127, 174), ovarian (Hellstrom *et al.*, Cancer Res., 2001, 61, 2420), head and neck (Shiga *et al.*, Head Neck, 2000, 22, 599) or pancreatic cancer (Ovotny *et al.*, Neoplasma, 2001, 48, 188). As more human tumour tissues are tested for expression of the erbB family of receptor tyrosine kinases it is expected that their widespread prevalence and importance will be further enhanced in the future.

As a consequence of the mis-regulation of one or more of these receptors (in particular erbB2), it is widely believed that many tumours become clinically more aggressive and so correlate with a poorer prognosis for the patient (Brabender *et al.*, Clin. Cancer Res., 2001, 7, 1850; Ross *et al.*, Cancer Investigation, 2001, 19, 554, Yu *et al.*, Bioessays, 2000, 22, 7, 673). In addition to these clinical findings, a wealth of pre-clinical information suggests that the erbB family of receptor tyrosine kinases are involved in cellular transformation. This includes the observations that many tumour cell lines overexpress one or more of the erbB receptors and that EGFR or erbB2 when transfected into non-tumour cells have the ability to transform these cells. This tumourigenic potential has been further verified as transgenic mice that overexpress erbB2 spontaneously develop tumours in the mammary gland. In addition to this, a number of pre-clinical studies have demonstrated that anti-proliferative effects can be induced by knocking out one or more erbB activities by small molecule inhibitors, dominant negatives or inhibitory antibodies (reviewed in Mendelsohn *et al.*, Oncogene, 2000, 19, 6550). Thus it has been recognised that inhibitors of these receptor tyrosine kinases should be of value as a selective inhibitor of the proliferation of mammalian cancer cells (Yaish *et al.*, Science, 1988, 242, 933, Kolibaba *et al.*, Biochimica et Biophysica Acta, 1997, 133, F217-F248; Al-Obeidi *et al.*, 2000, Oncogene, 19, 5690-5701; Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565). In addition to this pre-clinical data, findings using inhibitory

antibodies against EGFR and erbB2 (c-225 and trastuzumab respectively) have proven to be beneficial in the clinic for the treatment of selected solid tumours (reviewed in Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565).

Amplification and/or activity of members of the ErbB type receptor tyrosine kinases have been detected and so have been implicated to play a role in a number of non-malignant proliferative disorders such as psoriasis (Ben-Bassat, Curr. Pharm. Des., 2000, 6, 933; Elder *et al.*, Science, 1989, 243, 811), benign prostatic hyperplasia (BPH) (Kumar *et al.*, Int. Urol. Nephrol., 2000, 32, 73), atherosclerosis and restenosis (Bokemeyer *et al.*, Kidney Int., 2000, 58, 549). It is therefore expected that inhibitors of erbB type receptor tyrosine kinases will be useful in the treatment of these and other non-malignant disorders of excessive cellular proliferation.

International Patent Applications WO 96/33977, WO 96/33978, WO 96/33979, WO 96/33980 and WO 96/33981 disclose that certain quinazoline derivatives which bear an anilino substituent at the 4-position possess receptor tyrosine kinase inhibitory activity.

A review of the structure activity relationship of various quinazoline derivatives is disclosed by G. W. Rewcastle *et al.* (J. Med. Chem. 1995, 38, 3428-3487), including a number of 5-substituted compounds. However, such 5-substituted compounds are stated to have low in-vitro activity as EGFR tyrosine kinase inhibitors compared to quinazolines substituted at the 6- and 7- positions.


WO 96/09294 discloses 4-anilinoquinazoline derivatives, including 5-chloro and 5-methoxy substituted quinazoline derivatives as protein tyrosine kinase inhibitors.

Co-pending International Patent Application PCT/GB01/02424 discloses that certain quinazoline derivatives which carry a 5-substituent are inhibitors of the Src family of non-receptor tyrosine kinases, such as c-Src, c-Yes and c-Fyn.

We have now found that surprisingly certain 5-substituted quinazoline derivatives possess potent anti-tumour activity. Without wishing to imply that the compounds disclosed in the present invention possess pharmacological activity only by virtue of an effect on a single biological process, it is believed that the compounds provide an anti-tumour effect by way of inhibition of one or more of the erbB family of receptor tyrosine kinases that are involved in the signal transduction steps which lead to the proliferation of tumour cells. In particular, it is believed that the compounds of the present invention provide an anti-tumour effect by way of inhibition of EGFR and/or erbB2 receptor tyrosine kinases.

Generally the compounds of the present invention possess potent inhibitory activity against the erbB receptor tyrosine kinase family, for example by inhibition of EGFR and/or erbB2 and/or erbB4 receptor tyrosine kinases, whilst possessing less potent inhibitory activity against other kinases. Furthermore, generally the compounds of the present invention possess substantially better potency against the erbB2 over that of the EGFR tyrosine kinase, thus potentially providing effective treatment for erbB2 driven tumours. The invention also includes compounds that are active against all or a combination of EGFR, erbB2 and erbB4 receptor tyrosine kinases, thus potentially providing treatments for conditions mediated by one or more of these receptor tyrosine kinases.

10 According to a first aspect of the invention there is provided a quinazoline derivative of the Formula I



15 wherein

R^1 is hydrogen or (1-6C)alkyl;

Z is a direct bond or is selected from O, S and $N(R^2)$, wherein R^2 is hydrogen or (1-6C)alkyl;

Q^1 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q^1 -Z-group are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, $N(R^3)$, CO, CH(OR³), CON(R³), $N(R^3)CO$, SO₂N(R³), $N(R^3)SO_2$, CH=CH and C≡C wherein R^3 is hydrogen or (1-6C)alkyl,

25 and wherein any CH₂ or CH₃ group within the Q^1 -Z- group optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,

- 6 -

- (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,
- 5 and wherein any heterocyclyl, (3-7C)cycloalkyl or (3-7C)cycloalkenyl group within the Q¹-Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl,
- 10 (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, amino(2-6C)alkanoyl, N-(1-6C)alkylamino(2-6C)alkanoyl, N,N-di-[(1-6C)alkyl]amino(2-6C)alkanoyl, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino,
- 15 N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, from a group of the formula:
- $$-X^1-R^4$$
- wherein X¹ is a direct bond or is selected from O and N(R⁵), wherein R⁵ is hydrogen or (1-6C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl,
- 20 di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, and from a group of the formula:
- $$-X^5-Q^6$$
- 25 wherein X⁵ is a direct bond or is selected from O, CO and N(R¹⁰), wherein R¹⁰ is hydrogen or (1-6C)alkyl, and Q⁶ is heterocyclyl, heterocyclyl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl or (3-7C)cycloalkenyl-(1-6C)alkyl, and wherein any heterocyclyl, (3-7C)cycloalkyl or (3-7C)cycloalkenyl group in Q⁶ optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno,
- 30 hydroxy, cyano, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (1-6C)alkoxycarbonyl, (1-6C)alkylsulphonyl, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl,

- 7 -


carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl,

di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,


N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo
5 or thioxo substituents; and


Q² is selected from a group of formula Ia, Ib, Ic, Id and Ie




Ia




Ib



Ic



Id



Ie

10 wherein G¹, G², G³, G⁴ and G⁵ are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

G⁶ and G⁷ are each, independently, selected from hydrogen and halogeno,

15 X² is a direct bond or is selected from O, S, SO, SO₂, N(R⁶), CH(OR⁶), CON(R⁶), N(R⁶)CO, SO₂N(R⁶), N(R⁶)SO₂, OC(R⁶)₂, C(R⁶)₂O, SC(R⁶)₂, C(R⁶)₂S, CO, C(R⁶)₂N(R⁶) and N(R⁶)C(R⁶)₂ wherein each R⁶ is, independently, hydrogen or (1-6C)alkyl, and Q³ is aryl, or heteroaryl,

- 8 -

or X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

X^3 is a direct bond or is selected from SO_2 , CO, $SO_2N(R^7)$ and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,

- 5 and any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphonamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphonamoyl, N,N-di-[(1-6C)alkyl]sulphonamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,

- 20 N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, or from a group of the formula




wherein Q^5 is selected from aryl, heteroaryl or heterocyclyl which is optionally substituted by

- 25 1 or 2 substituents selected from halogeno, hydroxy, (1-6C)alkyl, (1-6C)alkoxy, amino, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any CH_2 or CH_3 group within a substituent on any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents,

- 30 and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or thioxo substituents,
or a pharmaceutically acceptable salt thereof.

According to a further aspect of the invention there is provided a quinazoline derivative of the Formula I



5

wherein

R^1 is hydrogen or (1-6C)alkyl;

Z is a direct bond or is selected from O, S and $N(R^2)$, wherein R^2 is hydrogen or (1-6C)alkyl;

- 10 Q^1 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q^1 - Z - group are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, $N(R^3)$, CO, CH(OR³), CON(R³), $N(R^3)$ CO, SO₂N(R³), $N(R^3)$ SO₂, CH=CH and C≡C
- 15 wherein R^3 is hydrogen or (1-6C)alkyl,


- and wherein any CH₂ or CH₃ group within the Q^1 - Z - group optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
- 20 (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,
- and wherein any heterocyclyl group within the Q^1 - Z - group optionally bears 1, 2 or 3
- 25 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl,

- 10 -

N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :




wherein X^1 is a direct bond or is selected from O and N(R^5), wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,




and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo or thioxo substituents;

Q^2 is selected from a group of formula Ia, Ib, Ic, Id and Ie


15




Ia




Ib



Ic



Id



Ie

wherein G^1 , G^2 , G^3 , G^4 and G^5 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl,

(2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy,
(1-6C)alkylamino and di-[(1-6C)alkyl]amino,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $CH(OR^6)$, $CON(R^6)$, $OC(R^6)_2$,
 $C(R^6)_2O$, $SC(R^6)_2$, $C(R^6)_2S$, CO, $C(R^6)_2N(R^6)$ and $N(R^6)C(R^6)_2$ wherein each R^6 is,
5 independently, hydrogen or (1-6C)alkyl, and Q^3 is aryl, or heteroaryl,

or X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a
nitrogen atom,

X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is,
independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,
10 and any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$
optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from
halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl,
(2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy,
(1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and
15 di-[(1-6C)alkyl]amino,

and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or
thioxo substituents;
or a pharmaceutically acceptable salt thereof.

According to a further aspect of the invention there is provided a quinazoline
20 derivative of the Formula I wherein each of R^1 and Q^2 has any of the meanings defined
hereinbefore and

Z is O; and

Q^1 is (3-7C)cycloalkyl or heterocyclyl,

and wherein any CH_2 or CH_3 group within the Q^1-Z - group optionally bears on each
25 said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent
selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio,
(1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,
(1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
(2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-
30 (2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl,
(1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,

and wherein any heterocyclyl group within the Q^1-Z - group optionally bears 1, 2 or 3
substituents, which may be the same or different, selected from halogeno, trifluoromethyl,

- 12 -


cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, 5 (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :




wherein X^1 is a direct bond or is selected from O and $N(R^5)$, wherein R^5 is hydrogen or 10 (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, 15 and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo or thioxo substituents;

According to a further aspect of the invention there is provided a quinazoline derivative of the Formula I wherein R^1 has any of the meanings defined hereinbefore and Z is O;

20 Q^1 is (3-7C)cycloalkyl, which optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, from a group of the formula:



wherein X^1 is a direct bond or is selected from O and $N(R^5)$, wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl or N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, 30 and from a group of the formula:




wherein X^5 is a direct bond or is selected from O, CO and $N(R^{10})$, wherein R^{10} is hydrogen or (1-6C)alkyl, and Q^6 is heterocyclyl or heterocyclyl-(1-6C)alkyl, which optionally bears 1 or 2


- 13 -

substituents, which may be the same or different, selected from halogeno, hydroxy, (1-6C)alkyl, (1-6C)alkoxy, amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (1-6C)alkoxycarbonyl, (1-6C)alkylsulphonyl, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkenyl, halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo or thioxo substituents; and


10 Q² is selected from a group of formula Ia, Ib and Id



Ia



Ib



Id


wherein G¹, G², G³, G⁴ and G⁵ are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino, G⁶ and G⁷ are each, independently, selected from hydrogen and halogeno,

- 14 -

X^2 is a direct bond or is selected from O, S, SO, SO₂, N(R⁶), CH(OR⁶), CON(R⁶), N(R⁶)CO, SO₂N(R⁶), N(R⁶)SO₂, OC(R⁶)₂, C(R⁶)₂O, SC(R⁶)₂, C(R⁶)₂S, CO, C(R⁶)₂N(R⁶) and N(R⁶)C(R⁶)₂ wherein each R⁶ is, independently, hydrogen or (1-6C)alkyl, and Q³ is aryl, or heteroaryl,

5 or X^2 is CO and Q³ is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

X^3 is a direct bond or is selected from SO₂, CO, SO₂N(R⁷), and C(R⁷)₂, wherein each R⁷ is, independently, hydrogen or (1-6C)alkyl, and Q⁴ is aryl or heteroaryl, and any aryl, heteroaryl or heterocyclyl group in the group - X^2 -Q³ and - X^3 -Q⁴ optionally
 10 bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
 15 N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



20 wherein X^4 is a direct bond or is selected from O and N(R⁹), wherein R⁹ is hydrogen or (1-6C)alkyl, and R⁸ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,
 25 (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, or from a group of the formula



wherein Q⁵ is selected from aryl, heteroaryl or heterocyclyl which is optionally substituted by 1 or 2 substituents selected from halogeno, hydroxy, (1-6C)alkyl, (1-6C)alkoxy, amino,
 30 (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any CH₂ or CH₃ group within a substituent on any aryl, heteroaryl or heterocyclyl group in the group - X^2 -Q³ and - X^3 -Q⁴ optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents,

- 15 -

and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or thioxo substituents.

According to a further aspect of the invention there is provided a quinazoline derivative of the Formula I wherein R^1 and Z have any of the meanings defined hereinbefore

5 and


Q^1 is (3-7C)cycloalkyl or heterocyclyl,

and wherein any heterocyclyl or (3-7C)cycloalkyl group within the Q^1 -Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, amino(2-6C)alkanoyl, N-(1-6C)alkylamino(2-6C)alkanoyl, N,N-di-[(1-6C)alkyl]amino(2-6C)alkanoyl, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, from a group of the formula:



wherein X^1 is a direct bond or is selected from O and $N(R^5)$, wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl,

25 and from a group of the formula:




wherein X^5 is a direct bond or is selected from O, CO and $N(R^{10})$, wherein R^{10} is hydrogen or (1-6C)alkyl, and Q^6 is heterocyclyl, heterocyclyl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl or (3-7C)cycloalkenyl-(1-6C)alkyl, and 30 wherein any heterocyclyl, (3-7C)cycloalkyl or (3-7C)cycloalkenyl group in Q^6 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, hydroxy, cyano, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (1-6C)alkoxycarbonyl,


- 16 -

- (1-6C)alkylsulphonyl, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,
- 5 N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo or thioxo substituents; and


Q² is selected from a group of formula Ia, Ib and Id



Ia



Ib



Id

10

- wherein G¹, G², G³, G⁴ and G⁵ are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy,
- 15 (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

G⁶ and G⁷ are hydrogen,

X² is a direct bond or is selected from O, S, SO, SO₂, N(R⁶), CH(OR⁶), CON(R⁶), N(R⁶)CO, SO₂N(R⁶), N(R⁶)SO₂, OC(R⁶)₂, C(R⁶)₂O, SC(R⁶)₂, C(R⁶)₂S, CO, C(R⁶)₂N(R⁶) and

$N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or (1-6C)alkyl, and Q^3 is aryl, or heteroaryl,

or X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

- 5 X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl, and any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



- wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl,

- and wherein any CH_2 or CH_3 group within a substituent on any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents,

and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or thioxo substituents,

or a pharmaceutically acceptable salt thereof.

- 30 In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups such as propyl, isopropyl and tert-butyl, and (3-7C)cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. However references to individual alkyl groups such as "propyl" are specific for the straight-chain

- 18 -

version only, references to individual branched-chain alkyl groups such as "isopropyl" are specific for the branched-chain version only and references to individual cycloalkyl groups such as "cyclopentyl" are specific for that 5-membered ring only. An analogous convention applies to other generic terms, for example (1-6C)alkoxy includes methoxy, ethoxy, 5 cyclopropyloxy and cyclopentyloxy, (1-6C)alkylamino includes methylamino, ethylamino, cyclobutylamino and cyclohexylamino, and di-[(1-6Calkyl)]amino includes dimethylamino, diethylamino, N-cyclobutyl-N-methylamino and N-cyclohexyl-N-ethylamino.

It is to be understood that, insofar as certain of the compounds of Formula I defined above may exist in optically active or racemic forms by virtue of one or more asymmetric 10 carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the above-mentioned activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory 15 techniques referred to hereinafter.

It is to be understood that the present invention includes in its definition any and all tautomeric forms of the compounds of the formula I which possess the above mentioned activity.

It is also to be understood that in so far as certain compounds of the formula 1 may 20 exist in solvated forms as well as unsolvated forms, for example, hydrated forms, the present invention includes any and all such solvated forms, which possess the above mentioned activity.

Suitable values for the generic radicals referred to above include those set out below.

A suitable value for any one of the 'Q' groups (Q^1 , Q^3 to Q^5) when it is aryl or for the 25 aryl group within a 'Q' group is, for example, phenyl or naphthyl, preferably phenyl.

A suitable value for any one of the 'Q' groups (Q^1 or Q^6) when it is (3-7C)cycloalkyl or for the (3-7C)cycloalkyl group within a 'Q' group is, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or bicyclo[2.2.1]heptyl and a suitable value for any one of the 'Q' groups (Q^1 or Q^6) when it is (3-7C)cycloalkenyl or for the 30 (3-7C)cycloalkenyl group within a 'Q' group is, for example, cyclobutenyl, cyclopentenyl, cyclohexenyl or cycloheptenyl.

A suitable value for any one of the 'Q' groups (Q^3 to Q^5) when it is heteroaryl or for the heteroaryl group within a 'Q' group is, for example, an aromatic 5- or 6-membered

monocyclic ring or a 9- or 10-membered bicyclic ring with up to five ring heteroatoms selected from oxygen, nitrogen and sulphur, which, unless specified otherwise, may be carbon or nitrogen linked. Examples of suitable values of "heteroaryl" include furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazenyl, 1,3-benzodioxolyl, benzofuranyl, indolyl, benzothienyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, benzothiazolyl, imidazopyridinyl, indazolyl, benzofurazanyl, quinolyl, isoquinolyl, quinazolinyl, quinoxalyl, cinnolinyl or naphthyridinyl. Such as furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3-benzodioxolyl, benzofuranyl, indolyl, benzothienyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, indazolyl, benzofurazanyl, quinolyl, isoquinolyl, quinazolinyl, quinoxalyl, cinnolinyl or naphthyridinyl.

A suitable value for any one of the 'Q' groups (Q^1 , Q^3 , Q^5 or Q^6) when it is heterocyclyl or for the heterocyclyl group within a 'Q' group is, for example, a non-aromatic saturated or partially saturated 3 to 10 membered monocyclic or bicyclic ring with up to five heteroatoms selected from oxygen, nitrogen and sulphur, which, unless specified otherwise, may be carbon or nitrogen linked. Examples of suitable values of "heterocyclyl" include oxiranyl, oxetanyl, azetidyl, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, pyrrolinyl, pyrrolidinyl, morpholinyl, tetrahydro-1,4-thiazinyl, 1,1-dioxotetrahydro-1,4-thiazinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, tetrahydropyrimidinyl, tetrahydrothienyl, tetrahydrothiopyranyl, decahydroisoquinolinyl or decahydroquinolinyl, preferably tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, morpholinyl, 1,4-oxazepanyl, 1,1-dioxotetrahydro-4H-1,4-thiazinyl, piperidinyl or piperazinyl, more preferably tetrahydrofuran-3-yl, tetrahydropyran-4-yl, tetrahydrothien-3-yl, tetrahydrothiopyran-4-yl, pyrrolidin-3-yl, morpholino, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-4-yl, piperidin-3-yl or piperazin-1-yl. A nitrogen or sulphur atom within a heterocyclyl group may be oxidized to give the corresponding N or S oxide, for example 1,1-dioxotetrahydrothienyl, 1-oxotetrahydrothienyl, 1,1-dioxotetrahydrothiopyranyl or 1-oxotetrahydrothiopyranyl. A suitable value for such a group which bears 1 or 2 oxo or thioxo substituents is, for example, 2-oxopyrrolidinyl, 2-thioxopyrrolidinyl,

2-oxoimidazolidinyl, 2-thioxoimidazolidinyl, 2-oxopiperidinyl, 2,5-dioxopyrrolidinyl, 2,5-dioxoimidazolidinyl or 2,6-dioxopiperidinyl.

A suitable value for any one of the 'Q' groups (such as Q³) when it is a nitrogen containing heterocyclyl group is, for example, a non-aromatic saturated or partially saturated 3 to 10 membered monocyclic or bicyclic ring with up to five heteroatoms selected from oxygen, nitrogen and sulphur, provided at least one heteroatom is nitrogen. Suitable values include, for example, those heterocyclic groups mentioned above that contain at least one nitrogen atom, for example azetidyl, pyrrolinyl, pyrrolidinyl, morpholinyl (including morpholino), tetrahydro-1,4-thiazinyl, 1,1-dioxotetrahydro-1,4-thiazinyl, piperidinyl (including piperidino), homopiperidinyl, piperazinyl, homopiperazinyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, tetrahydropyrimidinyl, decahydroisoquinolinyl or decahydroquinolinyl.

A suitable value for a 'Q' group when it is heterocyclyl-(1-6C)alkyl is, for example, heterocyclylmethyl, 2-heterocyclylethyl and 3-heterocyclylpropyl. The invention comprises corresponding suitable values for 'Q' groups when, for example, rather than a heterocyclyl-(1-6C)alkyl group, an (3-7C)cycloalkyl-(1-6C)alkyl or (3-7C)cycloalkenyl-(1-6C)alkyl is present.

Suitable values for any of the 'R' groups (R¹ to R⁹), or for any of the 'G' groups (G¹ to G⁷) within Q², or for various groups within Q², or for Q¹ or for various groups within Q¹, or for various groups within the Q¹-Z- group include:-

for halogeno	fluoro, chloro, bromo and iodo;
for (1-6C)alkyl:	methyl, ethyl, propyl, isopropyl and <u>tert</u> -butyl;
for (2-8C)alkenyl:	vinyl, isopropenyl, allyl and but-2-enyl;
for (2-8C)alkynyl:	ethynyl, 2-propynyl and but-2-ynyl;
25 for (1-6C)alkoxy:	methoxy, ethoxy, propoxy, isopropoxy and butoxy;
for (2-6C)alkenyloxy:	vinylloxy and allyloxy;
for (2-6C)alkynyloxy:	ethynyloxy and 2-propynyloxy;
for (1-6C)alkylthio:	methylthio, ethylthio and propylthio;
for (1-6C)alkylsulphinyl:	methylsulphinyl and ethylsulphinyl;
30 for (1-6C)alkylsulphonyl:	methylsulphonyl and ethylsulphonyl;
for (1-6C)alkylamino:	methylamino, ethylamino, propylamino, isopropylamino and butylamino;
for di-[(1-6C)alkyl]amino:	dimethylamino, diethylamino, <u>N</u> -ethyl-

- 21 -

- N-methylamino and diisopropylamino;
 for (1-6C)alkoxycarbonyl: methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and tert-butoxycarbonyl;
 for N-(1-6C)alkylcarbamoyl: N-methylcarbamoyl, N-ethylcarbamoyl and
 5 N-propylcarbamoyl;
 for N,N-di-[(1-6C)alkyl]carbamoyl: N,N-dimethylcarbamoyl, N-ethyl-
N-methylcarbamoyl and N,N-diethylcarbamoyl;
 for (2-6C)alkanoyl: acetyl and propionyl;
 for (2-6C)alkanoyloxy: acetoxy and propionyloxy;
 10 for (2-6C)alkanoylamino: acetamido and propionamido;
 for amino(2-6C)alkanoyl: aminoacetyl and 2-aminopropionyl;
 for N-(1-6C)alkylamino(2-6C)alkanoyl: N-methylaminoacetyl and 2-(N-
 methylaminopropionyl;
 for N,N-di-[(1-6C)alkyl]amino(2-6C)alkanoyl: N,N-di-methylaminoacetyl;
 15 for N-(1-6C)alkyl-(2-6C)alkanoylamino: N-methylacetamido and N-methylpropionamido;
 for N-(1-6C)alkylsulphamoyl: N-methylsulphamoyl and N-ethylsulphamoyl;
 for N,N-di-[(1-6C)alkyl]sulphamoyl: N,N-dimethylsulphamoyl;
 for (1-6C)alkanesulphonylamino: methanesulphonylamino and ethanesulphonylamino;
 for N-(1-6C)alkyl-(1-6C)alkanesulphonylamino: N-methylmethanesulphonylamino and
 20 N-methylethanesulphonylamino;
 for (3-6C)alkenoylamino: acrylamido, methacrylamido and crotonamido;
 for N-(1-6C)alkyl-(3-6C)alkenoylamino: N-methylacrylamido and N-methylcrotonamido;
 for (3-6C)alkynoylamino: propiolamido;
 for N-(1-6C)alkyl-(3-6C)alkynoylamino: N-methylpropiolamido;
 25 for amino-(1-6C)alkyl: aminomethyl, 2-aminoethyl, 1-aminoethyl and
 3-aminopropyl;
 for (1-6C)alkylamino-(1-6C)alkyl: methylaminomethyl, ethylaminomethyl,
 1-methylaminoethyl, 2-methylaminoethyl,
 2-ethylaminoethyl and 3-methylaminopropyl;
 30 for di-[(1-6C)alkyl]amino-(1-6C)alkyl: dimethylaminomethyl, diethylaminomethyl,
 1-dimethylaminoethyl, 2-dimethylaminoethyl and
 3-dimethylaminopropyl;
 for halogeno-(1-6C)alkyl: chloromethyl, 2-chloroethyl, 1-chloroethyl and

- 22 -

- for hydroxy-(1-6C)alkyl: 3-chloropropyl;
hydroxymethyl, 2-hydroxyethyl, 1-hydroxyethyl and
3-hydroxypropyl;
- for (1-6C)alkoxy-(1-6C)alkyl: methoxymethyl, ethoxymethyl, 1-methoxyethyl,
5 2-methoxyethyl, 2-ethoxyethyl and
3-methoxypropyl;
- for cyano-(1-6C)alkyl: cyanomethyl, 2-cyanoethyl, 1-cyanoethyl and
3-cyanopropyl;
- for carboxy-(1-6C)alkyl: carboxymethyl, 2-carboxyethyl, 1-carboxyethyl and
10 3-carboxypropyl;
- for (1-6C)alkylthio-(1-6C)alkyl: methylthiomethyl, ethylthiomethyl,
2-methylthioethyl, 1-methylthioethyl and
3-methylthiopropyl;
- for (1-6C)alkylsulphinyl-(1-6C)alkyl: methylsulphinylmethyl, ethylsulphinylmethyl,
15 2-methylsulphinylethyl, 1-methylsulphinylethyl and
3-methylsulphinylpropyl;
- for (1-6C)alkylsulphonyl-(1-6C)alkyl: methylsulphonylmethyl, ethylsulphonylmethyl,
2-methylsulphonylethyl, 1-methylsulphonylethyl and
3-methylsulphonylpropyl;
- 20 for (2-6C)alkanoylamino-(1-6C)alkyl: acetamidomethyl, propionamidomethyl and
2-acetamidoethyl;
- for (1-6C)alkoxycarbonyl-(1-6C)alkyl: methoxycarbonylmethyl, 2-methoxycarbonylethyl
and 2-ethoxycarbonylethyl;
- for (1-6C)alkoxycarbonylamino-(1-6C)alkyl: methoxycarbonylaminomethyl,
25 ethoxycarbonylaminomethyl,
tert-butoxycarbonylaminomethyl and
2-methoxycarbonylaminoethyl;
- for (1-6C)alkoxycarbonylamino-(1-6C)alkyl: methoxycarbonylaminomethyl,
ethoxycarbonylaminomethyl,
30 tert-butoxycarbonylaminomethyl and
2-methoxycarbonylaminoethyl;
- for carbamoyl-(1-6C)alkyl: carbamoylmethyl, 1-carbamoylethyl,
2-carbamoylethyl and 3-carbamoylpropyl;

- 23 -

for (2-6C)alkanoyl-(1-6C)alkyl: acetylmethyl and 2-acetylethyl;
 for N-(1-6C)alkylcarbamoyl-(1-6C)alkyl: N-methylcarbamoylmethyl,
N-ethylcarbamoylmethyl, N-propylcarbamoylmethyl,
 1-(N-methylcarbamoyl)ethyl,
 5 1-(N-ethylcarbamoyl)ethyl,
 2-(N-methylcarbamoyl)ethyl,
 2-(N-ethylcarbamoyl)ethyl and
 3-(N-methylcarbamoyl)propyl; and
 for N,N-di[(1-6C)alkyl]carbamoyl-(1-6C)alkyl: N,N-dimethylcarbamoylmethyl,
 10 N,N-diethylcarbamoylmethyl,
 2-(N,N-dimethylcarbamoyl)ethyl, and
 3-(N,N-dimethylcarbamoyl)propyl.

When in this specification reference is made to a (1-4C)alkyl group it is to be understood that such groups refer to alkyl groups containing up to 4 carbon atoms. A skilled person will realise that representative examples of such groups are those listed above under (1-6C)alkyl that contain up to 4 carbon atoms, such as methyl, ethyl, propyl and butyl. A similar convention is adopted for the other groups listed above such as (1-4C)alkoxy; (2-4C)alkenyl, (2-4C)alkynyl and (2-4C)alkanoyl.

When, as defined hereinbefore, in the group of the formula $-X^2-Q^3$, X^2 is, for example, a $OC(R^6)_2$ linking group, it is the oxygen atom, not the carbon atom, of the $OC(R^6)_2$ linking group which is attached to the phenyl ring in the formula Ia and the carbon atom is attached to the Q^3 group. It is to be understood that when X^3 is $SO_2N(R^7)$, as defined hereinbefore, the sulphur atom of the $SO_2N(R^7)$ linking group is attached to the nitrogen atom in formulae 1b, 1c, 1d or 1e and the nitrogen atom of the $SO_2N(R^7)$ linking group is attached to Q^4 .

As defined hereinbefore, adjacent carbon atoms in any (2-6C)alkylene chain within a Q^1-Z - group may be optionally separated by the insertion into the chain of a group such as O, $CON(R^3)$ or $C\equiv C$. For example, insertion of a $C\equiv C$ group into the ethylene chain within a 2-morpholinoethoxy group gives rise to a 4-morpholinobut-2-ynyloxy group.

When, as defined hereinbefore, any CH_2 or CH_3 group within a Q^1-Z - group optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents, there are suitably 1 or 2 halogeno or (1-6C)alkyl substituents present on each said CH_2 group and there are suitably 1, 2 or 3 such substituents present on each said CH_3 group.

When, as defined hereinbefore, any CH₂ or CH₃ group within a Q¹-Z- group optionally bears on each said CH₂ or CH₃ group a substituent as defined hereinbefore, suitable substituents so formed include, for example, hydroxy-substituted heterocyclyl- (1-6C)alkoxy groups such as 2-hydroxy-3-piperidinopropoxy and 2-hydroxy-3-morpholinopropoxy, hydroxy-substituted heterocyclyl- (1-6C)alkylamino groups such as 2-hydroxy-3-piperidinopropylamino and 2-hydroxy-3-morpholinopropylamino, heterocyclyl-substituted (1-6C)alkylamino-(1-6C)alkyl groups such as 2-morpholinoethylaminomethyl, 2-piperazin-1-ylethylaminomethyl and 3-morpholinopropylaminomethyl and halogen substituted alkyl groups, for example di-fluoromethyl and 2,2-difluoroethyl.

Similar considerations apply to the attachments and substitutions within the -X²-Q³ and -X³-Q⁴ groups present in Q².

It is to be understood that when, as defined hereinbefore, any CH₂ or CH₃ group within a Q¹-Z- group optionally bears on each said CH₂ or CH₃ group a substituent as defined hereinbefore, the optional substituent may be present on any CH₂ or CH₃ group within a Q¹-Z- group, including those on the hereinbefore defined substituents that may be present on an aryl, heteroaryl or heterocyclyl groups within Q¹-Z-. For example, if Q¹ is a 1-(1-6C)alkyl-piperidin-4-yl group, the (1-6C)alkyl group may be optionally substituted by, for example a (2-6C)alkanoyl group to give a 1-((2-6C)alkanoyl-(1-6C)alkyl)-piperidin-4-yl group such as 1-(acetylmethyl)piperidin-4-yl or 1-(2-acetylethyl)piperidin-4-yl. Other suitable groups that may be so formed by Q¹ include, (1-6C)alkoxycarbonyl-(1-6C)alkyl substituted heterocyclyl groups, such as 1-(methoxycarbonylmethyl)piperidin-4-yl or 1-(2-methoxycarbonylethyl)piperidin-4-yl, carbamoyl-(1-6C)alkyl substituted heterocyclyl groups such as 1-(carbamoylmethyl)piperidin-4-yl, or (1-6C)alkoxy-(1-6C)alkyl substituted heterocyclyl groups, such as 1-(2-methoxyethyl)piperidin-4-yl.

A suitable pharmaceutically-acceptable salt of a compound of the Formula I is, for example, an acid-addition salt of a compound of the Formula I, for example an acid-addition salt with an inorganic or organic acid such as hydrochloric, hydrobromic, sulphuric, trifluoroacetic, citric or maleic acid; or, for example, a salt of a compound of the Formula I which is sufficiently acidic, for example an alkali or alkaline earth metal salt such as a calcium or magnesium salt, or an ammonium salt, or a salt with an organic base such as methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

- 25 -

Particular novel compounds of the invention include, for example, quinazoline derivatives of the Formula I, or pharmaceutically-acceptable salts thereof, wherein, unless otherwise stated, each of R^1 , Z, Q^1 and Q^2 has any of the meanings defined hereinbefore or in paragraphs (a) to (zzz) hereinafter :-


- 5 (a) R^1 is hydrogen;
- (b) Z is a direct bond or is selected from O, and $N(R^2)$ wherein R^2 is hydrogen or (1-6C)alkyl;
- (c) Z is O;
- (d) Z is a direct bond or is O, Q^1 is (3-7C)cycloalkyl or heterocyclyl,
- 10 and wherein any CH_2 group within the Q^1 -Z- group optionally bears on each said CH_2 group a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,
- and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl,
- 15 hydroxy, amino, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and (2-6C)alkanoyl, or optionally bears 1 substituent selected from a group of the formula :
- $-X^1-R^4$
- 20 wherein X^1 is a direct bond or is selected from O and $N(R^5)$, wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl,
- 25 and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo or thioxo substituents;
- (e) the Q^1 -Z- group is selected from cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy, cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy and cycloheptylmethoxy,
- 30 or Z is a direct bond or is selected from O and NH and Q^1 is oxetan-3-yl, tetrahydrofuran-3-yl, 3- or 4-tetrahydropyranyl, 3- or 4-oxepanyl, 1-, 2- or 3-pyrrolidinyl, morpholino, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yl, piperidin-4-yl, 1-, 3- or 4-homopiperidinyl, piperazin-1-yl, homopiperazin-1-yl,

- azetidin-3-yl, tetrahydrothien-3-yl, 1,1-dioxotetrahydrothien-3-yl, 1-oxotetrahydrothien-3-yl, tetrahydrothiopyran-3-yl, tetrahydrothiopyran-4-yl, 1-oxotetrahydrothiopyran-3-yl, 1,1-dioxotetrahydrothiopyran-3-yl, 1-oxotetrahydrothiopyran-4-yl, 1,1-dioxotetrahydrothiopyran-4-yl, 1-, 2- or 3-pyrrolidinylmethyl, morpholinomethyl, 5 piperidinomethyl, 3- or 4-piperidinylmethyl, 1-, 3- or 4-homopiperidinylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-1-ylpropyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-2-ylpropyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-piperidin-3-ylethyl, 2-piperidin-4-ylethyl, 2-homopiperidin-1-ylethyl, 10 3-homopiperidin-1-ylpropyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, 2-homopiperazin-1-ylethyl or 3-homopiperazin-1-ylpropyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q¹-Z- group are optionally separated by the insertion into the chain of a group selected from O, NH, CONH, NHCO, CH=CH and C≡C,

- 15 and wherein any CH₂ or CH₃ group within the Q¹-Z- group optionally bears on each said CH₂ or CH₃ group a substituent selected from hydroxy, amino, methoxy, methylsulphonyl, methylamino and dimethylamino,

- and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, 20 hydroxy, amino, carbamoyl, methyl, ethyl and methoxy, or optionally bears 1 substituent selected from a group of the formula :



- wherein X¹ is a direct bond or is selected from O and NH and R⁴ is 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, aminomethyl, 2-aminoethyl, 25 3-aminopropyl, methylaminomethyl, 2-methylaminoethyl, 3-methylaminopropyl, 2-ethylaminoethyl, 3-ethylaminopropyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, acetamidomethyl, methoxycarbonylaminomethyl, ethoxycarbonylaminomethyl and tert-butoxycarbonylaminomethyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo 30 substituents;

(f) the Q¹-Z- group is selected from cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy, cyclopropylmethoxy, tetrahydrofuran-3-yloxy, tetrahydrofurfuryloxy, 3- or

- 4-tetrahydropyranyloxy, 2-tetrahydropyran-4-ylethoxy, 2-tetrahydropyran-3-ylethoxy,
3-tetrahydropyran-4-ylpropoxy, 3-tetrahydropyran-3-ylpropoxy, tetrahydrothiopyran-3-yloxy,
2-tetrahydrothiopyran-3-ylethoxy, tetrahydrothiopyran-4-yloxy,
2-tetrahydrothiopyran-4-ylethoxy, 1-oxotetrahydrothiopyran-3-yloxy,
5 2-(1-oxotetrahydrothiopyran-3-yl)ethoxy, 1,1-dioxotetrahydrothiopyran-3-yloxy,
2-(1,1-dioxotetrahydrothiopyran-3-yl)ethoxy, 1-oxotetrahydrothiopyran-4-yloxy,
2-(1-oxotetrahydrothiopyran-4-yl)ethoxy, 1,1-dioxotetrahydrothiopyran-4-yloxy,
2-(1,1-dioxotetrahydrothiopyran-4-yl)ethoxy, 3-tetrahydrothiopyran-3-ylpropoxy,
3-(1,1-dioxotetrahydrothiopyran-3-yl)propoxy, 3-(1-oxotetrahydrothiopyran-3-yl)propoxy,
10 3-tetrahydrothiopyran-4-ylpropoxy, 3-(1-oxotetrahydrothiopyran-4-yl)propoxy,
3-(1,1-dioxotetrahydrothiopyran-4-yl)propoxy, tetrahydrothien-3-yloxy,
1,1-dioxotetrahydrothien-3-yloxy, 1-oxotetrahydrothien-3-yloxy,
2-tetrahydrothien-3-ylethoxy, 2-(1,1-dioxotetrahydrothien-3-yl)ethoxy,
2-(1-oxotetrahydrothien-3-yl)ethoxy, 3-tetrahydrothien-3-ylpropoxy,
15 3-(1,1-dioxotetrahydrothien-3-yl)propoxy, 3-(1-oxotetrahydrothien-3-yl)propoxy,
azetidin-3-yloxy, 2-azetidin-3-ylethoxy, 3-azetidin-3-ylpropoxy, pyrrolidin-1-yl, morpholino,
piperidino, piperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy,
pyrrolidin-3-yloxy, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy,
3-pyrrolidin-2-ylpropoxy, 2-morpholinoethoxy, 3-morpholinopropoxy,
20 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethoxy,
3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propoxy, 2-piperidinoethoxy,
3-piperidinopropoxy, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylmethoxy,
2-piperidin-3-ylethoxy, piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy,
homopiperidin-3-yloxy, homopiperidin-4-yloxy, homopiperidin-3-ylmethoxy,
25 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-piperazin-1-ylethoxy,
3-piperazin-1-ylpropoxy, 2-homopiperazin-1-ylethoxy, 3-homopiperazin-1-ylpropoxy,
2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-3-ylamino,
pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino,
2-morpholinoethylamino, 3-morpholinopropylamino,
30 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethylamino,
3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propylamino, 2-piperidinoethylamino,
3-piperidinopropylamino, piperidin-3-ylamino, piperidin-4-ylamino,
piperidin-3-ylmethylamino, 2-piperidin-3-ylethylamino, piperidin-4-ylmethylamino,

2-piperidin-4-ylethylamino, homopiperidin-3-ylamino, homopiperidin-4-ylamino,
 homopiperidin-3-ylmethylamino, 2-homopiperidin-1-ylethylamino,
 3-homopiperidin-1-ylpropylamino, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino,
 2-homopiperazin-1-ylethylamino or 3-homopiperazin-1-ylpropylamino,

5 and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q¹-Z-
 group are optionally separated by the insertion into the chain of a group selected from O, NH,
 CH=CH and C≡C,

and wherein any CH₂ or CH₃ group within the Q¹-Z- group optionally bears on each
 said CH₂ or CH₃ group a substituent selected from hydroxy, amino, methoxy,
 10 methylsulphonyl, methylamino and dimethylamino,

and wherein any heterocyclyl group within the Q¹-Z group optionally bears
 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro,
 trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, allyl, 2-propynyl, acetyl,
 acetylmethyl, methoxycarbonyl, methoxycarbonylmethyl, methylsulphonyl, 2-methoxyethyl,
 15 carbamoyl, N-methylcarbamoyl, N,N-di-methylcarbamoyl, carbamoylmethyl,
N-methylcarbamoylmethyl and N,N-di-methylcarbamoylmethyl,

and wherein any heterocyclyl group within the Q¹-Z group optionally bears 1 or 2 oxo
 substituents;

(g) the Q¹-Z- group is selected from cyclopentyloxy, cyclohexyloxy,
 20 tetrahydrofuran-3-yloxy, tetrahydropyran-3-yloxy, tetrahydropyran-4-yloxy,
 tetrahydrothiopyran-3-yloxy, tetrahydrothiopyran-4-yloxy,
 1,1-dioxotetrahydrothiopyran-3-yloxy, 1,1-dioxotetrahydrothiopyran-4-yloxy,
 1-oxotetrahydrothiopyran-3-yloxy, 1-oxotetrahydrothiopyran-4-yloxy,
 tetrahydrothien-3-yloxy, 1,1-dioxodotetrahydrothien-3-yloxy, 1-oxotetrahydrothien-3-yloxy,
 25 pyrrolidin-3-yloxy, pyrrolidin-2-yloxy, piperidin-3-yloxy, piperidin-4-yloxy,
 homopiperidin-3-yloxy, homopiperidin-4-yloxy and azetidin-3-yloxy,

and wherein any azetidiny, pyrrolidinyl, piperidinyl or homopiperidinyl group within
 the Q¹-Z- group is optionally N- substituted by a substituent, T¹, selected from (1-4C)alkyl,
 (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, carbamoyl,
 30 N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl and (1-4C)alkylsulphonyl,

and wherein adjacent carbon atoms in any (2-4C)alkylene chain within a N-substituent, T¹, are optionally separated by the insertion into the chain of a group selected from O, NH and CO,

and wherein any CH₂ or CH₃ group within the N-substituent, T¹, optionally bears on
 5 each said CH₂ or CH₃ group a substituent selected from hydroxy, amino, methylamino, di-methylamino, ethylamino, diethylamino, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, acetyl, methoxycarbonyl and ethoxycarbonyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

10 (h) the Q¹-Z- group is selected from cyclopentyloxy, tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, tetrahydrothiopyran-4-yloxy, 1,1-dioxotetrahydrothiopyran-4-yloxy, 1-oxotetrahydrothiopyran-4-yloxy, tetrahydrothien-3-yloxy, 1,1-dioxodotetrahydrothien-3-yloxy, 1-oxotetrahydrothien-3-yloxy, pyrrolidin-3-yloxy, pyrrolidin-2-yloxy, piperidin-3-yloxy, piperidin-4-yloxy, homopiperidin-3-yloxy,
 15 homopiperidin-4-yloxy and azetidin-3-yloxy,

and wherein the azetidiny, pyrrolidinyl, piperidinyl or homopiperidinyl group within the Q¹-Z- group is optionally N-substituted by a substituent selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, allyl, 2-propynyl, acetyl, propionyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, methylsulphonyl,
 20 ethylsulphonyl, 2-methoxyethyl, carbamoylmethyl, N-methylcarbamoylmethyl, N,N-di-methylcarbamoylmethyl, 2-carbamoylethyl, 2-(N-methylcarbamoyl)ethyl, 2-(N,N-di-methylcarbamoyl)ethyl, acetylmethyl, 2-acetylethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo
 25 substituents;

(i) the Q¹-Z- group is piperidin-4-yloxy which group is optionally N-substituted by a substituent selected from methyl, ethyl, n-propyl, iso-propyl, allyl, 2-propynyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, methylsulphonyl, 2-methoxyethyl, carbamoylmethyl, N-methylcarbamoylmethyl,
 30 N,N-di-methylcarbamoylmethyl, acetylmethyl, 2-acetylethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

(j) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $CH(OR^6)$, $CON(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or (1-6C)alkyl, and Q^3 is aryl, or heteroaryl,

10 and any aryl or heteroaryl in the group X^2-Q^3 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

15 (k) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is aryl, or heteroaryl,

20 and any aryl or heteroaryl in the group X^2-Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(l) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl and (2-8C)alkynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is aryl, which optionally bears 1 or 2 substituents, which may be the same or different, selected from

halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(m) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both
5 hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl and (2-8C)alkynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is heteroaryl,
10 which optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(n) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both
15 hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, fluoro, chloro, bromo, trifluoromethyl, methyl, ethyl, vinyl, allyl and ethynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is a phenyl or
20 naphthyl, group which is optionally substituted with 1 or 2 substituents selected from fluoro, chloro, bromo, trifluoromethyl, nitro, methyl, ethyl, isopropyl, methoxy, vinyl, ethynyl and cyano;

(o) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

25 G^1 is hydrogen, G^2 is selected from hydrogen, fluoro, chloro, bromo, trifluoromethyl, methyl, ethyl, vinyl, allyl and ethynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$ and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is a phenyl group which is optionally substituted with 1 or 2 substituents selected from fluoro, chloro, bromo,
30 trifluoromethyl, nitro, methyl, ethyl, isopropyl, methoxy, vinyl, ethynyl and cyano;

(p) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

- 32 -

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, (1-6C)alkyl, (2-8C)alkenyl and (2-8C)alkynyl,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$ and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is selected from a furyl,

5 thienyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, 1,2,4-triazolyl, 1,2,5-thiadiazolyl, pyridyl, pyrimidinyl, 1,3-benzodioxolyl, quinolinyl, isoquinolinyl and quinazolinyl group which is optionally substituted with one or two substituents selected from fluoro, chloro, bromo, trifluoromethyl, nitro, methyl, ethyl, isopropyl, vinyl, ethynyl, methoxy and cyano;

(q) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both
10 hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, fluoro, chloro, bromo, trifluoromethyl, methyl, ethyl, vinyl, allyl, ethynyl and cyano,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $OC(R^6)_2$, $SC(R^6)_2$, CO and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or methyl, and Q^3 is selected from a
15 2-furyl, 2- or 3-thienyl, 2-,4- or 5-oxazolyl, 3-,4- or 5-isoxazolyl, 2-,4-or 5-1H-imidazolyl, 2-,4-or 5-thiazolyl, 3- (1H-1,2,4-triazolyl), 3- (1,2,5-thiadiazolyl), 2- 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1,3-benzodioxol-4-yl, 1,3-benzodioxol-5-yl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl and 2-, 3-, 4-, 5-, 6-, 7- or 8-quinazolinyl group, which is optionally substituted with one or two substituents selected from nitro, fluoro,
20 chloro, bromo, methyl, ethyl, methoxy, trifluoromethyl, ethynyl, and cyano;

(r) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 is hydrogen and G^2 is selected from hydrogen, fluoro, chloro, bromo, methyl, and ethynyl,

25 X^2 is a direct bond or is selected from O, S, NH, $N(CH_3)$, OCH_2 , CO and $NHCH_2$, and Q^3 is phenyl optionally substituted with 1 or 2 substituents selected from fluoro, chloro, bromo, nitro, methyl, methoxy, ethyl, ethynyl, cyano and nitro, or Q^3 is a heteroaryl moiety selected from 2-1H-imidazolyl, 4-(1,3-thiazolyl), 2-thienyl, 3-(1,2,5-thiadiazolyl), 3-isoxazolyl 2-, 3- or 4-pyridyl, 2-pyrimidinyl, 1,3-benzodioxol-5-yl
30 and 8-quinolinyl, which moiety is optionally substituted with one or two substituents selected from fluoro, chloro, bromo, trifluoromethyl, methyl, methoxy, ethynyl and cyano;

(s) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 and G^2 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

5 X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

and the nitrogen containing heterocyclyl group in the group X^2 - Q^3 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, 10 (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(t) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 and G^2 each, independently, is selected from hydrogen, halogeno, trifluoromethyl, 15 cyano, hydroxy, amino, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

and the nitrogen containing heterocyclyl group in the group X^2 - Q^3 optionally bears 1 20 or 2 substituents, which may be the same or different, selected from halogeno, cyano, hydroxy, amino, (1-6C)alkyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino, and Q^3 optionally bears 1 or 2 oxo substituents;

(u) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

25 G^1 and G^2 each, independently, is selected from hydrogen, fluoro, chloro, bromo, cyano, hydroxy, methyl, ethyl and ethynyl,

X^2 is CO and Q^3 is selected from pyrrolidin-1-yl, piperidino, homopiperidino, morpholino, piperazin-1-yl, homopiperazin-1-yl, decahydroquinolin-1-yl, and decahydroisoquinolin-2-yl,

30 and wherein Q^3 optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, cyano, hydroxy, methyl and ethyl,

and wherein Q^3 optionally bears 1 or 2 oxo substituents;

(v) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 is selected from fluoro, chloro, bromo and ethynyl and G^2 is hydrogen,

X^2 is CO and Q^3 is selected from piperidino, homopiperidino,

5 decahydroquinolin-1-yl, and decahydroisoquinolin-2-yl,

and wherein Q^3 optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, cyano, hydroxy, methyl and ethyl,

and wherein Q^3 optionally bears 1 or 2 oxo substituents;

(w) Q^2 is a group of the formula Ib or Ic as hereinbefore defined wherein G^6 and G^7 are
10 both hydrogen,

G^1 , G^3 , G^4 and G^5 each, independently, is selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

15 X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,

and Q^4 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy,
20 (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino

(x) Q^2 is a group of the formula Ib or Ic as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^3 is hydrogen,

25 G^1 is selected from hydrogen, fluoro, chloro, bromo, trifluoromethyl, cyano, hydroxy, amino, methyl, ethyl, vinyl, ethynyl, methylamino and di-methylamino,

G^4 and G^5 each, independently, is selected from hydrogen, fluoro, chloro, bromo, cyano, hydroxy, amino, methyl, ethyl, vinyl, ethynyl, methylamino and di-methylamino,

X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is,
30 independently, hydrogen or methyl, and Q^4 is selected from a phenyl, 2-furyl, 3-furyl, 2-(1,3-oxazolyl), 4-(1,3-oxazolyl), 3-, 4- or 5-isoxazolyl, 2-imidazolyl, 4-imidazolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5- pyrimidinyl, 1,2,4-triazol-3-yl, 2-thienyl, 3-thienyl, 2-(1,3-thiazolyl), 4-(1,3-thiazolyl), 3-, 4- or 5- isothiazolyl and 1,2,5-thiadiazol-3-yl group which group

optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, methoxy, vinyl, ethynyl and trifluoromethyl;

(y) Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^6 and G^7 are hydrogen,

5 G^4 is selected from hydrogen, fluoro, chloro, bromo, cyano, methyl and ethynyl,
 X^3 is selected from SO_2 and CH_2 , and Q^4 is selected from phenyl and 2-, 3- or 4-pyridyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, nitro, cyano and methyl;

(z) Q^2 is a group of the formula Id or Ie as hereinbefore defined wherein G^6 and G^7 are
 10 both hydrogen,

G^1 , G^4 and G^5 each, independently, is selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

15 X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,

and Q^4 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy,
 20 (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

(aa) Q^2 is a group of the formula Id or Ie as hereinbefore defined wherein G^6 and G^7 are both hydrogen,

G^1 is selected from hydrogen, fluoro, chloro, bromo, trifluoromethyl, cyano, hydroxy,
 25 amino, methyl, ethyl, vinyl, ethynyl, methylamino and di-methylamino,

G^4 and G^5 each, independently, is selected from hydrogen, fluoro, chloro, bromo, cyano, hydroxy, amino, methyl, ethyl, vinyl, ethynyl, methylamino and di-methylamino,

X^3 is a direct bond or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently hydrogen or methyl, and Q^4 is selected from a phenyl, 2-furyl, 3-furyl, 2-(1,3-oxazolyl), 4-(1,3-oxazolyl), 3-, 4- or 5-isoxazolyl, 2-imidazolyl, 4-imidazolyl, 2-, 3- or
 30 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1,2,4-triazol-3-yl, 2-thienyl, 3-thienyl, 2-(1,3-thiazolyl), 4-(1,3-thiazolyl), 3-, 4- or 5-isothiazolyl and 1,2,5-thiadiazol-3-yl group which group

- 36 -

optionally bears 1 or 2 substituents, which may be the same or different, selected fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, methoxy, vinyl, ethynyl and trifluoromethyl;


(bb) Q^2 is a group of the formula Id as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

5 G^4 is selected from hydrogen, fluoro, chloro, bromo, cyano, methyl and ethynyl,

X^3 is selected from SO_2 and CH_2 , and Q^4 is selected from phenyl and 2-, 3- or 4-pyridyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, nitro, cyano and methyl;

(cc) Q^1 is selected from (3-7C)cycloalkyl and a 4, 5, 6 or 7 membered heterocyclyl ring
10 linked to Z by a carbon atom,

and wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from formyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, aminoalkanoyl, (1-4C)alkoxycarbonyl, carbamoyl, sulphamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, N-(1-4C)alkylsulphamoyl, N,N-di-(1-4C)alkylsulphamoyl and
15 (1-4C)alkylsulphonyl, or from a group of the formula :




wherein X^1 is a direct bond, and R^4 is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-6C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,
20 N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl,

and wherein any CH or CH_2 group within a (3-7C)cycloalkyl or heterocyclyl group within Q^1 group optionally bears 1 substituent on each said CH group or 1 or 2 substituents on each said CH_2 group, which may be the same or different, selected from halogeno and
25 (1-6C)alkyl, or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,
30 N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,

and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo substituents,


- 37 -

- (dd) Z is O and Q¹ is selected from a 4, 5 or 6 membered heterocyclyl ring containing at least 1 nitrogen atom, said ring being linked to Z by a carbon atom, and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from formyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, carbamoyl, N-(1-4C)alkylcarbamoyl and N,N-di-(1-4C)alkylcarbamoyl, or from a group of the formula:



- wherein X¹ is a direct bond, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-6C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl or (2-4C)alkanoyl-(1-4C)alkyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

- 15 (ee) Z is O and Q¹ is selected from azetidin-3-yl, pyrrolidin-3-yl, morpholin-3-yl, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl and tetrahydro-4H-1,4-thiazin-3-yl (preferably pyrrolidin-3-yl, piperidin-3-yl or piperidin-4-yl), and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from formyl, cyano, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, carbamoyl, sulphamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, N-(1-4C)alkylsulphamoyl, N,N-di-(1-4C)alkylsulphamoyl, (1-4C)alkylsulphonyl and from a group of the formula :



- wherein X¹ is a direct bond, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

- 30 (ff) Z is O and Q¹ is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl, and wherein any NH group within a pyrrolidinyl or piperidinyl group in Q¹ optionally bears a substituent selected from methyl, allyl, acetyl, carbamoyl, methoxymethyl, 2-methoxyethyl,

- 38 -

carbamoylmethyl, N-methylcarbamoylmethyl, N,N-di-methylcarbamoylmethyl, 2-carbamoylethyl, 2-(N-methylcarbamoyl)ethyl, 2-(N,N-di-methylcarbamoyl)ethyl, acetylmethyl; 2-acetylethyl, cyanomethyl, 2-cyanoethyl, fluoromethyl, chloromethyl, 2-fluoroethyl and 2-chloroethyl,

5 and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 oxo substituent;

(gg) Z is O and Q¹ is selected from a 5 or 6 membered heterocyclyl ring containing at least 1 heteroatom selected from O and S and no nitrogen heteroatoms, and wherein said heterocyclyl ring is linked to Z by a carbon atom,

10 and wherein said 5 or 6 membered heterocyclyl ring optionally bears 1, 2 or 3 substituents selected from halogeno, (1-6C)alkyl, hydroxy, amino, carboxy, (1-6C)alkoxy and (1-6C)alkylthio

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

15 (hh) Z is O and Q¹ is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl, tetrahydropyran-4-yl, tetrahydrothiopyran-4-yl, tetrahydrothiopyran-3-yl, 1,1-dioxotetrahydrothiopyran-3-yl, 1,1-dioxotetrahydrothiopyran-4-yl, 1-oxotetrahydrothiopyran-3-yl, 1-oxotetrahydrothiopyran-4-yl, tetrahydrothien-3-yl, 1,1-dioxodotetrahydrothien-3-yl, 1-oxotetrahydrothien-3-yl,

20 and wherein any 5 or 6 membered heterocyclyl ring within Q¹ optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, (1-4C)alkyl, hydroxy, amino, carboxy, (1-4C)alkoxy and (1-4C)alkylthio, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;


(ii) Z is O and Q¹ is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl and 25 tetrahydropyran-4-yl,

and wherein Q¹ optionally bears an oxo substituent;


(jj) Q¹ is selected from (3-7C)cycloalkyl and (3-7C)cycloalkyl-(1-6C)alkyl, which optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, 30 (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,

- 39 -

N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,
N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and
N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, from a group of the formula :




- 5 wherein X^1 is a direct bond or is selected from O and N(R^5), wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,
 10 (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, and from a group of the formula:



- wherein X^5 is a direct bond or is selected from O, CO and N(R^{10}), wherein R^{10} is hydrogen or (1-6C)alkyl, and Q^6 is a nitrogen containing heterocyclyl or nitrogen containing
 15 heterocyclyl-(1-6C)alkyl, and wherein any heterocyclyl group in Q^6 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, hydroxy, cyano, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (1-6C)alkoxycarbonyl, (1-6C)alkylsulphonyl carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, halogeno-(1-6C)alkyl,
 20 hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, and wherein any heterocyclyl group in Q^1 optionally bears 1 or 2 oxo substituents;

- 25 (kk) Q^1 is selected from (3-7C)cycloalkyl and (3-7C)cycloalkyl-(1-6C)alkyl, which is substituted by 1 substituent selected from, amino, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, and from a group of the formula:




- 30 wherein X^5 is a direct bond or is selected from O, CO and N(R^{10}), wherein R^{10} is hydrogen or (1-6C)alkyl, and Q^6 is nitrogen containing heterocyclyl or nitrogen containing heterocyclyl-(1-6C)alkyl, and wherein any heterocyclyl group in Q^6 optionally bears 1 or 2

substituents, which may be the same or different, selected from halogeno, hydroxy, cyano, formyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, (1-4C)alkylsulphonyl carbamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, halogeno-(1-4C)alkyl, 5 hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, and wherein the (3-7C)cycloalkyl group in Q¹ optionally bears 1 or 2 further substituents, 10 which may be the same or different, selected from halogeno, cyano, hydroxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl and (1-4C)alkoxy

and wherein any heterocyclyl group in Q¹ optionally bears 1 or 2 oxo substituents;

(II) Z is O and Q¹ is (3-7C)cycloalkyl substituted by 1 substituent selected from, amino, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, 15 di-[(1-4C)alkyl]amino-(1-4C)alkyl, and from a group of the formula:



wherein X⁵ is a direct bond, and Q⁶ is a 5, 6 or 7 membered nitrogen containing heterocyclyl or a 5, 6, or 7 membered nitrogen containing heterocyclyl-(1-6C)alkyl, and wherein Q⁶ optionally bears 1 or 2 substituents, which may be the same or different, selected from 20 halogeno, hydroxy, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, (1-4C)alkylsulphonyl carbamoyl, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl, and wherein the (3-7C)cycloalkyl group in Q¹ optionally bears 1 or 2 further substituents, which may be the same or different, selected from fluoro, chloro, cyano, hydroxy and 25 (1-4C)alkyl,

and wherein any heterocyclyl group in Q¹ optionally bears 1 or 2 oxo substituents;

(mm) Z is O and Q¹ is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cycloheptylmethyl, 2-cyclopropylethyl, 2-cyclobutylethyl, 2-cyclopentylethyl, 2-cyclohexylethyl, 2-cycloheptylethyl, 3-cyclopentylpropyl, 3-cyclohexylpropyl and 2-cyclohexylpropyl, which is substituted by a substituent selected from amino, methylamino, ethylamino, propylamino, di-methylamino, di-ethylamino, aminomethyl, 2-aminoethyl, 3-aminopropyl, 2-aminopropyl, methylaminomethyl, ethylaminomethyl 2-methylaminoethyl, 2-

- ethylaminoethyl, 3-methylaminopropyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 2-dimethylaminopropyl, diethylaminomethyl, 2-diethylaminoethyl, 3-diethylaminopropyl, ethylmethylaminomethyl, 2-ethylmethylaminoethyl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, 2-pyrrolin-1-yl, 2-pyrrolin-2-yl, 2-pyrrolin-3-yl, 2-pyrrolin-4-yl, 2-pyrrolin-5-yl, 3-pyrrolin-1-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl, 3-pyrrolin-4-yl, 3-pyrrolin-5-yl, morpholino, morpholin-2-yl, morpholin-3-yl, piperidino, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, piperazin-1-yl, piperazin-2-yl, piperazin-3-yl, tetrahydro-4H-1,4-thiazin-2-yl, tetrahydro-4H-1,4-thiazin-3-yl, tetrahydro-4H-1,4-thiazin-4-yl, homopiperidin-1-yl and homopiperazin-1-yl,
- 10 and where any heterocyclyl ring in Q¹ optionally bears 1 or 2 substituents selected from fluoro, chloro, hydroxy, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, (1-4C)alkylsulphonyl carbamoyl, N-(1-4C)alkylcarbamoyl and N,N-di-[(1-4C)alkyl]carbamoyl, and wherein the (3-7C)cycloalkyl group in Q¹ optionally bears 1 or 2 further substituents,
- 15 which may be the same or different, selected from fluoro, chloro, hydroxy, methyl and ethyl, and wherein any heterocyclyl group in Q¹ optionally bears 1 or 2 oxo substituents;
- (nn) Z is O and Q¹ is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, which is substituted by a substituent selected from amino, methylamino, ethylamino, propylamino, di-methylamino di-ethylamino, aminomethyl, 2-aminoethyl, 3-aminopropyl, 2-aminopropyl, methylaminomethyl, ethylaminomethyl 2-methylaminoethyl, 2-ethylaminoethyl, 3-methylaminopropyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 2-dimethylaminopropyl, diethylaminomethyl, 2-diethylaminoethyl, 3-diethylaminopropyl, ethylmethylaminomethyl and 2-ethylmethylaminoethyl,
- 20 aminopropyl, 2-aminopropyl, methylaminomethyl, ethylaminomethyl 2-methylaminoethyl, 2-ethylaminoethyl, 3-methylaminopropyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 2-dimethylaminopropyl, diethylaminomethyl, 2-diethylaminoethyl, 3-diethylaminopropyl, ethylmethylaminomethyl and 2-ethylmethylaminoethyl,
- (oo) Z is O and Q¹ is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, which is substituted by a substituent selected from pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, morpholino, morpholin-2-yl, morpholin-3-yl, piperidino, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, piperazin-1-yl, piperazin-2-yl, piperazin-3-yl, tetrahydro-4H-1,4-thiazin-2-yl, tetrahydro-4H-1,4-thiazin-3-yl, tetrahydro-4H-1,4-thiazin-4-yl, homopiperidin-1-yl and homopiperazin-1-yl, and wherein any heterocyclyl ring in Q¹
- 25 optionally bears 1 or 2 substituents selected from fluoro, chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, acetyl, propionyl, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, carbamoyl,
- 30

N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-(methyl)carbamoyl and
N,N-di-(ethyl)carbamoyl,

and wherein any cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl group in Q¹ is optionally further substituted by 1 or 2 substituents, which may be the same or different,

5 selected from fluoro, chloro and methyl,

and wherein any heterocyclyl group in Q¹ optionally bears an oxo substituent;

(pp) Z is O and Q¹ is selected from cyclopentyl and cyclohexyl, which is substituted by a substituent selected from pyrrolidin-1-yl, morpholino, piperidino and piperazin-1-yl, and

wherein any pyrrolidinyl, morpholino, piperidino or piperazinyl group in Q¹ optionally bears 1

10 or 2 substituents selected from fluoro, chloro, hydroxy, methyl, methoxy, amino,

methylamino, ethylamino, dimethylamino, diethylamino, acetyl, methoxycarbonyl,

methylsulphonyl, ethylsulphonyl, carbamoyl, N-methylcarbamoyl and

N,N-di-(methyl)carbamoyl,

and wherein any cyclopentyl or cyclohexyl group in Q¹ is optionally further substituted by 1 or
15 2 substituents, which may be the same or different, selected from fluoro and methyl,

and wherein any heterocyclyl group in Q¹ optionally bears an oxo substituent;

(qq) Z is O and Q¹ is cyclohexyl, which is substituted (conveniently at the 4-position) by a substituent selected from pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, piperidino,

piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, piperazin-1-yl, piperazin-2-yl and piperazin-3-yl

20 and wherein any pyrrolidinyl, piperidino or piperazinyl group in Q¹ optionally bears 1 or 2

substituents selected from fluoro, chloro, hydroxy, methyl, methoxy, amino, methylamino,

ethylamino, dimethylamino, diethylamino, acetyl, methoxycarbonyl, methylsulphonyl,

ethylsulphonyl, carbamoyl, N-methylcarbamoyl and N,N-di-(methyl)carbamoyl,

and wherein any heterocyclyl group in Q¹ optionally bears an oxo substituent;

25 (rr) Z is O and Q¹ is cyclohexyl, which is substituted (conveniently at the 4-position) by a

substituent selected from amino, methylamino, ethylamino, di-methylamino di-ethylamino,

aminomethyl, 2-aminoethyl, methylaminomethyl, ethylaminomethyl 2-methylaminoethyl,

dimethylaminomethyl and 2-dimethylaminoethyl;

(ss) Z is O and Q¹ is 4-(piperazin-1-yl)cyclohexyl, wherein the piperazin-1-yl group is

30 optionally substituted at the 4-position by a substituent selected from (1-3C)alkyl,

(2-4C)alkanoyl, (1-3C)alkoxycarbonyl, (1-3C)alkylsulphonyl, carbamoyl,

N-(1-3C)alkylcarbamoyl and N,N-di-[(1-3C)alkyl]carbamoyl,

and wherein the piperazin-1-yl group in Q¹ optionally bears an oxo substituent;

- (tt) Z is O and Q¹ is 4-(piperazin-1-yl)cyclohexyl, wherein the piperazin-1-yl group is optionally substituted at the 4-position by (1-3C)alkyl, for example methyl;
- (uu) Z is O and Q¹ is piperidin-4-yl optionally substituted at the 1 position by a substituent selected from methyl, allyl, 2-methoxyethyl and carbamoylmethyl,
- 5 and wherein the piperidin-4-yl group optionally bears an oxo substituent;
- (vv) Q¹Z is 1-methylpiperidin-4-yloxy;
- (ww) Q¹Z is selected from pyrrolidin-1-ylmethoxy, 2-pyrrolidin-1-yloxy, 3-pyrrolidin-1-ylpropoxy, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-yloxy, 3-pyrrolidin-2-ylpropoxy, piperidinomethoxy, 2-piperidinoethoxy,
- 10 3-piperidinopropoxy, piperidin-2-ylmethoxy, 2-piperidin-2-yloxy, 3-piperidin-2-ylpropoxy, piperidin-3-ylmethoxy, 2-piperidin-3-yloxy, 3-piperidin-3-yl propoxy, piperidin-4-ylmethoxy, 2-piperidin-4-yloxy, 3-piperidin-4-ylpropoxy, homopiperidin-1-ylmethoxy, 2-homopiperidin-1-yloxy, 3-homopiperidin-1-ylpropoxy, homopiperidin-2-ylmethoxy,
- 15 2-homopiperidin-2-yloxy, 3-homopiperidin-2-ylpropoxy, homopiperidin-3-ylmethoxy, 2-homopiperidin-3-yloxy, 3-homopiperidin-3-ylpropoxy, piperazin-1-ylmethoxy, 2-piperazin-1-yloxy, 3-piperazin-1-ylpropoxy, piperazin-2-ylmethoxy, 2-piperazin-2-yloxy, 3-piperazin-2-ylpropoxy, homopiperazin-1-ylmethoxy, 2-homopiperazin-1-yloxy, 3-homopiperazin-1-ylpropoxy, homopiperazin-2-ylmethoxy,
- 20 2-homopiperazin-2-yloxy, 3-homopiperazin-2-ylpropoxy, homopiperazin-3-ylmethoxy, 2-homopiperazin-3-yloxy, 3-homopiperazin-3-ylpropoxy,
- and wherein any heterocyclyl group within the Q¹-Z group optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, cyano, formyl, trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, allyl, ethynyl, 2-propynyl,
- 25 acetyl, acetylmethyl, fluoromethyl, 2-fluoroethyl, chloromethyl, 2-chloroethyl, methoxycarbonyl, methoxycarbonylmethyl, methylsulphonyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,Ndi-methylcarbamoylmethyl,
- and wherein any heterocyclyl group within the Q¹-Z group optionally bears 1 or 2 oxo
- 30 substituents;
- (xx) Q¹Z is selected from pyrrolidin-1-ylmethoxy, pyrrolidin-2-ylmethoxy, piperidinomethoxy, piperidin-2-ylmethoxy, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy and piperazin-1-ylmethoxy, and wherein any heterocyclyl group

- within the Q¹-Z group optionally bears on an NH a substituent selected from methyl, ethyl, formyl, ethynyl, 2-propynyl, acetyl, acetylmethyl, fluoromethyl, 2-fluoroethyl, chloromethyl, 2-chloroethyl, cyanomethyl, methoxycarbonyl, methoxycarbonylmethyl, methylsulphonyl, methoxymethyl, 2-methoxyethyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-di-methylcarbamoylmethyl,
- and wherein any heterocyclyl group within the Q¹-Z group optionally bears an oxo substituent;
- (yy) Q¹Z is selected from pyrrolidin-1-ylmethoxy, pyrrolidin-2-ylmethoxy, piperidinomethoxy, piperidin-2-ylmethoxy, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy and piperazin-1-ylmethoxy, and wherein any NH group within a heterocyclyl group on Q¹-Z is substituted by a substituent selected from methyl and ethyl,
- (zz) G⁶ and G⁷ are hydrogen;
- (aaa) G⁶ is hydrogen and G⁷ is fluorine, or G⁷ is hydrogen and G⁶ is fluorine;
- (bbb) X² is selected from S, N(R⁶)C(R⁶)₂ and OC(R⁶)₂, wherein each R⁶ is, independently, hydrogen or (1-3C)alkyl;
- (ccc) X² is selected from S and OC(R⁶)₂, wherein each R⁶ is, independently, hydrogen or methyl;
- (ddd) Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,
- G² is selected from hydrogen, halogeno, (1-6C)alkyl, (1-6C)alkoxy, (2-8C)alkenyl and (2-8C)alkynyl,
- X² is selected from S and OC(R⁶)₂, wherein R⁶ is, independently, hydrogen or methyl, and Q³ is selected from phenyl, 2- or 3-furyl, 2- or 3-thienyl, 2-,4- or 5-(1,3-oxazolyl), 3-,4- or 5-isoxazolyl, 2-,4-or 5-1H-imidazolyl, 2-,4-or 5-thiazolyl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 1H-1,3,4-triazol-2-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-oxadiazol-5-yl, 3-, 4- or 5-1H-pyrazolyl, 2- 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, 1,2-benzisoxazol-3-yl, 1,3-benzodioxol-4-yl, 1,3-benzodioxol-5-yl, 2-imidazo[1,2-a]pyridyl, 3-benzo[d]isoxazolyl and 8-quinolinyl,
- and wherein Q³ optionally bears 1 or 2 substituents, which may be the same or different,
- selected from halogeno, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino,

(1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and

5 N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl,

10 di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,

N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl;

(eee) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

15 G^2 is selected from hydrogen, fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X^2 is $OC(R^6)_2$, R^6 is, independently, hydrogen or methyl, and Q^3 is phenyl, and

wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy,


20 formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl,

(1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl,

N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl,

(2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino,

25 N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl, hydroxy-(1-4C)alkyl,

30 (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl,

(1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,

N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl,

(2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;

(fff) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from hydrogen, fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

5 X^2 is S, and Q^3 is phenyl, and

wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl,

10 (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :

15 $-X^4-R^8$

wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,

20 N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;

(ggg) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,


25 X^2 is $OC(R^6)_2$, R^6 is hydrogen, and Q^3 is phenyl, and wherein Q^3 is optionally substituted by 1 or 2 substituents selected from fluoro and cyano (for example Q^3 is selected from 2-fluorophenyl, 3-fluorophenyl, 2-cyanophenyl, 3-cyanophenyl, 2,5-difluorophenyl and 2,6-difluorophenyl);

(hhh) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are
30 hydrogen,

G^2 is selected from hydrogen, halogeno, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X^2 is S, and

- Q^3 is selected from 2-thienyl, 3-thienyl, 1,3-oxazol-4-yl, 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 1H-1,3,4-triazol-2-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 3-1H-pyrazolyl, 2-pyridyl, 3-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 2-pyrazine and 8-quinoliny, 5
- wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, 10
- N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



- 15 wherein X^4 is a direct bond or is selected from O and N(R^9), wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, 20
- (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;
- (iii) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from hydrogen, halogeno, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X^2 is $OC(R^6)_2$, R^6 is, independently, hydrogen or methyl, and

- 25 Q^3 is selected from 2-thienyl, 3-thienyl, 1,3-oxazol-4-yl, 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 5-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-1H-pyrazolyl, 1H-1,2,4-triazol-3-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-pyridyl, 3-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 2-pyrazine and 1H-1,3,4-triazolyl-2-yl, and wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, 30
- selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl,

N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and N(R^9), wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,

10 N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;

(iii) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from hydrogen, fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and

15 (2-4C)alkynyl,

X^2 is selected from S and OC(R^6)₂, R^6 is hydrogen,

Q^3 is selected from 2-thienyl, 3-thienyl, 1,3-oxazol-4-yl, 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-, 4- or 5-1H-pyrazolyl, 1H-1,3,4-triazol-2-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-

20 pyridyl, 3-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl and 2-pyrazine, and

wherein any NH group within Q^3 is substituted by a substituent selected from (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, formyl, carbamoyl, sulphamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, (1-4C)alkylsulphonyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl

25 amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl,

N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl and

(1-4C)alkoxycarbonyl-(1-4C)alkyl, and wherein Q^3 optionally bears 1 substituent on a ring carbon atom selected from (1-4C)alkyl, amino, (1-4C)alkylamino and di-(1-4C)alkylamino;

30 (kkk) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from hydrogen, fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and

(2-4C)alkynyl,

- X^2 is selected from S and $OC(R^6)_2$, R^6 is, independently, hydrogen or methyl, and Q^3 is selected from 1,2-benzisoxazol-3-yl, 1,3-benzodioxol-4-yl, 1,3-benzodioxol-5-yl, 2-imidazo[1,2-a]pyridyl, 3-benzo[d]isoxazolyl and 8-quinolinyl, and wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different,
- 5 selected from halogeno, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl,
- 10 (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



- 15 wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,
- 20 (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl;
- (III) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X^2 is S,

- 25 Q^3 is selected from 2-1H-imidazolyl, 2-thiazolyl and 2-pyridyl; and Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, cyano, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, methoxycarbonyl, ethoxycarbonyl, formyl, carbamoyl, mercapto, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-methylcarbamoyl, N,N-di-ethylcarbamoyl, methylsulphonyl,
- 30 hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methy-amino)ethyl, di-methylaminomethyl, 2-(dimethylamino)ethyl, 2-(diethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-methylcarbamoylmethyl, N-ethylcarbamoylmethyl 2-

(N-methylcarbamoyl)ethyl, N,N-di-methylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl;

(mmm)Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are

5 hydrogen,

G² is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X² is OCH₂,

Q³ is selected from 3- isoxazolyl, 1,3-oxazol-4-yl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 2-pyridyl, 3-pyridyl and 2-pyrazinyl,

10 and Q³ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, cyano, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, 15 cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methylaminomethyl, 2-(methylamino)ethyl, di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-(ethyl)carbamoylmethyl 2-(N-(methyl)carbamoyl)ethyl, N,N-di-methylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylarbamoyl)ethyl, acetylmethyl, 20 methoxycarbonylmethyl and 2-methoxycarbonylethyl;

(nnn) Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

G² is chloro,

X² is OCH₂,

25 Q³ is selected from a 5 or 6-membered heteroaryl group containing at least 1 sp² hybridized nitrogen atom in the ortho position relative to X², for example isoxazolyl, 1,3-oxazol-4-yl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 2-pyridyl and 2-pyrazinyl, and Q³ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, cyano, hydroxy, carboxy, amino, methyl, ethyl, methoxy, ethoxy, 30 ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-methylcarbamoyl, N,N-di-ethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl,

methlyaminomethyl, 2-(methlyamino)ethyl, di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-(ethyl)carbamoylmethyl 2- (N-methyl)carbamoyl)ethyl, N,N-di-methylcarbamoylmethyl, N,N-di-ethylcarbamoylmethyl, 2-(N,N-di-methylcarbamoyl)ethyl, acetylmethyl,

5 methoxycarbonylmethyl and 2-methoxycarbonylethyl;

(ooo) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is chloro,

X^2 is OCH_2 ,

10 Q^3 is selected from 2-pyridyl, 3-pyridyl, 2-pyrazinyl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 5-methylisoxazol-3-yl, 1-methyl-1H-imidazol-5-yl and 1,3-oxazol-4-yl;

(ppp) Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is chloro,

15 X^2 is OCH_2 ,

Q^3 is phenyl optionally substituted by 1 or 2 substituents selected from fluoro and cyano (for example Q^3 is selected from phenyl, 2-fluorophenyl, 3-fluorophenyl, 2-cyanophenyl, 3-cyanophenyl, 2,5-difluorophenyl and 2,6-difluorophenyl);

(qqq) Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^6 and G^7 are
20 hydrogen,

G^4 is selected from hydrogen, halogeno, cyano, hydroxy, amino, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkylamino and di-[(1-4C)alkyl]amino,


X^3 is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-4C)alkyl, and Q^4 is selected from phenyl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 3-, 4- or 5-
25 isoxazolyl, 2-imidazolyl, 4-imidazolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl,

1,2,4-triazol-3-yl, 2-thienyl, 3-thienyl, 1,3-thiazol-2-yl, 1,3-thiazol-4-yl, 3-, 4- or 5-isothiazolyl, 1,2,3-thiadiazol-4-yl and 1,2,5-thiadiazol-3-yl, and wherein Q^4 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno,

trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl,

30 mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino,

N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl,
N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and
N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



- 5 wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl,
 10 (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;
 (rrr) Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^4 , G^6 and G^7 are hydrogen,

- X^3 is $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or methyl, and Q^4 is selected from phenyl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 3-isoxazolyl, 2-pyridyl, 3-pyridyl and
 15 4-thiazolyl and wherein Q^4 optionally bears 1 or 2 substituents selected from fluoro, chloro, cyano, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-methylcarbamoyl, N,N-di-ethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-
 20 cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methylamino)ethyl, di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-ethylcarbamoylmethyl 2- (N-methylcarbamoyl)ethyl, N,N-dimethylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl,
 25 methoxycarbonylmethyl and 2-methoxycarbonylethyl;
 (sss) Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^4 , G^6 and G^7 are hydrogen,
 X^3 is CH_2 and Q^4 is selected from phenyl, 2-fluorophenyl, 2-chlorophenyl, 2-methoxyphenyl, 2-cyanophenyl, 3-fluorophenyl, 2,5-dimethylphenyl, 2,6-di-fluorophenyl, 2-pyridyl, 3-pyridyl
 30 and 4-thiazolyl,
 (ttt) Q^2 is a group of the formula Id as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G⁴ is selected from hydrogen, halogeno, cyano, hydroxy, amino, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkylamino and di-[(1-4C)alkyl]amino,

X³ is selected from SO₂, CO and C(R⁷)₂, wherein each R⁷ is, independently, hydrogen or (1-4C)alkyl, and Q⁴ is selected from phenyl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 3-, 4- or 5-isoxazolyl, 2-imidazolyl, 4-imidazolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1,2,4-triazol-3-yl, 2-thienyl, 3-thienyl, 2-(1,3-thiazolyl), 4-(1,3-thiazolyl), 2-(1,3-thiazolyl), 3-, 4- or 5-isothiazolyl, 1,2,3-thiadiazol-4-yl and 1,2,5-thiadiazol-3-yl, and wherein Q⁴ optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X⁴ is a direct bond or is selected from O and N(R⁹), wherein R⁹ is hydrogen or methyl, and R⁸ is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;

(uuu) Q² is a group of the formula Id as hereinbefore defined wherein G¹, G⁴, G⁶ and G⁷ are hydrogen,

X³ is C(R⁷)₂, wherein each R⁷ is, independently, hydrogen or methyl, and Q⁴ is selected from phenyl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 3-isoxazolyl, 2-pyridyl and 4-thiazolyl and wherein Q⁴ optionally bears 1 or 2 substituents selected from fluoro, chloro, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, formyl, mercapto, acetyl, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methyamino)ethyl,

- di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(diethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-methylcarbamoylmethyl, N-ethylcarbamoylmethyl 2-(N-methylcarbamoyl)ethyl, N,N-dimethylcarbamoylmethyl, N,N-di-ethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl;
- (vvv) Q^2 is a group of the formula Id as hereinbefore defined wherein G^1 , G^4 , G^6 and G^7 are hydrogen,
- X^3 is CH_2 and Q^4 is phenyl optionally substituted by fluorine or chlorine, for example 3-fluorophenyl;
- 10 (www) Q^2 is selected from a group of the formula Ia, formula Ib or formula Id as hereinbefore defined;
- (xxx) Q^5 is 5 or 6 membered heterocyclyl ring linked to Q^3 or Q^4 by a ring nitrogen, and wherein Q^5 is optionally substituted by 1 or 2 substituents selected from fluoro, chloro, bromo, hydroxy, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino and
- 15 di-[(1-4C)alkyl]amino;
- (yyy) Q^5 is selected from pyrrolidin-1-yl, piperidino, morpholino and tetrahydro-4H-1,4-thiazin-4-yl; and
- (zzz) Q^1Z is selected from tetrahydrofuran-2-ylmethoxy, 2-tetrahydrofuran-2-ylethoxy, 3-tetrahydrofuran-2-ylpropoxy, tetrahydrofuran-3-ylmethoxy, 2-tetrahydrofuran-3-ylethoxy, 3-tetrahydrofuran-3-ylpropoxy, tetrahydropyran-2-ylmethoxy, 2-tetrahydropyran-2-ylethoxy, 3-tetrahydropyran-2-ylpropoxy, tetrahydropyran-3-ylmethoxy, 2-tetrahydropyran-3-ylethoxy, 3-tetrahydropyran-3-ylpropoxy, tetrahydropyran-4-ylmethoxy, 2-tetrahydropyran-4-ylethoxy, 3-tetrahydropyran-4-ylpropoxy, morpholin-2-ylmethoxy, 2-morpholin-2-ylethoxy, 3-morpholin-2-ylpropoxy, morpholin-3-ylmethoxy, 2-morpholin-3-ylethoxy, 3-morpholin-3-ylpropoxy,
- 25 morpholinomethoxy, 2-morpholinoethoxy and 3-morpholinopropoxy.

A particular embodiment of the present invention is a quinazoline derivative of the Formula I wherein:

- the Q^1Z - group is selected from cyclopentyloxy, cyclohexyloxy, tetrahydrofuran-3-yloxy, tetrahydropyran-3-yloxy, tetrahydropyran-4-yloxy,
- 30 tetrahydrothiopyran-3-yloxy, tetrahydrothiopyran-4-yloxy,
- 1,1-dioxotetrahydrothiopyran-3-yloxy, 1,1-dioxotetrahydrothiopyran-4-yloxy,
- 1-oxotetrahydrothiopyran-3-yloxy, 1-oxotetrahydrothiopyran-4-yloxy,
- tetrahydrothien-3-yloxy, 1,1-dioxodotetrahydrothien-3-yloxy, 1-oxotetrahydrothien-3-yloxy,

pyrrolidin-3-yloxy, pyrrolidin-2-yloxy, piperidin-3-yloxy, piperidin-4-yloxy,
homopiperidin-3-yloxy, homopiperidin-4-yloxy and azetidin-3-yloxy,

and wherein any azetidiny, pyrrolidinyl, piperidinyl or homopiperidinyl group within
the Q¹-Z- group is optionally N- substituted by a substituent T¹ selected from (1-4C)alkyl,
5 (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, (1-4C)alkoxycarbonyl, carbamoyl,
N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl and (1-4C)alkylsulphonyl,


and wherein adjacent carbon atoms in any (2-4C)alkylene chain within the N-
substituent T¹ are optionally separated by the insertion into the chain of a group selected from
O, NH and CO,

10 and wherein any CH₂ or CH₃ group within the N- substituent T¹ optionally bears on
each said CH₂ or CH₃ group a substituent selected from hydroxy, amino, methylamino,
di-methylamino, ethylamino, diethylamino, carbamoyl, N-methylcarbamoyl,
N,N-dimethylcarbamoyl, acetyl, methoxycarbonyl and ethoxycarbonyl,


and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo
15 substituents;

R¹ is hydrogen;


Q² is selected from a group of formula Ia, Ib and Id



Ia



Ib



Id

20

wherein G² and G³ are hydrogen,

G¹ is selected from hydrogen, fluoro, chloro, bromo, cyano, (1-4C)alkyl and
(2-3C)alkynyl,

G⁴ is selected from hydrogen, fluoro, chloro, bromo, cyano, (1-4C)alkyl and (2-3C)ethynyl,

X² is selected from O, S, NH, N(CH₃), OCH₂, CO and NHCH₂,

and Q³ is a group selected from phenyl, 2-furyl, 2- or 3-thienyl, 2-,4- or 5-oxazolyl, 3-,4- or 5-isoxazolyl, 2-,4-or 5-1H-imidazolyl, 2-,4-or 5-thiazolyl, 3- or 4-(1H-1,2,4-triazolyl), 3- or 5-(1,2,5-thiadiazolyl), 2- 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1,3-benzodioxol-4-yl, 1,3-benzodioxol-5-yl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl and 2-, 3-, 4-, 5-, 6-, 7- or 8-quinazolinyl, which group is optionally substituted with one or two substituents, which may be the same or different, selected from
10 nitro, fluoro, chloro, bromo, (1-4C)alkyl, trifluoromethyl, (1-4C)alkoxy, (2-3C)alkynyl and cyano,

or X² is CO and Q³ is a group selected from pyrrolidin-1-yl, piperidino, homopiperidino, morpholino, piperazin-1-yl, homopiperazin-1-yl, decahydroquinolin-1-yl, and decahydroisoquinolin-2-yl, which group optionally bears 1 or 2 substituents which may be
15 the same or different selected from fluoro, chloro, bromo, cyano, hydroxy and (1-4C)alkyl, and any heterocyclyl group represented by Q³ optionally bears 1 or 2 oxo substituents,

X³ is selected from SO₂ and CH₂,

Q⁴ is a group selected from phenyl, phenyl, 2-furyl, 3-furyl, 2-(1,3-oxazolyl), 4-(1,3-oxazolyl), 3-, 4- or 5-isoxazolyl, 2-imidazolyl, 4-imidazolyl, 2-, 3-or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1,2,4-triazol-3-yl, 2-thienyl, 3-thienyl, 2-(1,3-thiazolyl), 4-(1,3-thiazolyl), 3-, 4- or 5- isothiazolyl and 1,2,5-thiadiazol-3-yl group which group optionally bears 1 or 2 substituents, which may be the same or different, with one or two substituents which may be the same or different selected from nitro, fluoro, chloro, bromo, cyano, (1-4C)alkyl, (1-4C)alkoxy, trifluoromethyl and (2-3C)alkynyl;

25 or a pharmaceutically acceptable salt thereof.

A further embodiment of the present invention is a quinazoline derivative of the Formula I wherein:


the Q¹-Z- group is selected from cyclopentyloxy,

1-methylazetidin-3-yloxy, 1-isopropylazetidin-3-yloxy, tetrahydrothien-3-yloxy,
30 1-oxotetrahydrothien-3-yloxy, 1,1-dioxotetrahydrothien-3-yloxy, tetrahydrofuran-3-yloxy, 1-methylpyrrolidin-3-yloxy, tetrahydropyran-4-yloxy, tetrahydrothiopyran-4-yloxy, 1-oxotetrahydrothiopyran-4-yloxy, 1,1-dioxotetrahydrothiopyran-4-yloxy, piperidin-4-yloxy, 1-methylpiperidin-4-yloxy, 1-ethylpiperidin-4-yloxy, 1-propylpiperidin-4-yloxy,


1-(2-methoxyethyl)piperidin-4-yloxy, 1-acetylpiperidin-4-yloxy,
1-acetylmethylpiperidin-4-yloxy, 1-allylpiperidin-4-yloxy, 1-(2-propynyl)piperidin-4-yloxy,
1-methoxycarbonylmethylpiperidin-4-yloxy, 1-carbamoylmethylpiperidin-4-yloxy and
1-methanesulphonylpiperidin-4-yloxy;

5 R^1 is hydrogen;


Q^2 is selected from a group of formula Ia, Ib and Id



Ia



Ib



Id

10 wherein G^2 and G^3 are hydrogen,

G^1 is selected from hydrogen, fluoro, chloro, bromo, cyano, methyl and ethynyl,

G^4 is selected from hydrogen, fluoro, chloro, bromo, cyano, and methyl,

X^2 is selected from O, S, NH, OCH_2 , SCH_2 and $NHCH_2$,

15 and Q^3 is a group selected from phenyl, 2-1H-imidazolyl, 4-(1,3-thiazolyl), 2-thienyl,
3-(1,2,5-thiadiazolyl), 3-isoxazolyl 2-, 3- or 4-pyridyl, 2-pyrimidinyl, 1,3-benzodioxol-5-yl
and 8-quinolinyl, which group is optionally substituted with one or two substituents selected
from fluoro, chloro, bromo, nitro, trifluoromethyl, methyl, methoxy, ethynyl and cyano,

or X^2 is CO and Q^3 is a group selected from piperidino, homopiperidino,
20 decahydroquinolin-1-yl, and decahydroisoquinolin-2-yl, which group optionally bears 1 or 2
substituents selected from fluoro, chloro, bromo, cyano, hydroxy and methyl,
and wherein any heterocyclyl group represented by Q^3 optionally bears 1 or 2 oxo
substituents,

X^3 is selected from SO_2 and CH_2 ,

Q^4 is selected from phenyl and 2-pyridyl which optionally bears 1 or 2 substituents, which may be the same or different, selected fluoro, chloro, bromo, nitro, trifluoromethyl, methyl, methoxy, ethynyl and cyano;

or a pharmaceutically acceptable salt thereof.


5 A further embodiment of the present invention is a quinazoline derivative of the Formula I wherein:

the Q^1 -Z- group is selected from cyclopentyloxy, tetrahydrofuran-3-yloxy, 1-methylpyrrolidin-3-yloxy, pyrrolidin-3-yloxy, tetrahydropyran-4-yloxy, piperidin-4-yloxy, 1-methylpiperidin-4-yloxy, 1-ethylpiperidin-4-yloxy, 1-(2-methoxyethyl)piperidin-4-yloxy, 10 1-acetylpiperidin-4-yloxy, 1-acetylmethylpiperidin-4-yloxy, 1-allylpiperidin-4-yloxy, 1-(2-propynyl)piperidin-4-yloxy, 1-methoxycarbonylmethylpiperidin-4-yloxy, 1-carbamoylmethylpiperidin-4-yloxy and 1-methanesulphonylpiperidin-4-yloxy;

R^1 is hydrogen;

Q^2 is a group of the formula Ia

15



Ia

20

wherein G^1 and G^2 are each independently selected from hydrogen, fluoro and chloro, X^2 is selected from O, S, OCH_2 , NH, $N(CH_3)$, CO and $NHCH_2$, and Q^3 is selected from phenyl, 2-thienyl, 2-1H-imidazolyl, 3-isoxazolyl, 4-thiazolyl, 3-(1,2,5-thiadiazolyl), 2-pyridyl, 3-pyridyl, 4-pyridyl and 8-quinolynyl, which is optionally substituted by 1 or 2 substituents 25 selected from fluoro, chloro, methyl, methoxy, nitro and cyano; or a pharmaceutically acceptable salt thereof.


Suitable values for the group $-X^2-Q^3$ in this embodiment include, for example phenoxy, 3-fluorophenoxy, 2,3-difluorophenoxy, phenylthio, 2-fluorobenzoyloxy, 2-chlorobenzoyloxy, 2-cyanobenzoyloxy, 3-fluorobenzoyloxy, 3-fluorobenzoyloxy, 3-methylbenzoyloxy, 4-fluorobenzoyloxy, 2-methoxybenzoyloxy, 2,6-difluorobenzoyloxy, 2,6-dichlorobenzoyloxy, 2, 5-dimethylbenzoyloxy, 4-methyl-2-nitrobenzoyloxy, 3-fluorophenylaminomethyl, 5-chloro-2-thienyl, 2-thienylcarbonyl, 1-methyl-2-1H-imidazolyl, 1-methyl-2-1H-imidazolylmethoxy, 5-methyl-3-isoxazolylmethoxy, 2-methyl-4-thiazolylmethoxy, 1,2,5-thiadiazol-3-ylmethoxy, 2-pyridyl, 3-pyridyl, 2-pyridylmethoxy, 3-pyridylmethoxy, 4-pyridylmethoxy, 6-chloro-3-pyridylmethoxy, 2-pyridylmethylamino, 2-pyridyl(methyl)amino, 2-pyridyl(methyl)amino, 2-pyrimidinyl and 8-quinolinylthio.

A further embodiment of the present invention is a quinazoline derivative of the Formula I wherein:

the Q^1-Z group is selected from piperidin-4-yloxy, 1-methylpiperidin-4-yloxy, 1-ethylpiperidin-4-yloxy, 1-(2-methoxyethyl)piperidin-4-yloxy, 1-acetyl piperidin-4-yloxy, 1-acetylmethylpiperidin-4-yloxy, 1-allylpiperidin-4-yloxy, 1-(2-propynyl)piperidin-4-yloxy, 1-methoxycarbonylmethylpiperidin-4-yloxy, 1-carbamoylmethylpiperidin-4-yloxy and 1-methanesulphonylpiperidin-4-yloxy;

R^1 is hydrogen;

Q^2 is a group of the formula Ia



Ia

wherein G^1 and G^2 are each independently selected from hydrogen and chloro, X^2 is selected from O, S and OCH_2 , and Q^3 is selected from phenyl, 2-thienyl, 2-1H-imidazolyl, 3-isoxazolyl, 4-thiazolyl, 3-(1,2,5-thiadiazolyl), 2-pyridyl, 3-pyridyl, 4-pyridyl and 8-quinolinyl, which is optionally substituted by 1 or 2 substituents selected from fluoro, chloro, methyl, methoxy, nitro and cyano;

or a pharmaceutically acceptable salt thereof.

Suitable values for the group $-X^2-Q^3$ in this embodiment include, for example phenoxy, 3-fluorophenoxy, 2,3-difluorophenoxy, phenylthio, 2-fluorobenzoyloxy, 2-chlorobenzoyloxy, 2-cyanobenzoyloxy, 3-fluorobenzoyloxy, 3-fluorobenzoyloxy, 3-methylbenzoyloxy, 4-fluorobenzoyloxy, 2-methoxybenzoyloxy, 2,6-difluorobenzoyloxy, 2,6-dichlorobenzoyloxy, 2, 5-dimethylbenzoyloxy, 4-methyl-2-nitrobenzoyloxy, 5-chloro-2-thienylmethoxy, 1-methyl-2-1H-imidazolylmethoxy, 1-methyl-2-1H-imidazolylmethoxy, 5-methyl-3-isoxazolylmethoxy, 2-methyl-4-thiazolylmethoxy, 1,2,5-thiadiazol-3-ylmethoxy, 2-pyridylmethoxy, 3-pyridylmethoxy, 2-pyridylmethoxy, 3-pyridylmethoxy, 4-pyridylmethoxy, 6-chloro-3-pyridylmethoxy, 2-pyrimidinylmethoxy and 8-quinolinylthio.

A further embodiment of the present invention is a quinazoline derivative of the Formula I wherein

the Q^1-Z - group is selected from cyclopentylmethoxy, tetrahydrofuran-3-ylmethoxy, tetrahydropyran-4-ylmethoxy, tetrahydrothiopyran-4-ylmethoxy, 1,1-dioxotetrahydrothiopyran-4-ylmethoxy, 1-oxotetrahydrothiopyran-4-ylmethoxy, tetrahydrothien-3-ylmethoxy, 1,1-dioxotetrahydrothien-3-ylmethoxy, 1-oxotetrahydrothien-3-ylmethoxy, pyrrolidin-3-ylmethoxy, pyrrolidin-2-ylmethoxy, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, homopiperidin-3-ylmethoxy, homopiperidin-4-ylmethoxy and azetidin-3-ylmethoxy,


and wherein the azetidyl, pyrrolidinyl, piperidinyl or homopiperidinyl group within the Q^1-Z - group is optionally N- substituted by a substituent selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, allyl, 2-propynyl, acetyl, propionyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, methylsulphonyl, ethylsulphonyl, 2-methoxyethyl, carbamoylmethyl, N-methylcarbamoylmethyl, N,N-di-methylcarbamoylmethyl, 2-carbamoylethyl, 2-(N-methylcarbamoyl)ethyl, 2-(N,N-di-methylcarbamoyl)ethyl, acetylmethyl, 2-acetylethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl,

and wherein any heterocyclyl group within the Q^1-Z - group optionally bears 1 or 2 oxo substituents;

R^1 is hydrogen;

Q^2 is a group of the formula Ia

- 61 -



Ia

wherein G^1 and G^2 each independently is selected from hydrogen, fluoro, chloro, bromo, methyl and ethynyl,

X^2 is CO and Q^3 is selected from pyrrolidin-1-yl, piperidino, homopiperidino, morpholino,
 5 piperazin-1-yl, homopiperazin-1-yl, decahydroquinolin-1-yl and decahydroisoquinolin-2-yl,

and wherein Q^3 optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, cyano, hydroxy, amino, methyl, ethyl, methylamino and di-methylamino,

and wherein Q^3 optionally bears 1 or 2 oxo substituents;

or a pharmaceutically acceptable salt thereof.


10 A further embodiment of the present invention is a quinazoline derivative of the Formula I wherein

the Q^1 -Z- group is 1-methylpiperidin-4-yl-oxy,

R^1 is hydrogen;

Q^2 is a group of the formula Ia

15



Ia

wherein G^1 and G^2 each independently is selected from hydrogen, chloro and ethynyl,

X^2 is CO and Q^3 is selected from piperidino, homopiperidino, decahydroquinolin-1-yl and decahydroisoquinolin-2-yl, which is optionally substituted by methyl,

20 and wherein Q^3 optionally bears 1 or 2 oxo substituents;

or a pharmaceutically acceptable salt thereof.

Suitable values for Q^2 in this embodiment include for example

- 3-chloro-4-(homopiperidin-1-ylcarbonyl)phenyl,
- 3-chloro-4-(decahydroquinolin-1-ylcarbonyl)phenyl,
- 3-chloro-4-(decahydroisoquinolin-1-ylcarbonyl)phenyl,
- 5 3-chloro-4-((3-methylpiperidin-1-yl)carbonyl)phenyl,
- 3-chloro-4-((4-methylpiperidin-1-yl)carbonyl)phenyl,
- 3-ethynyl-4-(decahydroquinolin-1-ylcarbonyl)phenyl and
- 3-ethynyl-4-(decahydroquinolin-1-ylcarbonyl)phenyl.


A further embodiment of the invention is a quinazoline derivative of the Formula I

10 wherein:

the Q^1 -Z- group is 1-methylpiperidin-4-yl,

R^1 is hydrogen;

Q^2 is a group of the formula Ib



Ib

15 wherein G^1 and G^3 are hydrogen,

G^4 is selected from fluoro, chloro, bromo, cyano and methyl

X^3 is selected from CH_2 and SO_2 , and Q^4 is phenyl or 2-pyridyl optionally

bearing one substituent selected from fluoro, chloro and bromo;

or a pharmaceutically acceptable salt thereof.

20 Suitable values for Q^2 in this embodiment include, for example

- 1-benzenesulphonylindol-5-yl, 1-benzylindol-5-yl, 1-(3-fluorobenzyl)indol-5-yl and
- 1-(2-pyridylmethyl)indol-5-yl.


A further embodiment of the invention is a quinazoline derivative of the Formula I

wherein:

25 the Q^1 -Z- group is 1-methylpiperidin-4-yl,

R^1 is hydrogen;

Q^2 is a group of the formula Id



Id

5 wherein G^1 is hydrogen,

G^4 is selected from fluoro, chloro, bromo, cyano and methyl

X^3 is selected from CH_2 and SO_2 , and Q^4 is selected from phenyl and 2-pyridyl

optionally bearing one substituent selected from fluoro, chloro and bromo;
or a pharmaceutically acceptable salt thereof.

10 Suitable values for Q^2 in this embodiment include, for example
1-(2-pyridylmethyl)indazol-5-yl and 1-(3-fluorobenzyl)indazol-5-yl.

A further embodiment of the invention is a quinazoline derivative of the Formula I
wherein:

Z is O;

15 Q^1 is selected from a 5 or 6 membered heterocyclyl ring containing at least 1 nitrogen,
atom, said ring being linked to Z by a carbon atom,
and wherein any NH group within a heterocyclyl group optionally bears a substituent selected
from (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, formyl, (2-4C)alkanoyl,
(1-4C)alkoxycarbonyl, carbamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl,
20 N-(1-4C)alkylsulphamoyl, N,N-di-(1-4C)alkylsulphamoyl and (1-4C)alkylsulphonyl,
or from a group of the formula :



wherein X^1 is a direct bond and R^4 is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl,

(1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-6C)alkyl, amino-(1-4C)alkyl,

25 (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,

N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl,
(2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

5 **R**¹ is hydrogen;

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

G² is selected from fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X² is selected from S and OC(R⁶)₂, R⁶ is hydrogen or (1-4C)alkyl;

10 **Q**³ is selected from 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl 5-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-, 4- or 5-1H-pyrazolyl, 1H-1,2,4-triazol-3-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-pyridyl, 3-pyridyl, 2-pyrazinyl and phenyl, and

wherein Q³ optionally bears 1 or 2 substituents, which may be the same or different, selected
15 from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl,
20 (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X⁴ is a direct bond or is selected from O and N(R⁹), wherein R⁹ is hydrogen or
25 methyl, and R⁸ is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;

30 or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl and tetrahydropyran-4-yl,

and wherein Q¹ optionally bears an oxo substituent;

5 R¹ is hydrogen; and


Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

G² is selected from fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X² is selected from S and OCH₂,

10 Q³ is selected from 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 5-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-, 4- or 5-1H-pyrazolyl, 1H-1,2,4-triazol-3-yl, 1,2,5-thiadiazol-3-yl, 1,2,5-thiadiazol-3-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-pyridyl, 3-pyridyl, 2-pyrazinyl and phenyl, and

wherein Q³ optionally bears 1 or 2 substituents, which may be the same or different, selected
 15 from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl,
 20 (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X⁴ is a direct bond or is selected from O and N(R⁹), wherein R⁹ is hydrogen or
 25 methyl, and R⁸ is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl;
 30 or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl, tetrahydropyran-4-yl, pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl, and

wherein any NH group within a heterocyclyl group in **Q**¹ optionally bears a substituent
5 selected from (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkanoyl, carbamoyl,

N-(1-4C)alkylcarbamoyl and N,N-di-(1-4C)alkylcarbamoyl, or from a group of the formula :



wherein **X**¹ is a direct bond and **R**⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,
10 N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl or (2-4C)alkanoyl-(1-4C)alkyl,

and wherein any heterocyclyl group within the **Q**¹-**Z**- group optionally bears 1 or 2 oxo substituents;

R¹ is hydrogen; and

15 **Q**² is a group of the formula Ia as hereinbefore defined wherein **G**¹, **G**⁶ and **G**⁷ are hydrogen,

G² is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X² is S,

Q³ is selected from 2-1H-imidazolyl, 2-thiazolyl and 2- pyridyl;


20 and **Q**³ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, cyano, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-
25 cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methylaminomethyl, 2-(methylamino)ethyl, dimethylaminomethyl, 2-(dimethylamino)ethyl, 2-(diethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-methylcarbamoylmethyl, N-ethylcarbamoylmethyl 2-
(N-methylcarbamoyl)ethyl, N,N-di-methylcarbamoylmethyl, N,N-di ethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-
30 methoxycarbonylethyl;

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl, tetrahydropyran-4-yl, pyrrolidin-3-yl, piperidin-2-yl, piperidin-3-yl and piperidin-4-yl and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkanoyl, carbamoyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, or from a group of the formula :



wherein X¹ is a direct bond, and R⁴ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl or (2-4C)alkanoyl-(1-4C)alkyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents;

R¹ is hydrogen; and

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

G² is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X² is OCH₂,

Q³ is phenyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methyamino)ethyl, dimethylaminomethyl, 2-(dimethylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-ethylcarbamoylmethyl, 2-(N-methylcarbamoyl)ethyl, N,N-dimethylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl;

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

5 Q^1 is selected from tetrahydrofuran-3-yl, tetrahydropyran-3-yl, tetrahydropyran-4-yl, pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,

wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkanoyl, carbamoyl, N-(1-4C)alkylcarbamoyl and N,N-di-(1-4C)alkylcarbamoyl, or from a group of the formula :

10 $-X^1-R^4$

wherein X^1 is a direct bond, and R^4 is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl or (2-4C)alkanoyl-(1-4C)alkyl,

15 and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo substituents;

R^1 is hydrogen; and

Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

20 G^2 is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X^2 is OCH_2 ,

Q^3 is selected from 3-isoxazolyl, 1,3-oxazol-4-yl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 2-pyridyl, 3-pyridyl and 2-pyrazinyl,

and Q^3 optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, cyano,

25 carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, formyl, acetyl,

methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl,

N,N-di-methylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl,

2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl,

2-carboxyethyl, aminomethyl, 2-aminoethyl, methylaminomethyl, 2-(methylamino)ethyl,

30 dimethylaminomethyl, 2-(dimethylamino)ethyl, 2-(diethylamino)ethyl, carbamoylmethyl,

2-carbamoyl-ethyl, N-methylcarbamoylmethyl, N-ethylcarbamoylmethyl 2-

(N-methylcarbamoyl)ethyl, N,N-dimethylcarbamoylmethyl, N,N-di-ethylcarbamoylmethyl, 2-

(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-(methoxycarbonyl)ethyl;
or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I

5 wherein:

Z is O;

Q¹ is selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, which is substituted by a substituent selected from pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, morpholino, morpholin-2-yl, morpholin-3-yl, piperidino, piperidin-2-yl, 10 piperidin-3-yl, piperidin-4-yl, piperazin-1-yl, piperazin-2-yl, piperazin-3-yl, tetrahydro-4H-1,4-thiazin-2-yl, tetrahydro-4H-1,4-thiazin-3-yl, tetrahydro-4H-1,4-thiazin-4-yl, homopiperidin-1-yl, homopiperazin-1-yl, amino, methylamino, ethylamino, propylamino, dimethylamino diethylamino, aminomethyl, 2-aminoethyl, 3-aminopropyl, methylaminomethyl, ethylaminomethyl 2-methylaminoethyl, 2-ethylaminoethyl, 3- 15 methylaminopropyl, dimethylaminomethyl, 2-dimethylaminoethyl, 3-dimethylaminopropyl, diethylaminomethyl and 2-diethylaminoethyl,

and wherein any heterocyclyl ring in Q¹ optionally bears 1 or 2 substituents selected from fluoro, chloro, hydroxy, (1-3C)alkyl, (1-3C)alkoxy, amino, methylamino, ethylamino, dimethylamino, diethylamino, acetyl, propionyl, methoxycarbonyl, ethoxycarbonyl, 20 methylsulphonyl, ethylsulphonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl and N,N-diethylcarbamoyl,

and wherein any cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl group in Q¹ is optionally further substituted by 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro and methyl,

25 and wherein any heterocyclyl group in Q¹ optionally bears an oxo substituent;

R¹ is hydrogen; and


Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

G² is selected from fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

30 X² is selected from S and OCH₂,

Q³ is selected from 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-, 4- or 5-1H-pyrazolyl, 1H-1,2,4-triazol-3-yl, 1,2,5-thiadiazol-3-yl,

1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-pyridyl, 3-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 2-pyrazinyl and phenyl, and wherein Q³ optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :



wherein X⁴ is a direct bond or is selected from O and N(R⁹), wherein R⁹ is hydrogen or methyl, and R⁸ is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl; or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is 4-(piperazin-1-yl)cyclohexyl, wherein the piperazin-1-yl group is optionally substituted at the 4-position by a substituent selected from (1-3C)alkyl, (2-4C)alkanoyl, (1-3C)alkoxycarbonyl, (1-3C)alkylsulphonyl, carbamoyl, N-(1-3C)alkylcarbamoyl and N,N-di-[(1-3C)alkyl]carbamoyl,

and wherein the piperazin-1-yl group in Q¹ optionally bears an oxo substituent;

R¹ is hydrogen; and

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen;

G² is selected from fluoro, chloro, methyl, ethyl and ethynyl,

X² is selected from OCH₂ and S,

Q³ is selected from phenyl, 2-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-isoxazolyl, 1,2,4-oxadiazol-3-yl, 2-pyridyl and 2-pyrazinyl, and Q³ optionally bears 1 or 2 substituents selected from fluoro, chloro, bromo, cyano, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-methylcarbamoyl, N,N-di-ethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methyamino)ethyl, di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-(ethyl)carbamoylmethyl, 2-(N-(methyl)carbamoyl)ethyl, N,N-di-[methyl]carbamoylmethyl, N,N-di-[ethyl]carbamoylmethyl, 2-(N,N-di-[methyl]carbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl; or a pharmaceutically acceptable salt thereof.

15 A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is 4-(piperazin-1-yl)cyclohexyl, wherein the piperazin-1-yl group is optionally substituted at the 4-position by (1-3C)alkyl,

20 and wherein the piperazin-1-yl group in Q¹ optionally bears an oxo substituent;

R¹ is hydrogen; and

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen;

G² is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

25 X² is OCH₂; and

Q³ is phenyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl methyl, ethyl, methoxy, ethoxy, ethynyl, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylamino, dimethylamino and methylsulphonyl; or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from pyrrolidin-1-ylmethyl,

5 2-pyrrolidin-2-ylethyl, piperidin-2-ylmethyl, 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl and 3-homopiperidin-1-ylpropyl,

10 and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl),

15 and wherein any heterocyclyl group within the Q¹-Z- group optionally bears an oxo substituent;

R¹ is hydrogen; and

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

20 G² is chloro,

X² is OCH₂,

Q³ is selected from phenyl optionally substituted by 1 or 2 substituents selected from fluoro and cyano (for example Q³ is selected from phenyl, 2-fluorophenyl, 3-fluorophenyl, 2-cyanophenyl, 3-cyanophenyl, 2,5-difluorophenyl and 2,6-difluorophenyl);

25 or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,

30 and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl,

- 73 -

cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl) and wherein any heterocyclyl group within the Q¹-Z- group optionally bears an oxo substituent;

5 **R¹** is hydrogen; and

Q² is a group of the formula Ia as hereinbefore defined wherein G¹, G⁶ and G⁷ are hydrogen,

 G² is chloro,

 X² is OCH₂,

10 **Q³** is selected from phenyl optionally substituted by 1 or 2 substituents selected from fluoro and cyano (for example **Q³** is selected from phenyl, 2-fluorophenyl, 3-fluorophenyl, 2-cyanophenyl, 3-cyanophenyl, 2,5-difluorophenyl and 2,6-difluorophenyl);
or a pharmaceutically acceptable salt thereof.

 Suitable values for **Q¹** in this embodiment include, for example 1-methylpiperidin-4-yl
15 or 1-methylpiperidin-3-yl.

 A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q¹ is selected from pyrrolidin-1-ylmethyl, 2-pyrrolidin-2-ylethyl, piperidin-2-ylmethyl,
20 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl and 3-homopiperidin-1-ylpropyl,

25 and wherein any NH group within a heterocyclyl group in **Q¹** optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl)

30 and wherein any heterocyclyl group within the **Q¹**-Z- group optionally bears an oxo substituent;

R¹ is hydrogen; and

Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is chloro,

X^2 is OCH_2 ,

5 Q^3 is selected from 2-pyridyl, 3-pyridyl, 2-pyrazinyl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 5-methyl-isoxazol-3-yl, 1-methyl-1H-imidazol-5-yl and 1,3-oxazol-4-yl; or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

10 Z is O;

Q^1 is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,

and wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, 15 cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl) and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears an oxo substituent;

R^1 is hydrogen; and

20 Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is chloro,

X^2 is OCH_2 ,

25 Q^3 is selected from 2-pyridyl, 3-pyridyl, 2-pyrazinyl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 5-methylisoxazol-3-yl, 1-methyl-1H-imidazol-5-yl and 1,3-oxazol-4-yl; or a pharmaceutically acceptable salt thereof.

Suitable values for Q^1 in this embodiment include, for example 1-methylpiperidin-4-yl or 1-methylpiperidin-3-yl.

A further embodiment of the invention is a quinazoline derivative of the Formula I 30 wherein:

Z is O;

Q^1 is selected from pyrrolidin-1-ylmethyl, 2-pyrrolidin-2-ylethyl, piperidin-2-ylmethyl, 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl and 3-homopiperidin-1-ylpropyl,

and wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl)

and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears an oxo substituent;

R^1 is hydrogen; and

Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^4 , G^6 and G^7 are hydrogen,

X^3 is CH_2 and Q^4 is selected from phenyl, 2-fluorophenyl, 2-chlorophenyl, 2-methoxyphenyl, 2-cyanophenyl, 3-fluorophenyl, 2,5-dimethylphenyl, 2,6-difluorophenyl, 2-pyridyl, 3-pyridyl and 4-thiazolyl;

or a pharmaceutically acceptable salt thereof.

A further embodiment of the invention is a quinazoline derivative of the Formula I wherein:

Z is O;

Q^1 is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,

and wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl (conveniently methyl, 2-methoxyethyl or carbamoylmethyl)

and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears an oxo substituent;

R^1 is hydrogen; and

Q^2 is a group of the formula Ib as hereinbefore defined wherein G^1 , G^3 , G^4 , G^6 and G^7 are hydrogen,

X^3 is CH_2 and Q^4 is selected from phenyl, 2-fluorophenyl, 2-chlorophenyl, 2-methoxyphenyl, 2-cyanophenyl, 3-fluorophenyl, 2,5-dimethylphenyl, 2,6-difluorophenyl,
 5 2-pyridyl, 3-pyridyl and 4-thiazolyl;
 or a pharmaceutically acceptable salt thereof.

Suitable values for Q^1 in this embodiment include, for example 1-methylpiperazin-4-yl or 1-methylpiperazin-3-yl.

A further particular preferred compound of the invention is, for example, a
 10 quinazoline derivative of the Formula I selected from:
 4-(3-Chloro-4-phenoxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-((3-fluorobenzoyloxy)anilino))-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-(3-Fluorobenzyl)indazol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(decahydroquinolin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-
 15 yloxy)quinazoline;
 4-(3-Chloro-4-(decahydroisoquinolin-2-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(3-methylpiperidin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 20 4-(3-Ethynyl-4-(decahydroquinolin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(2, 6-difluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 25 4-(3-Chloro-4-(2-fluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(2-cyanobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and
 4-(3-Chloro-4-(thiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 or a pharmaceutically acceptable acid addition salt thereof.

A further particular preferred compound of the invention is, for example, a
 30 quinazoline derivative of the Formula I selected from:
 4-(1-Benzylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-Benzenesulphonylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-(2-Cyanobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

- 4-(1-(2-Methoxybenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-(2-Chlorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-(2, 5-Dimethylbenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(1-(2-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and
 5 4-(1-(3-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline
 or a pharmaceutically acceptable acid addition salt thereof.

A further particular preferred compound of the invention is, for example, a quinazoline derivative of the Formula I selected from:

- 4-(3-Chloro-4-(3-fluorophenoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 10 4-(3-Chloro-4-(2-thienoyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-((1-methyl-1*H*-imidazol-2-yl)thio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and
 4-(3-Chloro-4-(1-cyanomethyl-1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 15 or a pharmaceutically acceptable acid addition salt thereof.


A further particular preferred compound of the invention is, for example, a quinazoline derivative of the Formula I selected from:

- 4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline;
 4-(4-Benzyloxy-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 20 4-(3-Methyl-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(4-(3-Fluorobenzyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Ethynyl 4-(3-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Ethynyl 4-(2, 6-difluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 25 4-(4-(2-Aminothiazol-4-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(pyrazin-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and
 4-(3-Chloro-4-(pyridin-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 or a pharmaceutically acceptable acid addition salt thereof.
- 30 A quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes, when used to prepare a quinazoline derivative of the Formula I are provided as a further feature of the invention and are illustrated

by the following representative process variants in which, unless otherwise stated, Q^1 , Z , R^1 and Q^2 have any of the meanings defined hereinbefore. Necessary starting materials may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described in conjunction with the following representative process variants and

5 within the accompanying Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

Process (a) The reaction, conveniently in the presence of a suitable base, of a quinazoline of the Formula II



II

10 wherein L^1 is a displaceable group and Q^1 and Z have any of the meanings defined hereinbefore except that any functional group is protected if necessary, with a compound of the Formula



15 wherein Q^2 and R^1 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means.

A suitable base is, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, di-isopropylethylamine,

20 N-methylmorpholine or diazabicyclo[5.4.0]undec-7-ene, or, for example, an alkali or alkaline earth metal carbonate, for example sodium carbonate, potassium carbonate, calcium carbonate, or, for example, an alkali metal hydride, for example sodium hydride.

A suitable displaceable group L^1 is, for example, a halogeno, alkoxy, aryloxy, mercapto, alkylthio, arylthio, alkylsulphinyl, arylsulphinyl, alkylsulphonyl, arylsulphonyl or

25 sulphonyloxy group, for example a chloro, bromo, methoxy, phenoxy, pentafluorophenoxy, methylthio, methanesulphonyl, methanesulphonyloxy or toluene-4-sulphonyloxy group. The reaction is conveniently carried out in the presence of a suitable inert solvent or diluent, for example an alcohol or ester such as methanol, ethanol, isopropanol or ethyl acetate, a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether

such as tetrahydrofuran or 1,4-dioxan, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulphoxide. The reaction is conveniently carried out at a temperature in the range, for example, 10 to 250°C, conveniently in the range 40 to 80°C.

5 The quinazoline of the Formula II may also be reacted with a compound of the formula Q^2NHR^1 in the presence of a protic solvent such as isopropanol, conveniently in the presence of an acid, for example hydrogen chloride gas in diethyl ether or dioxane, or hydrochloric acid. Alternatively, this reaction may be conveniently carried out in an aprotic solvent, such as dioxane or a dipolar aprotic solvent such as N,N-dimethylacetamide in the presence of an
10 acid, for example hydrogen chloride gas in diethyl ether or dioxane, or hydrochloric acid. The above reactions are conveniently carried out at a temperature in the range, for example, 0 to 150°C, preferably at or near the reflux temperature of the reaction solvent.

 The quinazoline derivative of the Formula II, wherein L^1 is halogeno, may be reacted with a compound of the formula Q^2NHR^1 in the absence of an acid or a base. In this reaction
15 displacement of the halogeno leaving group L^1 results in the formation of the acid HL^1 in-situ and the autocatalysis of the reaction. Conveniently the reaction is carried out in a suitable inert organic solvent, for example isopropanol, dioxane or N,N-dimethylacetamide. Suitable conditions for this reaction are as described above.

 The quinazoline derivative of the Formula I may be obtained from this process in the
20 form of the free base or alternatively it may be obtained in the form of a salt with the acid of the formula $H-L^1$ wherein L^1 has the meaning defined hereinbefore. When it is desired to obtain the free base from the salt, the salt may be treated with a suitable base, for example, an alkali or alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide.

25 Protecting groups may in general be chosen from any of the groups described in the literature or known to the skilled chemist as appropriate for the protection of the group in question and may be introduced by conventional methods. Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so
30 as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

 Specific examples of protecting groups are given below for the sake of convenience, in which "lower", as in, for example, lower alkyl, signifies that the group to which it is applied

preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned are, of course, within the scope of the invention.

5 A carboxy protecting group may be the residue of an ester-forming aliphatic or arylaliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (for example isopropyl, and tert-butyl); lower alkoxy- lower alkyl groups (for example methoxymethyl, ethoxymethyl and
10 isobutoxymethyl); lower acyloxy-lower alkyl groups, (for example acetoxymethyl, propionyloxymethyl, butyryloxymethyl and pivaloyloxymethyl); lower alkoxycarbonyloxy-lower alkyl groups (for example 1-methoxycarbonyloxyethyl and 1-ethoxycarbonyloxyethyl); aryl-lower alkyl groups (for example benzyl, 4-methoxybenzyl, 2-nitrobenzyl, 4-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (for
15 example trimethylsilyl and tert-butyldimethylsilyl); tri(lower alkyl)silyl-lower alkyl groups (for example trimethylsilylethyl); and (2-6C)alkenyl groups (for example allyl). Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, base-, metal- or enzymically-catalysed cleavage.

Examples of hydroxy protecting groups include lower alkyl groups (for example
20 tert-butyl), lower alkenyl groups (for example allyl); lower alkanoyl groups (for example acetyl); lower alkoxycarbonyl groups (for example tert-butoxycarbonyl); lower alkenyloxycarbonyl groups (for example allyloxycarbonyl); aryl-lower alkoxycarbonyl groups (for example benzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl); tri(lower alkyl)silyl (for example
25 trimethylsilyl and tert-butyldimethylsilyl) and aryl-lower alkyl (for example benzyl) groups.

Examples of amino protecting groups include formyl, aryl-lower alkyl groups (for example benzyl and substituted benzyl, 4-methoxybenzyl, 2-nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-4-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (for example tert-butoxycarbonyl); lower alkenyloxycarbonyl (for example
30 allyloxycarbonyl); aryl-lower alkoxycarbonyl groups (for example benzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl and 4-nitrobenzyloxycarbonyl); lower alkanoyloxyalkyl groups (for example pivaloyloxymethyl); trialkylsilyl (for example

trimethylsilyl and tert-butyldimethylsilyl); alkylidene (for example methyldiene) and benzylidene and substituted benzylidene groups.


Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base-, metal- or enzymically-catalysed hydrolysis for groups such as

5 2-nitrobenzyloxycarbonyl, hydrogenation for groups such as benzyl and photolytically for groups such as 2-nitrobenzyloxycarbonyl. For example a tert butoxycarbonyl protecting group may be removed from an amino group by an acid catalysed hydrolysis using trifluoroacetic acid.

The reader is referred to Advanced Organic Chemistry, 4th Edition, by J. March,

10 published by John Wiley & Sons 1992, for general guidance on reaction conditions and reagents and to Protective Groups in Organic Synthesis, 2nd Edition, by T. Green *et al.*, also published by John Wiley & Son, for general guidance on protecting groups.

Quinazoline starting materials of the Formula II may be obtained by conventional procedures. For example, a 3,4-dihydroquinazolin-4-one of Formula III



III

15 wherein Q¹ and Z have any of the meanings defined hereinbefore except that any functional group is protected if necessary, may be reacted with a halogenating agent such as thionyl chloride, phosphoryl chloride or a mixture of carbon tetrachloride and triphenylphosphine whereafter any protecting group that is present is removed by conventional means. The

20 reaction is conveniently carried out in a suitable inert solvent, for example 1,2-dichloroethane or N,N-dimethylformamide conveniently in the presence of an base such as an organic base, for example di-isopropylethylamine. The reaction is conveniently carried out at a temperature in the range, for example, 0 to 150°C, preferably at or near the reflux temperature of the reaction solvent.


25 The 4-chloroquinazoline so obtained may be converted, if required, into a 4-pentafluorophenoxyquinazoline by reaction with pentafluorophenol in the presence of a suitable base such as potassium carbonate and in the presence of a suitable solvent such as N,N-dimethylformamide.

The compound of Formula I may be also be prepared using a telescoped process stating from the compound of Formula III, wherein the compound of the Formula Q^2NHR^1 is reacted with the compound of Formula II following halogenation of the compound of Formula III. The use of such a telescoped process avoids the need to isolate the compound of Formula II prior to reaction with the compound of formula Q^2NHR^1 .

The 3,4-dihydroquinazolin-4-one of Formula III may be obtained using conventional procedures. For example when Z is an oxygen atom the compound of Formula III may be prepared as illustrated by Reaction Scheme 1 starting with the compound of Formula IV.

Reaction Scheme 1

10



wherein Q^1 is as hereinbefore defined, X is (1-6C)alkyl (for example methyl) or benzyl, and Pg is a suitable amine protecting group.

15 Notes:

Step (1)

When X is (1-6C)alkyl, it may be may be cleaved from the compound of formula IV by conventional methods, such as by treatment of the compound of Formula IV with, for example:

- 20 (i) an alkali metal (1-6C)alkylsulphide such as sodium ethanethiolate;
 (ii) an alkali metal diarylphosphide such as lithium diphenylphosphide;
 (iii) a boron or aluminium trihalide such as boron tribromide;
 (iv) magnesium bromide in pyridine; or
 (v) pyridine hydrochloride in pyridine.

Generally the cleavage reaction is carried out at a temperature in the range of from, for example, 40 to 150°C.

When X is benzyl, it may be may be cleaved from the compound of formula IV by, for example, acid catalysed hydrolysis, for example by treatment of the compound of Formula IV
5 with trifluoroacetic acid. Conveniently the reaction is carried out at a temperature in the range of 30 to 120°C, for example 70°C.

Step (2)

The protecting group Pg is added to the 3,4-dihydro-5-hydroxyquinazolin-4-one of
Formula IVa using conventional techniques. For example a suitable Pg is a
10 pivaloyloxymethyl group that may be added to the compound of Formula IVa by reacting the compound of Formula IVa with chloromethylpivalate in the presence of a suitable base such as sodium hydride.

Step (3)


The Q¹O group may be introduced by coupling the compound of Formula IVb with a
15 compound of the Formula Q¹OH in the presence of a suitable dehydrating agent. Suitable conditions for the coupling reaction are described below with reference to process (b).

Step (4)

The protecting group Pg may be removed using conventional methods, for example
when Pg is a pivaloyloxymethyl group it may be removed by treating the compound of
20 Formula IVc with a methanolic ammonia solution.

The compound of formula IV may, for example, be prepared starting from an aniline of the Formula V as illustrated in Reaction Scheme 2.

Reaction Scheme 2



wherein X is as hereinbefore defined.

Notes:

Steps 1 and 2 may be carried out using analogous conditions to the processes described in Organic Syntheses, Coll. Vol. 1, p 327-330; J. Org. Chem. 1969, 34, 3484-3489.


- 5 Step 3 may be carried out using analogous conditions to the process described in J. Org. Chem. 1952, 17, 141-148; J. Med. Chem. 1994, 37, 2106-2111.

Anilines of Formula V are commercially available compounds, or they are known in the literature, or can be prepared using well known processes in the art.

In an alternative process the compound of formula III may be obtained according to

10 Reaction Scheme 1a:


Reaction Scheme 1a



- The reaction is conveniently performed in the presence of a suitable base. Suitable bases are
 15 as herein described under process (a), for example sodium hydride. The reaction is conveniently performed in a suitable inert solvent, for example N,N-dimethylacetamide.

- Reaction Scheme 1a is particularly suitable for the preparation of compounds of the formula III in which Z is O. The 5-fluoro-3,4-dihydroquinazoline starting material is commercially available or can be prepared using conventional methods, for example as described in J. Org.
 20 Chem. 1952, 17, 164-176.

Compounds of the Formula Q^2NHR^1 are commercially available compounds, or they are known in the literature, or can be prepared using conventional synthetic methods. For example when Q^2 is a group of the formula Ia the compound of the formula Q^2NHR^1 may be prepared in accordance with Reaction Scheme 3 or Reaction Scheme 4.

Reaction Scheme 3

wherein X^2 is as hereinbefore defined, for example, NR^6 , S, O or $\text{NR}^6\text{C}(\text{R}^6)_2$ and $G^1, G^2, G^6,$
 5 G^7, L^1, Q^3 and R^6 are as hereinbefore defined, except any functional group is protected if
 necessary, and whereafter any protecting group that is present is removed by conventional
 means.

Notes

Step 1 may be performed under analogous conditions to those used in process (a)
 10 described above. The compounds of the formula HX^2Q^3 are commercially available, or they
 are known in the literature, or can be prepared using well known processes in the art.

The reduction of the nitro group in step 2 may be carried out under standard
 conditions, for example by catalytic hydrogenation over a platinum/carbon, palladium/carbon
 or nickel catalyst, treatment with a metal such as iron, titanium chloride, tin II chloride or
 15 indium, or treatment with another suitable reducing agent such as sodium dithionite.

In an variation of process (a) the reduction of the nitro-compound in step 2 of reaction
 scheme 3 may be carried out as described above, followed directly with reaction with the
 compound of of formula II in a telescoped process, thereby avoiding the need to isolate the
 compound of the formula Q^2NHR^1 .

20 When Q^2 is a compound of the formula 1a in which X^2 is OCH_2 compounds of the
 formula Q^2NHR^1 may, for example, be prepared by reacting the starting nitro compound
 shown in Reaction Scheme 3 in which L^1 is OH with a compound of the formula Q^3CH_2 -
 halide (for example Q^3CH_2 -bromide). The nitro group may then be reduced to an amino
 group by using step 2 of the process in Reaction Scheme 3. Such compounds may also be
 25 prepared by reacting the starting nitro compound shown in Reaction Scheme 3 in which L^1 is
 halide with a compound of the formula $\text{Q}^3\text{CH}_2\text{OH}$, followed by reduction of the nitro group as
 described above in Reaction Scheme 3.


Compounds of the formula Q^3CH_2OH are known or may be prepared using known methods, for example by reduction of the corresponding ester of the formula Q^3COOR , wherein R is, for example (1-6C)alkyl, or benzyl, with a suitable reducing agent, for example sodium borohydride.

- 5 When Q^2 is a compound of the formula 1a in which X^2 is OCH_2 compounds of the formula Q^2NHR^1 may, for example, be prepared by coupling the starting nitro compound shown in Reaction Scheme 3 in which L^1 is OH with a compound of the formula Q^3CH_2OH , conveniently in the presence of a suitable dehydrating agent. Suitable conditions for performing this reaction are analogous to those described below in relation to Process(b).


- 10 When Q^2 is a compound of the formula 1a in which X^2 is $C(R^6)_2NR^6$ or $NR^6C(R^6)_2$ compounds of the formula Q^2NHR^1 may, for example, be prepared according to Reaction Scheme 3a:

Reaction Scheme 3a

15



- wherein G¹, G², G⁶, G⁷, L¹, Q³ and R⁶ are as hereinbefore defined, except any functional group is protected if necessary, and whereafter any protecting group that is present is removed by conventional means. The first step of Reaction Scheme 3a may be performed under analogous conditions to those used in process (a) described above. The starting nitro compounds and the compounds of the formula Q^3NR^6H and $Q^3C(R^6)_2L^1$ are commercially available, or they are known in the literature, or can be prepared using well known processes in the art. The reduction of the nitro group in step 2 may be carried out under analogous conditions to those described above for Reaction Scheme 3.

Reaction Scheme 4

wherein G^1 , G^2 , G^6 , G^7 , Q^3 and R^6 are as hereinbefore defined, except any functional group is protected if necessary, and whereafter any protecting group that is present is removed by conventional means, and L^1 is a suitable leaving group such as halide, for example chloro.

Notes

Suitable for the preparation of those compounds wherein X^2 is CO or CH_2NR^6 .


Step 1 may be carried out under analogous conditions to those used in process (a) described above.

10 The reduction of the nitro group in step 2 may be carried out as described above in Reaction Scheme 3.


When Q^2 is a compounds of the formula Ia in which X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom, the compound of the formula Q^2NHR^1 may be prepared by coupling the starting nitro compound shown in Reaction Scheme 15 3 in which L^1 is carboxy with a compound of the formula Q^3H in the presence of a suitable coupling agent, for example O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (HATU). Suitable conditions for this reaction are analogous to those described in relation to process (j) below.

When Q^2 is a compound of the formula Ib, Ic, Id and Ie, the compound of the formula 20 Q^2NHR^1 may be prepared, for example, by reacting an appropriate nitro-indole or nitro indazole with a compound of the formula Q^4X^3 -halide, suitably in the presence of a base. The reaction is conveniently carried out in a suitable inert solvent, under analogous conditions to those described for Process (a). The nitro group on the resulting indole or indazole may then be reduced to an amino group using an analogous process to that described in step 2 in 25 Reaction Scheme 3.

Process (b) For the production of those compounds of the Formula I wherein Z is an oxygen atom, the coupling, conveniently in the presence of a suitable dehydrating agent, of an alcohol of the Formula



- 5 wherein Q^1 has any of the meanings defined hereinbefore except that any functional group is protected if necessary with a quinazoline of the Formula VI




VI

- wherein R^1 and Q^2 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by
10 conventional means.

A suitable dehydrating agent is, for example, a carbodiimide reagent such as dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide or a mixture of an azo compound such as diethyl or di-tert-butyl azodicarboxylate and a phosphine such as triphenylphosphine. The reaction is conveniently carried out in the presence of a suitable inert
15 solvent or diluent, for example a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride and at a temperature in the range, for example, 0 to 150°C, preferably at or near ambient temperature.

The quinazoline of the Formula VI may be obtained by conventional procedures. For example, by cleavage of the group represented by X from the compound of the Formula VII

20



VII

wherein X is as defined hereinbefore and R^1 and Q^2 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means.

The cleavage reaction is conveniently carried out as hereinbefore described in relation to step (1) in Reaction Scheme 1.

The compound of Formula VII may be prepared by for example reacting the compound of the Formula (IV) as hereinbefore defined, with a halogenating agent such as thionyl chloride, phosphoryl chloride or a mixture of carbon tetrachloride and triphenylphosphine. The resulting compound is then reacted with a compound of the Formula

10 Q^2NHR^1

wherein Q^2 and R^1 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means. The halogenation reaction may be performed under analogous conditions to those described above in relation to the reaction with the compound of the


15 Formula III. The subsequent reaction with the compound of the Formula Q^2NHR^1 may be performed under analogous conditions to those described above in relation to the reaction with the compound of the Formula II.

Process(c) For the production of those compounds of the Formula I wherein Z is an oxygen atom, the reaction, conveniently in the presence of a suitable base, of an alcohol of the

20 Formula

Q^1-OH

wherein Q^1 has any of the meanings defined hereinbefore except that any functional group is protected if necessary with a quinazoline of the Formula VIII




VIII

25 wherein R^1 and Q^2 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means.

A suitable base includes, for example a strong non-nucleophilic base such as an alkali metal hydride, for example sodium hydride, or an alkali metal amide, for example lithium di-isopropylamide (LDA).

The reaction is conveniently carried out in the presence of a suitable inert solvent or diluent, for example a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulphoxide. The reaction is conveniently carried out at a temperature in the range, for example, 10 to 250°C, preferably in the range 40 to 150°C. Conveniently, this reaction may also be performed by heating the reactants in a sealed vessel using a suitable heating apparatus such as a microwave heater.

The quinazoline of the Formula VIII may be obtained by conventional procedures. For example, a quinazoline of the Formula IX



IX

wherein L^1 is a displaceable group as defined hereinbefore (such as halogeno, for example chloro), may be reacted with a compound of the Formula



wherein Q^2 and R^1 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means. The reaction may be performed under analogous conditions to those described above under Process (a).

The quinazoline of Formula IX may be obtained using conventional methods, for example 5-fluoro-3,4-dihydroquinazolin-4-one may be reacted with a halogenating agent such as thionyl chloride, phosphoryl chloride or a mixture of carbon tetrachloride and triphenylphosphine whereafter any protecting group that is present is removed by conventional means, as described in relation to Process (a) above.

Process (d) For the production of those compounds of the Formula I wherein Q¹ or Q² contains a primary or secondary amino group, the cleavage of the corresponding compound of Formula I wherein Q¹ or Q² contains a protected primary or secondary amino group.

Suitable protecting groups for an amino group are, for example, any of the protecting groups disclosed hereinbefore for an amino group. Suitable methods for the cleavage of such amino protecting groups are also disclosed hereinbefore. In particular, a suitable protecting group is a lower alkoxy carbonyl group such as a *tert*-butoxycarbonyl group which may be cleaved under conventional reaction conditions such as under acid-catalysed hydrolysis, for example in the presence of trifluoroacetic acid.

10. Process (e) For the production of those compounds of the Formula I wherein Q¹ or Q² contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, the alkylation, conveniently in the presence of a suitable base as defined hereinbefore, of a quinazoline derivative of the formula I wherein Q¹ or Q² contains a hydroxy group or a primary or secondary amino group as appropriate.


15 A suitable alkylating agent is, for example, any agent known in the art for the alkylation of hydroxy to alkoxy or substituted alkoxy, or for the alkylation of amino to alkylamino or substituted alkylamino, for example an alkyl or substituted alkyl halide, for example a (1-6C)alkyl chloride, bromide or iodide or a substituted (1-6C)alkyl chloride, bromide or iodide, conveniently in the presence of a suitable base as defined hereinbefore, in a suitable inert solvent or diluent as defined hereinbefore and at a temperature in the range, for example, 10 to 140°C, conveniently at or near ambient temperature. An analogous procedure may be used to introduce optionally substituted (2-6C)alkenyloxy, (2-6C)alkenylamino, (2-6C)alkynyloxy or (2-6C)alkynylamino groups into Q¹ or Q². Examples of substituents that may be introduced to a compound of formula I using this process (e) include, for example the substitution of an NH group in a heterocyclyl represented by Q¹ with an ethyl, allyl, 2-propynyl, 2-methoxyethyl or acetylmethyl, carbamoyl or methanesulphonyl group.

Process (f) For the production of those compounds of the Formula I wherein Q¹ contains an amino-hydroxy-disubstituted (1-6C)alkoxy group (such as 2-hydroxy-3-piperidinopropoxy, 2-hydroxy-3-methylaminopropoxy, 3-dimethylamino-2-hydroxypropoxy or 3-[N-(3-dimethylaminopropyl)-N-methylamino]-2-hydroxypropoxy), the reaction of a compound of the Formula I wherein Q¹ contains an epoxy-substituted (1-6C)alkoxy group with a heterocycle or an appropriate amine.

The reaction is conveniently carried out in the presence of a suitable inert diluent or

carrier as defined hereinbefore and at a temperature in the range 10 to 150°C, preferably at or near ambient temperature.

Process (g) The reaction, conveniently in the presence of a suitable base as defined hereinbefore, of a quinazoline of the Formula X



5

wherein L¹ is a displaceable group as defined hereinbefore and R¹ and Q² have any of the meanings defined hereinbefore except that any functional group is protected if necessary, with a compound of the Formula



10 wherein Q¹ and Z have any of the meanings defined hereinbefore except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means.

The reaction is conveniently carried out in the presence of a suitable base, such as an alkali metal hydride, for example sodium hydride.

15 The reaction is conveniently carried out in the presence of a suitable inert diluent or carrier as defined hereinbefore and at a temperature in the range 10 to 150°C, preferably at or near 125°C.

The compound of Formula X may be prepared using an analogous procedure to that described for the preparation of Formula VIII, except that the 5-fluoro atom is replaced by L¹.

20 Process (h) For the production of those compounds of the Formula I wherein Q¹ contains an amino-substituted (1-6C)alkoxy group (such as 3-piperidinopropoxy, 3-methylaminopropoxy or 3-dimethylaminopropoxy), the reaction of a compound of the Formula I wherein Q¹ contains a halogeno-substituted (1-6C)alkoxy group with an appropriate amine or a heterocycle containing a ring NH group.

25 The reaction is conveniently carried out in the presence of a suitable inert diluent or carrier as defined hereinbefore and at a temperature in the range 10 to 150°C, preferably at or near ambient temperature.

Process (i) For the production of those compounds of the Formula I wherein a heterocycl


group in Q^1 or Q^3 contains an S- or N-oxide the oxidation of a ring N or S atom in compound of the Formula (I). Suitable oxidizing agents include for example, a peracid (such as m-chloroperbenzoic acid) or perphthalic acid. The reaction is conveniently carried out in a suitable inert solvent or diluent (such as dichloromethane) at a suitable temperature (such as – 5 to 50°C).

Process (j) For the production of those compounds wherein Q^2 is a group of the formula 1a wherein:

(i) the group X^2-Q^3 is $CON(R^6)Q^3$ wherein R^6 and Q^3 are as hereinbefore defined, or

(ii) the group X^2-Q^3 is COQ^3 and Q^3 is a nitrogen linked heterocyclcyl group,

10 the coupling of the carboxy substituted quinazoline of the formula XI



XI

or a reactive derivative thereof, with an amine of the formula $NH(R^6)Q^3$ or Q^3H (when Q^3 is a nitrogen containing heterocyclcyl group, for example hompiperidine) as appropriate, wherein R^1 , R^6 , Q^1 , Q^3 , Z , G^1 , G^2 , G^6 and G^7 are as hereinbefore defined except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means. The coupling reaction is conveniently carried out in the presence of a suitable coupling agent, such as a carbodimide, or a suitable peptide coupling agent, for example O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate. The coupling reaction is conveniently carried out in an inert solvent such as 1-methyl-2-pyrrolidinone, preferably in the presence of a suitable base, such as an organic amine, for example di-isopropylethylamine.

By 'reactive derivative' of a compound by the formula XI is meant a derivative of carboxylic acid of formula XI that will react with the amine to give the corresponding amide. Such reactive derivatives include, for example, an acid chloride of the compound of formula XI.

The compound of formula XI may be prepared using process (a) above by reacting a compound of the formula II with an appropriately protected 4-aminobenzoic acid (for example an alkyl or aryl 4-aminobenzoate), followed by removal of the carboxylic acid protecting group using a conventional deprotection technique for the removal of such groups (for

5 example acid hydrolysis in HCl)

Process (k) For the production of those compounds wherein Q^2 is a group of the formula Ia wherein:

(i) X^2Q^3 is OCH_2Q^3 and Q^3 is aryl or heteroaryl, or

(ii) X^2Q^3 is OQ^3 and Q^3 is heteroaryl,

10 the reaction the reaction, optionally in the presence of a base, of a compound of formula I wherein Q^2 is a group of the formula Ia as defined hereinbefore except that the group X^2Q^3 in formula Ia is OH, with a compound of the formula $L^1CH_2Q^3$ or L^1Q^3 wherein L^1 is a displaceable group, and Q^3 is as defined hereinbefore except that any functional group is protected if necessary, and whereafter any protecting group that is present is removed by

15 conventional means. Suitable displaceable groups are, for example halogeno such as chloro, or alkanesulphonyloxy, such as mesyloxy. The reaction is conveniently carried out in an inert solvent such as or a dipolar aprotic solvent for example N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulphoxide, or acetonitrile. The reaction is conveniently carried out in the presence of a suitable base, as described in

20 relation to process (a) above, such as an alkali metal carbonate, for example potassium carbonate. Generally the reaction is suitably performed at a temperature of from -10 to 120°C , conveniently at or near ambient temperature.

The starting material, the compound of formula I wherein Q^2 is a group of the formula Ia wherein X^2Q^3 is OH may be prepared using an analogous process to one of those described

25 herein for the preparation of compounds of the Formula I. For example, Process (a) may be used wherein a compound of the Formula II as hereinbefore described is reacted with an appropriate 4-aminophenol. Compounds of the formula $L^1CH_2Q^3$ or L^1Q^3 are known or may be prepared using conventional procedures.

Process (l) For the production of those compounds of the formula I wherein Q^1 contains an

30 (2-6C)alkanoylamino or substituted (2-6C)alkanoylamino, the acylation of a quinazoline derivative of the formula I wherein Q^1 contains an amino group. A suitable acylating agent is, for example, any agent known in the art for the acylation of amino to acylamino, for example an acyl halide, for example a (2-6C)alkanoyl chloride or bromide, conveniently in the

presence of a suitable base, as defined hereinbefore, an alkanoic acid anhydride or mixed anhydride, for example a (2-6C)alkanoic acid anhydride such as acetic anhydride or the mixed anhydride formed by the reaction of an alkanoic acid and a (1-4C)alkoxycarbonyl halide, for example a (1-4C)alkoxycarbonyl chloride, in the presence of a suitable base as defined


5 hereinbefore. In general the acylation is carried out in a suitable inert solvent or diluent as defined hereinbefore and at a temperature, in the range, for example, -30°C to 120°C , conveniently at or near ambient temperature.

An analogous process may be used to prepare compounds of the formula I containing an (1-6C)alkanesulphonylamino group or substituted alkanesulphonylamino group except the

10 corresponding (1-6C)alkanesulphonylhalide or or substituted alkanesulphonylhalide (for example methanesulphonyl chloride) is used in place of the acylhalide.

Process (m): For the production of those compounds of the Formula I wherein Q^2 is an indole or indazole derivative of the formula Ib, Ic, Id or Ie as hereinbefore defined, the reaction, conveniently in the presence of a suitable base, of a compound of the formula


15 $\text{Q}^4\text{X}^3\text{L}^1$, wherein L^1 is a displaceable group and Q^4 and X^3 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, with a compound of the formula XII,




XII

20 wherein Q^1 and R^1 have any of the meanings defined hereinbefore except that any functional group is protected if necessary, and Q^{2a} is selected from a compound of the formula Ib', Ic', Id' and Ie'


- 96 -




Ib'



Ic'



Id'



Ie'

wherein G^1 , G^4 , G^5 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means.

- 5 Suitable displaceable groups represented by L^1 are as hereinbefore described herein in relation to process (a), such as halogeno, for example chloro.


The reaction is conveniently carried out in the presence of a base. Suitable bases are as hereinbefore described in relation to Process (a), for example sodium hydride.

- 10 The reaction is conveniently performed in a suitable inert solvent, for example or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide or N-methylpyrrolidin-2-one. The reaction is conveniently carried out at a temperature in the range, for example, 10 to 250°C, preferably in the range 20 to 80°C.

- 15 The starting material of the formula XII may be obtained using analogous processes to those described herein, for example using process (a) by reacting a quinazoline of the Formula II as hereinbefore described with the appropriate amino substituted indole or indazole derivative of the compounds of the formulae Ib', Ic', Id' or Ie'.

Compounds of the formula $Q^4X^3L^1$ are known or may be prepared using conventional procedures.

Process (n) For the production of those compounds of the Formula I wherein Q^2 is a group of the formula Ia wherein Q^3 is 1,2,4-oxadiazol-4-yl, or 5-alkyl substituted 1,2,4-oxadiazol-4-yl, the ring closure of the hydroxyamidine of the formula XIII




XIII

- 5 wherein Q^1 , Z, R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with an orthoester of the formula $RCO(OCH_3)$ or carboxylic acid of the formula $RCOOH$, wherein R is hydrogen or (1-6C)alkyl, whereafter any protecting group that is present is removed by conventional means. The reaction is conveniently performed in the presence of an acid, such as formic acid. The
- 10 reaction may be performed at a temperature of from ambient to 120°C , preferably at or near to ambient temperature. Process (n) is especially suitable for those compounds of the formula I wherein X^3 is OCH_2 .

The hydroxyamidine of the formula XIII may be prepared by conventional procedures, for example by reacting a compound of the formula XIV

15




XIV

wherein Q^1 , Z, R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with hydroxylamine, conveniently

in the presence of a base, whereafter any protecting group that is present is removed by conventional means. Suitable bases that may be used in this reaction are as hereinbefore defined in relation to process (a), for example an alkali metal carbonate such a potassium carbonate. The reaction is conveniently carried out in the presence of a suitable inert solvent
 5 such as a polar solvent, for example in aqueous ethanol. The reaction is suitably carried out at a temperature of from 30 to 100°C, for example from 70 to 100°C.


The compound of the formula XIV may be prepared using an analogous process to those described herein for the preparation of compounds of the formula I. For example process (a) may be used by reacting a quinazoline of the formula II as hereinbefore defined
 10 with an aniline of the formula XV:



wherein R¹, X³, G¹, G², G⁶ and G⁷ have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, whereafter any protecting group that is present
 15 is removed by conventional means. Suitable conditions for this reaction are as described for process (a).

Anilines of formula XV are known or may be prepared using conventional techniques, for example using analogous processes to those described above for the preparation of starting materials of the formula Q²NHR¹. For example when X³ is OCH₂ the compound of the
 20 formula XV may be prepared may, for example, be prepared by reacting the starting nitro compound shown in Reaction Scheme 3 in which L¹ is OH with chloroacetonitrile. The nitro group may then be reduced to an amino group by using step 2 of the process in Reaction Scheme 3.

Process (o) For the production of those compounds of the Formula I wherein Q² is a group
 25 of the formula Ia wherein Q³ is 5-amino-1,3,4-oxadiazol-5-yl, the cyclisation, conveniently in the presence of a base, of the compound of the formula XVI




XVI

wherein Q^1 , Z , R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with a cyanogen halide (for example
 5 cyanogen bromide), whereafter any protecting group that is present is removed by conventional means.

Suitable bases that may be used in this reaction are as hereinbefore defined in relation to process (a), for example potassium hydrogen carbonate. The reaction is conveniently carried out in the presence of a suitable inert solvent such as a polar solvent, for example in
 10 aqueous ethanol. The reaction is suitably carried out at a temperature of from -10 to 40°C , for example from -5°C to ambient temperature.

Compounds of the formula XVI may be prepared by conventional procedures, for example by reacting, conveniently in the presence of a base, an ester of formula XVII




XVII

15 wherein R is (1-6C)alkyl or benzyl and Q^1 , Z , R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with hydrazine, whereafter any protecting group that is present is removed by conventional means. Suitable bases that may used are as hereinbefore defined in relation to process (a), for

example an organic base such as pyridine. The reaction is conveniently performed in a suitable inert solvent such as, for example ethanol.

The ester of formula XVII may be prepared may be prepared using an analogous process to those described herein for the preparation of compounds of the formula I. For example when X^3 is OCH_2 compounds of the formula XVII may be prepared according to Reaction Scheme 5.

Reaction Scheme 5



wherein R is (1-6C)alkyl or benzyl and Q¹, Z, R¹, G¹, G², G⁶, G⁷ and each L¹ have any of the


meanings defined hereinbefore, except that any functional group is protected if necessary,

(i) is suitably performed under analogous conditions to those described herein for process (a) hereinbefore

(ii) is suitable performed under analogous conditions to those described for process (k) hereinbefore.

Process (p) For the production of those compounds of the Formula I wherein Q² is a group of the formula Ia wherein Q³ is aryl or heteroaryl and X³ is S, the coupling of the compound of formula XVIII

- 101 -



XVIII

wherein Q^1 , Z , R^1 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with a compound of the formula Q^3SH , wherein Q^3 is aryl or heteroaryl as defined hereinbefore, except that any functional group is
 5 protected if necessary, in the presence of cuprous bromide, and a base, whereafter any protecting group that is present is removed by conventional means.

Suitable bases for the reaction include, for example organic bases such as diazabicyclo[5.4.0]undec-7-ene. The reaction is suitably performed under anhydrous conditions in an inert solvent, for example a hydrocarbon solvent such as toluene. The
 10 reaction is suitably performed at a temperature of from 80 to 150°C, for example from 90 to 130°C.

Compounds of the formula XVIII may be prepared using analogous processes for the preparation of the compounds of the formula I described herein. For example a compound of formula XVIII may be prepared using an analogous process to Process (a) by reacting a
 15 compound of the formula II with an appropriate 4-iodoaniline.

Process (q) For the production of those compounds of the Formula I wherein Q^1 or Q^2 contains an (1-6C)alkylamino, substituted (1-6C)alkylamino group or a nitrogen linked heterocyclyl group (for example when Q^1 is 1-(4-methylpiperazin-1-yl)cyclohexan-4-yl), the reductive amination of an aldehyde or ketone group in a compound of formula 1, with a
 20 (1-6C)alkylamino, substituted (1-6C)alkylamino group or a heterocyclyl group containing an NH group in the presence of a suitable reducing agent. A suitable reducing agent is, for example, a hydride reducing agent, for example an alkali metal aluminium hydride such as lithium aluminium hydride, formic acid or, preferably, an alkali metal borohydride such as sodium borohydride, sodium cyanoborohydride, sodium triethylborohydride, sodium
 25 trimethoxyborohydride and sodium triacetoxyborohydride. The reaction is conveniently

performed in a suitable inert solvent or diluent, for example tetrahydrofuran or diethyl ether for the more powerful reducing agents such as lithium aluminium hydride, and, for example, methylene chloride or a protic solvent such as methanol and ethanol for the less powerful reducing agents such as sodium triacetoxymethylborohydride and sodium cyanoborohydride. The
5 reaction is conveniently performed at a temperature in the range, for example, 10 to 100°C, conveniently at or near ambient temperature.

An analogous reductive amination to that described above may be used to introduce an alkyl or substituted alkyl group onto a primary or secondary amine group in a compound of the formula I by reductive amination with a corresponding ketone in the presence of a suitable
10 reducing agent. For example, for the production of those compounds of the Formula I wherein Q¹ or Q² contains a N-methyl group, the corresponding compound containing an NH group may be reacted with formaldehyde in the presence of a suitable reducing agent as described above.

Process (r) The conversion of one compound of the Formula I into another compound of
15 the Formula I.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above (for example as in process (r)), and as such are
20 included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using
25 concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group.

Particular examples of modifications include the oxidation of alkylthio to alkylsulphinyl or
30 alkylsulphonyl; the substitution of an NH group in Q¹ or Q² by the reaction with an optionally substituted alkyl halide, an optionally substituted alkenyl halide, an optionally substituted alkynyl halide or optionally substituted alkanoyl halide

When a pharmaceutically-acceptable salt of a quinazoline derivative of the Formula I is required, for example an acid-addition salt, it may be obtained by, for example, reaction of said quinazoline derivative with a suitable acid using a conventional procedure.

Biological Assays

- 5 The inhibitory activities of compounds were assessed in non-cell based protein tyrosine kinase assays as well as in cell based proliferation assays before their *in vivo* activity was assessed in Xenograft studies.

a) Protein Tyrosine Kinase phosphorylation Assays

- 10 This test measures the ability of a test compound to inhibit the phosphorylation of a tyrosine containing polypeptide substrate by an enzyme selected from the EGFR kinase, erbB2 kinase and erb4 kinase.

- Recombinant intracellular fragments of EGFR, erbB2 and erbB4 (accession numbers X00588, X03363 and L07868 respectively) were cloned and expressed in the baculovirus/Sf21 system. Lysates were prepared from these cells by treatment with ice-cold
15 lysis buffer (20mM N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid (HEPES) pH7.5, 150mM NaCl, 10% glycerol, 1% Triton X-100, 1.5mM MgCl₂, 1mM ethylene glycol-bis(β-aminoethyl ether) N',N',N',N'-tetraacetic acid (EGTA), plus protease inhibitors and then cleared by centrifugation.

- Constitutive kinase activity of these recombinant proteins was determined by their
20 ability to phosphorylate a synthetic peptide (made up of a random co-polymer of Glutamic Acid, Alanine and Tyrosine in the ratio of 6:3:1). Specifically, Maxisorb™ 96-well immunoplates were coated with synthetic peptide (0.2μg of peptide in a 200μl phosphate buffered saline (PBS) solution and incubated at 4°C overnight). Plates were washed in 50mM HEPES pH 7.4 at room temperature to remove any excess unbound synthetic peptide. EGFR,
25 erbB2 or erbB4 activities were assessed by incubation in peptide coated plates for 20 minutes at room temperature in 100mM HEPES pH 7.4 at room temperature, adenosine trisphosphate (ATP) at Km concentration for the respective enzyme, 10mM MnCl₂, 0.1mM Na₃VO₄, 0.2mM DL-dithiothreitol (DTT), 0.1% Triton X-100 with test compound in DMSO (final concentration of 2.5%). Reactions were terminated by the removal of the liquid components
30 of the assay followed by washing of the plates with PBS-T (phosphate buffered saline with 0.5% Tween 20).

The immobilised phospho-peptide product of the reaction was detected by immunological methods. Firstly, plates were incubated for 90 minutes at room temperature with anti-phosphotyrosine primary antibodies that were raised in the mouse (4G10 from Upstate Biotechnology). Following extensive washing, plates were treated with Horseradish Peroxidase (HRP) conjugated sheep anti-mouse secondary antibody (NXA931 from Amersham) for 60 minutes at room temperature. After further washing, HRP activity in each well of the plate was measured colorimetrically using 22'-Azino-di-[3-ethylbenzthiazoline sulphonate (6)] diammonium salt crystals (ABTS™ from Roche) as a substrate.

Quantification of colour development and thus enzyme activity was achieved by the measurement of absorbance at 405nm on a Molecular Devices ThermoMax microplate reader. Kinase inhibition for a given compound was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of phosphorylation in this assay. The range of phosphorylation was calculated from the positive (vehicle plus ATP) and negative (vehicle minus ATP) control values.

15 **b) KB cell proliferation assay**

This assay measures the ability of a test compound to inhibit the proliferation of KB cells (human naso-pharyngeal carcinoma obtained from the American Type Culture Collection (ATCC)).

KB cells (human naso-pharyngeal carcinoma obtained from the ATCC) were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% foetal calf serum, 2 mM glutamine and non-essential amino acids at 37°C in a 7.5% CO₂ air incubator. Cells were harvested from the stock flasks using Trypsin/ethylenediaminetetraacetic acid (EDTA). Cell density was measured using a haemocytometer and viability was calculated using trypan blue solution before being seeded at a density of 1.25x10³ cells per well of a 96 well plate in DMEM containing 2.5% charcoal stripped serum, 1mM glutamine and non-essential amino acids at 37°C in 7.5% CO₂ and allowed to settle for 4 hours.

Following adhesion to the plate, the cells are treated with or without EGF (final concentration of 1ng/ml) and with or without compound at a range of concentrations in dimethylsulphoxide (DMSO) (1% final) before incubation for 4 days. Following the incubation period, cell numbers were determined by removal of the media by aspiration and incubating with 50µl of 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT)

(stock 5mg/ml) for 2 hours. MTT solution was then removed by aspiration, allowed to air dry and the cells dissolved upon the addition of 100µl of DMSO.

Absorbance of this solubilised cells was read at 540nm to quantify cell biomass.

Inhibition of proliferation was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of proliferation. The range of proliferation was calculated from the positive (vehicle plus EGF) and negative (vehicle minus EGF) control values.

c) H16N-2 cell proliferation assay

This assay measures the ability of a test compound to inhibit heregulin β or EGF driven proliferation of H16N-2 cells. These non-neoplastic epithelial cells respond in a proliferative manner to stimulation with either EGF or heregulin β (Ram, G.R. and Ethier, S.P. (1996) *Cell Growth and Differentiation*, 7, 551-561) were isolated human mammary tissue (Band, V. and Sager, R. Tumour progression in breast cancer. In: J. S. Rhim and A. Dritschilo (eds.), *Neoplastic Transformation in human Cell Culture*, pp 169-178. Clifton, NJ: Humana Press, 1991) and were obtained from the Dana-Farber Cancer Institute, 44 Binney Street, Boston, Massachusetts 02115.

H16N-2 cells were routinely cultured in culture medium (a 1:1 mix of Gibco F12 and Ham's αMEM media containing 1 % foetal calf serum, 10mM HEPES, 1µg/ml Insulin, 12.5ng/ml EGF, 2.8µM Hydrocortisone, 2nM Estradiol 5µM Ascorbic Acid, 10µg/ml Transferrin, 0.1mM Phosphoethanolamine, 15nM Sodium Selenite, 2mM Glutamine, 10nM Tri-iodo-thyrone, 35µg/ml Bovine pituitary Extract and 0.1mM Ethanolamine) at 37°C in a 7.5% CO₂ air incubator. Cells were harvested from the stock flasks using Trypsin/ethylenediaminetetraacetic acid (EDTA). Cell density was measured using a haemocytometer and viability was calculated using trypan blue solution before being seeded at a density of 1.0x10³ cells per well of a 96 well plate in the above media at 37°C in 7.5% CO₂ and allowed to settle for 72 hours.

Following this, the cells were starved of serum for 24 hours upon the addition of starvation medium (a 1:1 mix of Gibco F12 and Ham's αMEM media containing, 10mM HEPES, 2nM Estradiol, 5µM Ascorbic Acid, 10µg/ml Transferrin, 0.1mM Phosphoethanolamine, 15nM Sodium Selenite, 2mM Glutamine, and 0.1mM Ethanolamine) and incubated at 37°C in 7.5% CO₂. The cells were then treated with or without compound at a range of concentrations in dimethylsulphoxide (DMSO) (1% final) for two hours before the

addition of exogenous ligand (at a final concentration of 100ng/ml of heregulin β or 5ng/ml of EGF) and incubation with both ligand and compound for 4 days at 37°C in 7.5% CO₂.

Following the incubation period, cell numbers were determined by removal of the media by aspiration and incubating with 50 μ l of 3-(4,5-Dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) (stock 5mg/ml) for 2 hours. MTT solution was then removed by aspiration, allowed to air dry and the cells dissolved upon the addition of 100 μ l of DMSO.

Absorbance of this solubilised cells was read at 540nm to quantify cell biomass.

Inhibition of proliferation was expressed as an IC₅₀ value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of proliferation.

10 The range of proliferation was calculated from the positive (vehicle plus ligand) and negative (vehicle minus ligand) control values.

d) *In vivo* LoVo Xenograft assay

This assay measures the ability of a test compound to inhibit the growth of a LoVo tumour cell xenograft (colorectal adenocarcinoma obtained from the ATCC) in Female Swiss athymic mice (Alderley Park, *nu/nu* genotype).

Female Swiss athymic (*nu/nu* genotype) mice were bred and maintained in Alderley Park in negative pressure Isolators (PFI Systems Ltd.). Mice were housed in a barrier facility with 12hr light/dark cycles and provided with sterilised food and water *ad libitum*. All procedures were performed on mice of at least 8 weeks of age. LoVo tumour cell xenografts were established in the hind flank of donor mice by sub-cutaneous injections of 1x10⁷ freshly cultured cells in 100 μ l of serum free media per animal. On day 5 post-implant, mice were randomised into groups of 7 prior to the treatment with compound or vehicle control that was administered once daily at 0.1ml/kg body weight. Tumour volume was assessed twice weekly by bilateral Vernier calliper measurement, using the formula (length x width) x $\sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$, where length was the longest diameter across the tumour, and width was the corresponding perpendicular. Growth inhibition from start of treatment was calculated by comparison of the mean changes in tumour volume for the control and treated groups, and statistical significance between the two groups was evaluated using a Students *t* test.

e) *In vivo* BT-474 Xenograft assay

30 This assay measures the ability of a test compound to inhibit the growth of a BT-474 tumour cell xenograft (human mammary carcinoma obtained from Dr Baselga, Laboratorio Recerca Oncologica, Paseo Vall D'Hebron 119-129, Barcelona 08035, Spain) in Female Swiss

athymic mice (Alderley Park, *nu/nu* genotype) (Baselga, J. *et al.* (1998) *Cancer Research*, **58**, 2825-2831).

Female Swiss athymic (*nu/nu* genotype) mice were bred and maintained in Alderley Park in negative pressure Isolators (PFI Systems Ltd.). Mice were housed in a barrier facility with 12hr light/dark cycles and provided with sterilised food and water *ad libitum*. All procedures were performed on mice of at least 8 weeks of age. BT-474 tumour cell xenografts were established in the hind flank of donor mice by sub-cutaneous injections of 1×10^7 freshly cultured cells in 100µl of serum free media with 50% Matrigel per animal. On day 14 post-implant, mice were randomised into groups of 10 prior to the treatment with compound or vehicle control that was administered once daily at 0.1ml/kg body weight. Tumour volume was assessed twice weekly by bilateral Vernier calliper measurement, using the formula $(\text{length} \times \text{width}) \times \sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$, where length was the longest diameter across the tumour, and width was the corresponding perpendicular. Growth inhibition from start of treatment was calculated by comparison of the mean changes in tumour volume for the control and treated groups, and statistical significance between the two groups was evaluated using a Students *t* test.

Although the pharmacological properties of the compounds of the Formula I vary with structural change as expected, in general activity possessed by compounds of the Formula I, may be demonstrated at the following concentrations or doses in one or more of the above tests (a), (b), (c), (d) and (e):-

- Test (a):- IC_{50} in the range, for example, 0.001 - 10 µM;
- Test (b):- IC_{50} in the range, for example, 0.001 - 20 µM;
- Test (c):- IC_{50} in the range, for example, 0.001 - 20 µM;
- Test (d):- activity in the range, for example, 1-200 mg/kg/day;
- Test (e):- activity in the range, for example, 1-200 mg/kg/day;

No physiologically unacceptable toxicity was observed in Test (d) or (e) at the effective dose for compounds tested of the present invention. Accordingly no untoward toxicological effects are expected when a compound of Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore is administered at the dosage ranges defined hereinafter.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a quinazoline derivative of the Formula I, or a

pharmaceutically-acceptable thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible
5 powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing
10 or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

15 The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 0.5 g of active agent (more suitably from 0.5 to 100 mg, for example from 1 to 30 mg) compounded with an
20 appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known
25 principles of medicine.

In using a compound of the Formula I for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.1 mg/kg to 75 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous
30 administration, a dose in the range, for example, 0.1 mg/kg to 30 mg/kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will be used. Oral administration is however

preferred, particularly in tablet form. Typically, unit dosage forms will contain about 0.5 mg to 0.5 g of a compound of this invention.

We have found that the compounds of the present invention possess anti-proliferative properties such as anti-cancer properties that are believed to arise from their erbB family
5 receptor tyrosine kinase inhibitory activity, particularly inhibition of the EGFR and/or erbB2 and/or erbB4 receptor tyrosine kinases, and especially the selective inhibition of erbB2 receptor tyrosine kinases. Accordingly the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by erbB receptor tyrosine kinases, i.e. the compounds may be used to produce a erbB receptor
10 tyrosine kinase inhibitory effect in a warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for the treatment of malignant cells characterised by inhibition of one or more of the erbB family of receptor tyrosine kinases. Particularly the compounds of the invention may be used to produce an anti-proliferative and/or pro-apoptotic and/or anti-invasive effect mediated alone or in part by the inhibition of
15 erbB receptor tyrosine kinases. Particularly, the compounds of the present invention are expected to be useful in the prevention or treatment of those tumours that are sensitive to inhibition of one or more of the erbB receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4 kinase that are involved in the signal transduction steps which drive proliferation and survival of these tumour cells. Accordingly the compounds of the present invention are
20 expected to be useful in the treatment and/or prevention of a number of hyperproliferative disorders by providing an anti-proliferative effect. These disorders include, for example psoriasis, benign prostatic hyperplasia (BPH), atherosclerosis and restenosis and, in particular, erbB2 receptor tyrosine kinase driven tumours. Such benign or malignant tumours may affect any tissue and include non-solid tumours such as leukaemia, multiple myeloma or lymphoma,
25 and also solid tumours, for example bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancers.

According to this aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament.

30 Thus according to this aspect of the invention there is provided the use of a quinazoline derivative of the formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

According to a further feature of this aspect of the invention there is provided a method for producing an anti-proliferative effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, as
5 hereinbefore defined.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

According to a further aspect of the invention there is provided the use of a
10 quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the prevention or treatment of those tumours which are sensitive to inhibition of erbB receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4, that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

15 According to a further feature of this aspect of the invention there is provided a method for the prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB family of receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4, that are involved in the signal transduction steps which lead to the proliferation and/or survival of tumour cells in a warm-blooded animal, such as man, in need of such
20 treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in the prevention or treatment
25 of those tumours which are sensitive to inhibition of one or more of the erbB family of receptor tyrosine kinases, such as EGFR and/or erbB2 and/or erbB4, that are involved in the signal transduction steps which lead to the proliferation and/or survival of tumour cells.

According to a further aspect of the invention there is provided the use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as
30 defined hereinbefore in the manufacture of a medicament for use in providing a EGFR and/or erbB2 and/or erbB4 kinase inhibitory effect.

According to a further feature of this aspect of the invention there is provided a method for providing a EGFR and/or an erbB2 and/or an erbB4 kinase inhibitory effect in a

warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in providing a EGFR and/or an erbB2 and/or an erbB4 kinase inhibitory effect.

According to a further aspect of the invention there is provided the use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in providing a selective erbB2 kinase inhibitory effect.

According to a further feature of this aspect of the invention there is provided a method for providing a selective erbB2 kinase inhibitory effect in a warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in providing a selective erbB2 kinase inhibitory effect.

By "a selective erbB2 kinase inhibitory effect" is meant that the quinazoline derivative of formula I is more potent against erbB2 receptor tyrosine kinase than it is against other kinases. In particular the quinazoline derivative of formula I is more potent against erbB2 receptor kinase than it is against EGFR tyrosine kinase. For example in a cellular assay (such as the H16N-2 assay described herein) the quinazoline derivative of formula I is at least 5 times, preferably at least 10 times more potent against erbB2 receptor tyrosine kinase driven proliferation than it is against EGFR tyrosine kinase driven proliferation, as determined from the relative IC₅₀ values

According to a further aspect of the present invention there is provided the use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of a cancer selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer.

According to a further feature of this aspect of the invention there is provided a

method for treating a cancer selected from selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer in a warm-blooded animal, such as man, in need
5 of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, for use in the treatment of a cancer
10 selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer.

The anti-proliferative treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the quinazoline derivative of the invention, conventional
15 surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents:

(i) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan and nitrosoureas); antimetabolites (for
20 example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside and hydroxyurea; antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and
25 taxotere); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);

(ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifyfene), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example
30 goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of α -reductase such as finasteride;

- (iii) Agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function);
- (iv) other inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erb2 antibody trastuzumab [Herceptin™] and the anti-erb1 antibody cetuximab [C225]) , farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;
- (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody bevacizumab [Avastin™], compounds such as those disclosed in International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);
- (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO00/40529, WO 00/41669, WO01/92224, WO02/04434 and WO02/08213;
- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- (ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor,

approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

Such conjoint treatment may be achieved by way of the simultaneous, sequential or
5 separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention within the dosage range described hereinbefore and the other pharmaceutically-active agent within its approved dosage range.

According to this aspect of the invention there is provided a pharmaceutical product comprising a quinazoline derivative of the formula I as defined hereinbefore and an additional
10 anti-tumour agent as defined hereinbefore for the conjoint treatment of cancer.

Although the compounds of the Formula I are primarily of value as therapeutic agents for use in warm-blooded animals (including man), they are also useful whenever it is required to inhibit the effects of the erbB receptor tyrosine protein kinases. Thus, they are useful as pharmacological standards for use in the development of new biological tests for the
15 evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, and in the search for new pharmacological agents.

The invention will now be illustrated by the following non limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius ($^{\circ}\text{C}$); operations were carried out at room or
20 ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}\text{C}$;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30mmHg) with a bath temperature of up to 60 $^{\circ}\text{C}$;
- (iii) chromatography means flash chromatography on silica gel; thin layer chromatography
25 (TLC) was carried out on silica gel plates;
- (iv) in general, the course of reactions was followed by TLC and / or analytical LC-MS, and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra and/or mass spectral data;
- 30 (vi) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard,

determined at 300 MHz using perdeuterio dimethyl sulphoxide (DMSO- d_6) as solvent unless otherwise indicated; the following abbreviations have been used: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; b, broad;

(viii) chemical symbols have their usual meanings; SI units and symbols are used;

5 (ix) solvent ratios are given in volume:volume (v/v) terms; and

(x) mass spectra were run with an electron energy of 70 electron volts in the chemical ionization (CI) mode using a direct exposure probe; where indicated ionization was effected by electron impact (EI), fast atom bombardment (FAB) or electrospray (ESP); values for m/z are given; generally, only ions which indicate the parent mass are reported; and unless

10 otherwise stated, the mass ion quoted is $(MH)^+$ which refers to the protonated mass ion; reference to M^+ is to the mass ion generated by loss of an electron; and reference to $M-H^+$ is to the mass ion generated by loss of a proton;

(xi) unless stated otherwise compounds containing an asymmetrically substituted carbon and/or sulphur atom have not been resolved;

15 (xii) where a synthesis is described as being analogous to that described in a previous example the amounts used are the millimolar ratio equivalents to those used in the previous example;

(xvi) the following abbreviations have been used:

	THF	tetrahydrofuran;
	DMF	<i>N,N</i> -dimethylformamide;
20	DMA	<i>N,N</i> -dimethylacetamide;
	NMP	1-methyl-2-pyrrolidinone;
	DCM	dichloromethane;
	DMSO	dimethylsulphoxide;
	HATU	O-(7-Azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -Tetramethyluronium
25		Hexafluoro-Phosphate;
	DCE	1,2-dichloroethane;
	<i>m</i> -CPBA	<i>Meta</i> -Chloroperbenzoic acid;
	IPA	Isopropyl alcohol; and
	ether	diethyl ether.

30

xvii) where a synthesis is described as leading to an acid addition salt (e.g. HCl salt), the stoichiometry of the salt was not determined. Unless otherwise stated, all NMR data is reported on free-base material, with isolated salts converted to the free-base form prior to

characterisation by treating a solution of the salt in aqueous methanol with a base such as ammonium hydroxide or sodium bicarbonate thereby precipitating the free base, or by chromatography on silica using an eluant containing a base such as ammonia. Alternatively the free base may be obtained by an extraction method wherein the compound is partitioned
5 between an organic solvent and a basic aqueous medium. The free base is then isolated from the organic medium by, for example evaporation of the organic solvent.

Example 1**4-(1-Benzylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

A solution of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (0.15 g) (reference example 2) and 5-amino-1-benzylindole (0.12 g) in IPA (5 ml) (reference example 7.1) containing HCl in ether (1N, 0.54 ml) was heated at reflux for 1 hr. The solution was cooled and the resulting precipitate filtered to give the title compound as a yellow solid (0.115 g, 46%); NMR spectrum (DMSO-d₆) 2.2 - 3.6 (m, 11H), 5.5 (s, 2H), 6.6 (s, 1H), 7.2 - 7.6 (m, 11H), 8.0 (m, 1H), 8.1 (m, 1H); Mass spectrum MH⁺ 464.

The procedure described above was repeated using the appropriate 4-chloroquinazoline and aniline. Thus were obtained the compounds described below:

Example 1.1**4-(3-Chloro-4-phenoxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-phenoxyaniline (reference example 8) in 51% yield; NMR Spectrum (DMSO-d₆) 1.90 (m, 2H), 2.17 (s, 3H), 2.20 (m, 4H), 2.65 (m, 2H), 4.78 (m, 1H), 6.94 (d, 2H), 7.10 (t, 1H), 7.19 (d, 1H), 7.24 (d, 1H), 7.36 (t, 3H), 7.59 (dd, 1H), 7.73 (t, 1H), 8.33 (d, 1H), 8.56 (s, 1H), 10.19 (s, 1H); Mass Spectrum MH⁺ 461.

Example 1.2**4-(3-Chloro-4-((3-fluorobenzoyloxy)anilino))-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(3-fluorobenzoyloxy)aniline (reference example 8.1) in 70% yield; NMR Spectrum (DMSO-d₆) 1.90 (m, 2H), 2.11 (m, 2H), 2.17 (s, 3H), 2.26 (t, 2H), 2.61 (m, 2H), 4.77 (m, 1H), 5.23 (s, 2H), 7.11-7.34 (m, 6H), 7.39-7.53 (m, 2H), 7.69 (t, 1H), 8.11 (d, 1H), 8.49 (s, 1H), 10.03 (s, 1H); Mass Spectrum MH⁺ 493.

Example 1.3**5-(1-Methylpiperidin-4-yloxy)-4-(4-phenoxyaniline)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 4-phenoxyaniline in 41% yield; NMR spectrum (DMSO-d₆) 2.2 - 2.5 (m, 4H), 2.7 (s, 3H), 3.1 (m, 2H), 3.5 (m, 2H), 5.1 (m, 2H), 7.1 (d, 2H), 7.2 (m, 3H), 7.4 (m, 2H), 7.6 (m, 2H), 7.7 (d, 1H), 7.8 (d, 1H), 8.0 (t, 1H), 8.8 (s, 1H); Mass spectrum MH⁺ 427.

Example 1.4**5-(1-Methylpiperidin-4-yloxy)-4-(4-phenylthioaniline)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 4-(phenylthio)aniline in 57% yield; NMR spectrum (DMSO-d₆) 2.3 - 2.5 (m, 4H), 2.8 (s, 3H), 3.2 (m, 2H), 3.5 (m, 2H), 5.1 (m, 1H), 7.4 (m, 7H), 7.6 (m, 2H), 7.8 (d, 1H), 7.9 (d, 1H), 8.0 (t, 1H), 8.9 (s, 1H); Mass spectrum MH⁺ 443.

Example 1.5**4-(1-Benzenesulphonylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

10 Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 5-amino-1-benzenesulphonylindole (reference example 7) in 26% yield; NMR spectrum (DMSO-d₆) 1.8 (m, 2H), 2.2 (s, 3H), 2.3 (m, 4H), 2.6 (m, 2H), 4.8 (m, 1H), 6.9 (d, 1H), 7.2 (d, 1H), 7.3 (d, 1H), 7.6 (m, 3H), 7.7 (m, 2H), 7.8 (d, 1H), 8.0 (m, 3H), 8.2 (s, 1H), 8.5 (s, 1H); Mass spectrum MH⁺ 514.

15 Example 1.6**4-(3-Chloro-4-(3-pyridyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(pyridyl-3-oxy)aniline (reference example 5) in 17% yield; NMR spectrum (DMSO-d₆, 373K) 2.2-2.5 (m, 2H), 2.5-2.8 (m, 2H), 2.8 (s, 3H), 2.8-3.6 (m, 4H), 5.0 (m, 1H), 7.2-7.3 (d, 1H), 7.3-7.35 (d, 1H), 7.35-7.5 (m, 3H), 7.7-7.75 (dd, 1H), 7.75-7.85 (t, 1H), 8.3 (d, 1H), 8.4 (d, 2H), 8.5 (s, 1H), 9.9 (bs, 1H), 10.3-10.5 (bs, 1H); Mass spectrum MH⁺ 462.

Example 1.7**25 4-(3-Chloro-4-(3-fluorophenoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(3-fluorophenoxy)aniline (reference example 5.1) in 35% yield; NMR spectrum (DMSO-d₆, 373K) 2.2 - 2.5 (m, 2H), 2.5 - 2.8 (m, 2H), 2.8 (s, 3H), 2.8 - 3.6 (m, 4H), 5.0 (m, 1H), 6.75 - 6.8 (dd, 1H), 6.8 - 6.85 (dt, 1H), 6.92 - 7.02 (td, 1H), 7.2 - 7.25 (d, 1H), 7.25 - 7.35 (d, 1H), 7.4 - 7.45 (m, 2H), 7.7 - 7.8 (dd, 1H), 7.8 - 7.85 (t, 1H), 8.3 (s, 1H), 8.6 (s, 1H), 9.9 - 10.1 (bs, 1H), 10.4 - 10.6 (bs, 1H); Mass spectrum MH⁺ 479.

Example 1.8**3-Chloro-4-(2,3-difluorophenoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2,3-difluorophenoxy)aniline (reference example 5.2) in 67% yield; NMR spectrum (DMSO-d₆, 373K) 2.3 - 2.4 (m, 2H), 2.5 - 2.6 (m, 2H), 2.8 (s, 3H), 3.0 - 3.6 (m, 4H), 5.0 (m, 1H), 6.8 - 6.9 (m, 1H), 7.15 - 7.25 (m, 2H), 7.3 - 7.35 (d, 1H), 7.45 - 7.5 (d, 1H), 7.7 - 7.75 (dd, 1H), 7.75 - 7.85 (t, 1H), 8.32 - 8.35 (d, 1H), 8.6 (s, 1H), 10.0 (bs, 1H); Mass spectrum MH⁺ 497.

10 Example 1.9**4-(3-Chloro-4-(2-pyrimidin-yloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-pyrimidin-yloxy)aniline (ex Bionet Ltd) in 90% yield; NMR spectrum (DMSO-d₆, 373K) 2.2 - 2.4 (m, 2H), 2.4 - 2.6 (m, 2H), 2.8 (s, 3H), 3.0 - 3.6 (m, 4H), 5.0 (m, 1H), 7.25 - 7.3 (t, 1H), 7.3 - 7.35 (d, 1H), 7.38 - 7.42 (d, 1H), 7.45 - 7.5 (d, 1H), 7.7 - 7.78 (dd, 1H), 7.78 - 7.82 (t, 1H), 8.25 (d, 1H), 8.6 (s, 1H), 8.66 (s, 1H), 8.68 (s, 1H), 10.0 (bs, 1H); Mass spectrum MH⁺ 463.

Example 1.10**20 4-(3-Chloro-4-(2-thenoyl)anilino)- 5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-thenoyl)aniline (prepared using the method of Example 15(6) of WO 96/15118) in 53% yield; NMR Spectrum (DMSO-d₆) 2.2 - 2.6 (m, 4H), 2.8 (s, 3H), 3.0 - 3.6 (m, 4H), 5.0 - 5.1 (m, 1H), 7.25 (t, 1H), 7.3 - 7.4 (m, 1H), 7.45 (d, 1H), 7.55 (d, 1H), 7.6 (d, 1H), 7.75 - 7.85 (m, 2H), 8.2 (d, 1H), 8.4 (d, 1H), 8.65 (s, 1H); Mass Spectrum MH⁺ 477, 479.

Example 1.11**4-(3-Chloro-4-((1-methyl-1H-imidazol-2-yl)methoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-((1-methyl-1H-imidazol-2-yl)methoxy)aniline (prepared using the method of Example 15(24) of WO 96/15118) in 44% yield; NMR Spectrum (DMSO-d₆) 2.2 - 2.6 (m, 4H), 2.8 (s, 3H), 3.1 - 3.6 (m, 4H), 3.75 (s, 3H), 4.9 - 5.0 (m, 1H), 5.25 (s, 2H), 6.9 (s, 1H), 7.2 (s, 1H), 7.25 (d, 1H), 7.35 (d, 1H), 7.4 (d, 1H), 7.55 (dd, 1H), 7.75 (t, 1H), 8.15 (d, 1H), 8.5 (s, 1H), 9.8 (bs, 1H); Mass Spectrum M-H⁺ 477.

Example 1.12**4-(3-Chloro-4-(N-(2-pyridylmethyl)amino)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-Chloro-4-(N-(2-pyridylmethyl)amino)aniline (reference example 9.1) in 46% yield; NMR Spectrum (DMSO-d₆) 1.90 (m, 2H), 2.15 (m, 2H), 2.20 (s, 3H), 2.30 (m, 2H), 2.61 (m, 2H), 4.50 (d, 2H), 4.79 (tt, 1H), 6.23 (t, 1H), 6.63 (d, 1H), 7.19 (d, 1H), 7.23 - 7.35 (m, 3H), 7.36 (d, 1H), 7.68 (d, 1H), 7.77 (ddd, 1H), 7.96 (d, 1H), 8.44 (s, 1H), 8.57 (d, 1H), 9.90 (s, 1H); Mass Spectrum MH⁺ 475.

Example 1.13**5-(1-Methylpiperidin-4-yloxy)-4-(3-methyl-4-[N-(2-pyridylmethyl)amino]anilino)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-methyl-4-[N-(2-pyridylmethyl)amino]aniline (reference example 5.3) in 53% yield; NMR Spectrum (DMSO-d₆) 2.27 (s, 3H), 2.30 (m, 2H), 2.50 (m, 2H), 2.75 (s, 3H), 3.15 (m, 2H), 3.55 (m, 2H), 4.52 (s, 2H), 5.10 (m, 1H), 6.43 (d, 1H), 7.27 (d, 1H), 7.32 (dd, 1H), 7.37-7.55 (m, 4H), 7.81 (dd, 1H), 7.91 (dd, 1H), 8.59 (d, 1H), 8.75 (s, 1H), 10.3 - 10.6 (d, 1H), 11.15 (bs, 1H); Mass Spectrum MH⁺ 455.

Example 1.14**4-(3-Chloro-4-[N-methyl-N-(2-pyridyl)amino]anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-[N-methyl-N-(2-pyridyl)amino]aniline (reference example 5.4) in

85% yield; NMR Spectrum (DMSO-d₆) 2.25 (m, 2H), 2.40 (m, 2H), 2.79 (s, 3H), 3.20 (m, 2H), 3.40 (s, 3H), 3.57 (m, 2H), 5.00 (m, 1H), 6.24 (d, 1H), 6.66 (dd, 1H), 7.36 (d, 1H), 7.40 - 7.50 (m, 3H), 7.76 (d, 1H), 7.80 (dd, 1H), 8.15 (d, 1H), 8.38 (d, 1H), 8.64 (s, 1H), 9.85 - 10.25 (d, 1H), 10.60 - 10.90 (d, 1H); Mass Spectrum MH⁺ 475.

5 **Example 1.15**

4-(3-Chloro-4-(2-pyridylamino)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-pyridylamino)aniline (reference example 5.5) in 89% yield;

10 NMR Spectrum (DMSO-d₆) 2.22 (m, 2H), 2.45 (m, 2H), 2.77 (s, 3H), 3.20 (m, 2H), 3.55 (m, 2H), 4.90-5.15 (m, 1H), 6.77 (dd, 1H), 6.97 (d, 1H), 7.37 (d, 1H), 7.41 (d, 1H), 7.53 (d, 1H), 7.60 (ddd, 1H), 7.80 (dd, 1H), 8.00 (d, 1H), 8.09 (d, 1H), 8.27 (d, 1H), 8.50 (s, 1H), 8.63 (s, 1H), 9.85 - 10.20 (d, 1H), 10.65 - 11.00 (d, 1H); Mass Spectrum MH⁺ 463.

Example 1.16

15 **5-(1-Methylpiperidin-4-yloxy)-4-(3-methyl-4-(2-pyridylamino)anilino)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-methyl-4-(2-pyridylamino)aniline (reference example 5.6) in 86% yield;

20 NMR Spectrum (DMSO-d₆) 2.27 (m, 5H), 2.50 (m, 2H), 2.77 (s, 3H), 3.20 (m, 2H), 3.55 (m, 2H), 5.01 (m, 1H), 6.66 (dd, 1H), 6.72 (d, 1H), 7.31 (d, 1H), 7.38 (d, 1H), 7.52 (dd, 1H), 7.55 (d, 1H), 7.67 (d, 1H), 7.70 (s, 1H), 7.77 (dd, 1H), 8.03 (d, 1H), 8.26 (s, 1H), 8.54 (s, 1H), 10.01 (s, 1H), 10.80 (s, 1H); Mass Spectrum MH⁺ 441.

Example 1.17

25 **4-(3-Methyl-4-[N-methyl-N-(2-pyridyl)amino]anilino)-5-(1-methylpiperidin-4-yloxy)-quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-methyl-4-[N-methyl-N-(2-pyridyl)amino]aniline (reference example 5.7) in 73% yield; NMR Spectrum (DMSO-d₆) 2.12 (s, 3H), 2.28 (m, 2H), 2.50 (m, 2H), 2.78 (s, 3H), 3.18 (m, 2H), 3.31 (s, 3H), 3.55 (m, 2H), 5.03 (m, 1H), 6.11 (d, 1H), 6.61 (dd, 1H), 7.24 (d, 1H), 7.34 (d, 1H), 7.40 (dd, 1H), 7.41 (d, 1H), 7.79 (dd, 1H), 7.82 (s, 1H), 7.85 (d, 1H), 8.13 (d, 1H), 8.58 (s, 1H), 10.13 (bs, 1H), 10.82 (s, 1H); Mass Spectrum MH⁺ 455.6.

Example 1.18**4-(3-Chloro-4-((3-fluorophenylamino)methyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-((3-fluorophenylamino)methyl)aniline (reference example 10) in 54% yield; NMR Spectrum (DMSO-d₆) 1.93 (m, 2H), 2.15 (m, 2H), 2.19 (s, 3H), 2.27 (m, 2H), 2.66 (m, 2H), 4.33 (d, 2H), 4.80 (tt, 1H), 6.32 (dd, 1H), 6.36 (dd, 1H), 6.60 (t, 1H), 7.09 (ddd, 1H), 7.27 (d, 1H), 7.38 (d, 1H), 7.42 (d, 1H), 7.52 (dd, 1H), 7.75 (dd, 1H), 8.28 (d, 1H), 8.59 (s, 1H); Mass Spectrum MH⁺ 494.5.

Example 1.19**4-(3-Chloro-4-(8-quinolylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(8-quinolylthio)aniline (reference example 11) in 61% yield; Mass spectrum MH⁺ 528.

Example 2**4-(3-Chloro-4-((1-methyl-1H-imidazol-2-yl)thio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

To a solution of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) (100 mg) and 3-chloro-4-((1-methyl-1H-imidazol-2-yl)thio)aniline (obtained as described in example 10 of WO 96/15118) (86 mg) in DMA (4 ml) was added 1.0 M HCl in diethyl ether (1.0 ml) and the resulting slurry heated at 80°C for 1 hour. The reaction mixture was then cooled to room temperature, diluted with acetone and the product filtered off. This solid was purified by column chromatography, eluting with DCM/methanol/880 NH₄OH (100/8/1), to give the title compound (116 mg, 67%); NMR spectrum (CDCl₃) 2.00 (m, 2H), 2.30 (m, 4H), 2.33 (s, 3H), 2.81 (m, 2H), 3.69 (s, 3H), 4.63 (m, 1H), 6.83 (d, 1H), 6.92 (d, 1H), 7.11 (d, 1H), 7.22 (s, 1H), 7.40 (dd, 1H), 7.45 (d, 1H), 7.62 (t, 1H), 8.12 (d, 1H), 8.65 (s, 1H), 10.14 (s, 1H); Mass spectrum MH⁺ 481.

The procedure described above was repeated using the appropriate 4-chloroquinazoline and aniline. Thus were obtained the compounds described below:

Example 2.1**4-(3-Chloro-4-(2-pyridyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-pyridyloxy)aniline (obtained as described in example 15 (12) of WO 96/15118) in 65 % yield; NMR spectrum (DMSO-d₆) 1.85 - 1.97 (m, 2H), 2.12 - 2.34 (m, 7H), 2.63 - 2.75 (m, 2H), 4.80 (m, 1H), 7.08 - 7.15 (m, 2H), 7.27 (d, 1H), 7.43 - 7.49 (m, 2H), 7.62 (dd, 1H), 7.75 (t, 1H), 7.85 - 7.90 (m, 1H), 7.99 - 8.12 (m, 1H), 8.28 (d, 1H), 8.58 (s, 1H), 10.22 (bs, 1H); Mass spectrum MH⁺ 462.

Example 2.2**5-(1-Methylpiperidin-4-yloxy)-4-(3-methyl-4-(2-pyridylmethoxy)anilino)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-methyl-4-(2-pyridylmethoxy)aniline (Example 13 of WO 96/15118) in 48% yield; NMR spectrum (DMSO-d₆) 1.82 - 1.97 (m, 2H), 2.10 - 2.20 (m, 2H), 2.20 (s, 3H), 2.23 - 2.36 (m, 5H), 2.56 - 2.68 (m, 2H), 4.80 (m, 1H), 5.20 (s, 2H), 7.04 (s, 1H), 7.22 (d, 1H), 7.29 - 7.41 (m, 2H), 7.50 - 7.60 (m, 3H), 7.70 (t, 1H), 7.87 (t, 1H), 8.47 (s, 1H), 8.60 (m, 1H), 9.99 (s, 1H); Mass spectrum MH⁺ 456.

Example 3**5-(1-tert-Butoxycarbonylpiperidin-4-yloxy)-4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)quinazoline**

Di-*iso*-propylethylamine (0.234 g), 5-(1-*tert*-butyloxycarbonylpiperidin-4-yloxy)-4-chloroquinazoline (0.218 g) (reference example 2.1) and 3-chloro-4-(3-fluorobenzoyloxy)aniline (0.162 g) (reference example 8.1) in DMA (10 ml) were heated at 90°C for 1.5 hours. The reaction was cooled, concentrated *in vacuo* and the residue purified by chromatography (using DCM to DCM-5% methanol as eluent) to give the title compound as a yellow solid (0.325 g, 94%); NMR spectrum (DMSO-d₆) 1.40 (m, 9H), 1.79 (m, 2H), 2.14 - 2.22 (m, 2H), 3.15 (m, 2H), 3.84 (m, 2H), 4.94 (m, 1H), 5.26 (s, 2H), 7.17 (m, 1H), 7.22 - 7.37 (m, 5H), 7.43 - 7.54 (m, 2H), 7.73 (t, 1H), 8.12 (d, 1H), 8.51 (s, 1H), 9.96 (s, 1H); Mass spectrum MH⁺ 579.

The procedure described above was repeated using the appropriate 4-chloroquinazoline and aniline. Thus were obtained the compounds described below:

Example 3.1**4-(3-Chloro-4-(1,5-dimethylpyrazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-(2-chloro-4-aminophenoxyethyl)-1,5-dimethylpyrazole (reference example 8.3) in IPA in 31% yield; NMR spectrum (DMSO-d₆) 2.0 (m, 2H), 2.1 - 2.2 (m, 5H), 2.2 - 2.3 (m, 5H), 2.6 (m, 2H), 3.6 (s, 3H), 4.8 (m, 1H), 5.0 (s, 2H), 6.1 (s, 1H), 7.3 (d, 1H), 7.3 (m, 2H), 7.5 (dd, 1H), 7.7 (t, 1H), 8.1 (d, 1H), 8.5 (s, 1H), 10.0 (s, 1H); Mass spectrum MH⁺ 493.

Example 3.2**4-(3-Chloro-4-(1-methylpyrazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-(2-chloro-4-aminophenoxyethyl)-1-methylpyrazole (reference example 8.4) in IPA in 35% yield; NMR spectrum (DMSO-d₆) 1.9 (m, 2H), 2.1 (m, 2H), 2.2 (s, 3H), 2.3 (m, 3H), 2.6 (m, 2H), 3.8 (s, 3H), 4.8 (m, 1H), 5.1 (s, 2H), 6.3 (d, 1H), 7.2 (d, 1H), 7.3 (m, 2H), 7.5 (m, 1H), 7.7 (m, 2H), 8.1 (d, 1H), 8.5 (s, 1H), 10.0 (bs, 1H); Mass spectrum MH⁺ 479.

Example 3.3**4-(3-Chloro-4-((3-methylisoxazol-5-yl)methoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 5-(2-chloro-4-aminophenoxyethyl)-3-methylisoxazole (reference example 9.5) in IPA in 63% yield; NMR spectrum (DMSO-d₆) 1.9 (m, 2H), 2.1 (m, 2H), 2.2 (s, 3H), 2.2 (s, 3H), 2.26 (m, 2H), 2.6 (m, 2H), 3.8 (s, 3H), 4.8 (m, 1H), 5.3 (s, 2H), 6.5 (s, 1H), 7.2 (d, 1H), 7.3 (d, 2H), 7.5 (m, 1H), 7.7 (t, 1H), 8.1 (d, 1H), 8.5 (s, 1H), 10.0 (bs, 1H); Mass spectrum MH⁺ 480.

Example 4**4-(4-(Azepan-1-ylcarbonyl)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Sodium hydride (60% dispersion in oil, 20 mg) was added to 4-hydroxy-1-methylpiperidine (46 mg) in DMA (0.5 ml) and the reaction stirred at 40°C for 30 minutes. 4-(4-(Azepan-1-ylcarbonyl)-3-chloroanilino)-5-fluoroquinazoline hydrochloride (reference example 4) (43.5 mg) was added and the reaction heated at 130°C for 3 hours. The reaction was quenched by addition of a few drops of water, then concentrated *in vacuo*. The residue

was purified by chromatography (using DCM-2% methanolic ammonia to DCM-5% methanolic ammonia as eluent) to give the title compound as a white solid (66 mg, 67%); NMR spectrum (DMSO-d₆) 1.55 (bs, 6H), 1.73 (bs, 2H), 1.88 - 1.97 (m, 2H), 2.14 - 2.22 (m, 5H), 2.26 (t, 2H), 2.64 - 2.74 (m, 2H), 3.23 (s, 2H), 3.58 (d, 2H), 4.80 (m, 1H), 7.27 (d, 1H), 7.37 (d, 1H), 7.39 (d, 1H), 7.64 (dd, 1H), 7.76 (t, 1H), 8.27 (d, 1H), 8.62 (s, 1H), 10.28 (s, 1H); Mass spectrum MH⁺ 494.

The procedure described above was repeated using the appropriate 5-fluoroquinazoline and alcohol. Thus was obtained the compound described below:

Example 4.1

10 4-(1-(3-Fluorobenzyl)indazol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(1-(3-fluorobenzyl)indazol-5-ylamino)-5-fluoroquinazoline hydrochloride (reference example 4.1) with 4-hydroxy-1-methylpiperidine in 41 % yield; NMR spectrum (DMSO-d₆) 1.87 - 1.97 (m, 2H), 2.02 - 2.30 (m, 5H), 2.27 (m, 2H), 2.58 - 2.67 (m, 2H), 4.82 (m, 1H), 5.68 (s, 2H), 7.02 - 7.12 (m, 3H), 7.23 (d, 1H), 7.31 - 7.38 (m, 2H), 7.54 (dd, 1H), 7.68 - 7.77 (m, 2H), 8.15 (s, 1H), 8.37 (d, 1H), 8.48 (s, 1H), 10.19 (s, 1H); Mass spectrum MH⁺ 483.

Example 4.2

4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) and tetrahydropyran-4-ol in 53% yield; NMR spectrum 1.80 - 1.95 (m, 2H), 2.20 (m, 2H), 3.55 (dt, 1H), 3.90 (dt, 1H), 4.95 (m, 1H), 5.24 (s, 2H), 7.16 (dt, 1H), 7.20 - 7.35 (m, 5H), 7.40 - 7.53 (m, 2H), 7.70 (t, 1H), 8.17 (d, 1H), 8.50 (s, 1H), 10.03 (bs, 1H); Mass Spectrum M⁺ 480.

Example 4.3

25 4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(1-methylpyrrolidin-3-yloxy)quinazoline

Obtained from 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) and 3-hydroxy-1-methylpyrrolidine in 50% yield; NMR spectrum 1.92 (m, 1H), 2.22 (m, 1H), 2.38 (m + s, 4H), 2.50 (m, 1H), 3.00 (dt, 1H), 3.17 (d, 1H), 5.23 (s, 2H), 5.32 (m, 1H), 7.15 (m, 2H), 7.23 - 7.37 (m, 4H), 7.43 (m, 1H), 7.70 (t, 1H), 7.77 (dd, 1H), 8.08 (d, 1H), 8.52 (s, 1H), 10.40 (bs, 1H); Mass Spectrum M⁺ 479.

Example 4.4**4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(tetrahydrofuran-3-yloxy)quinazoline**

Obtained from 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) and 3-hydroxytetrahydrofuran in 64% yield; Mass Spectrum M^+ 466.

5 Example 5**4-(4-(2-Bromobenzoyloxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Potassium carbonate (290 mg) was added to a mixture of 2-bromobenzylchloride (128 mg) and 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.2) in DMF (15 ml). The mixture was stirred at room temperature overnight. The reaction mixture was then concentrated *in vacuo* and the residue purified by chromatography using 0-4% methanolic ammonia in DCM as eluent to afford the title compound as a pink solid (90 mg, 31%); NMR Spectrum (DMSO- d_6) 1.90 (m, 2H), 2.11 (m, 2H), 2.17 (s, 3H), 2.26 (m, 2H), 2.61 (m, 2H), 4.78 (m, 1H), 5.21 (s, 2H), 7.20 - 7.34 (m, 4H), 7.42 - 7.54 (m, 2H), 7.61 - 7.73 (m, 3H), 8.13 (d, 1H), 8.50 (s, 1H), 10.05 (bs, 1H); Mass spectrum MH^+ 555.

The procedure described above was repeated using the appropriate 4-hydroxyanilinoquinazoline and alkyl halide or mesylate. Thus were obtained the compounds described below:

Example 5.1**4-(3-Chloro-4-(1,2,5-thiadiazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.2) and 3-bromomethyl-1,2,5-thiadiazole (obtained as described in *J. Heterocycl. Chem.* 1984, 21, 1157-60); NMR Spectrum (DMSO- d_6) 1.8 - 2.0 (m, 2H), 2.05 - 2.4 (m, 7H), 2.5 - 2.7 (m, 2H), 4.7 - 4.8 (m, 1H), 5.55 (s, 2H), 7.2 (d, 1H), 7.35 (d, 2H), 7.5 (d, 1H), 7.7 (t, 1H), 8.15 (s, 1H), 8.5 (s, 1H), 8.95 (s, 1H); Mass Spectrum MH^+ 483.

Example 5.2**4-(4-Benzoyloxy-3-fluoroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and benzyl chloride in 22% yield; Mass Spectrum MH^+ 459.

Example 5.3**4-(3-Fluoro-4-(2-fluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and 2-fluorobenzyl chloride in 37%
5 yield; Mass Spectrum MH^+ 477.

Example 5.4**4-(4-(2,6-Difluorobenzoyloxy)-3-fluoroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and 2,6-difluorobenzyl chloride in
10 27% yield; Mass Spectrum $M-H^+$ 493.

Example 5.5**4-(4-(2-Cyanobenzoyloxy)-3-fluoroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and 2-cyanobenzyl chloride in 12%
15 yield; Mass Spectrum MH^+ 484.

Example 5.6**4-(3-Fluoro-4-(2-pyridylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and 2-picoyl chloride
20 hydrochloride in 11% yield; Mass Spectrum MH^+ 460.

Example 5.7**4-(3-Fluoro-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.3) and 5-methyl-3-chloromethylisoxazole in 38% yield; Mass Spectrum MH^+ 464.
25

Example 5.8**4-(3-Chloro-4-(3,4-difluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.2) and 3,4-difluorobenzyl chloride in
30 53% yield; Mass Spectrum MH^+ 511.

Example 5.10**4-(3-Chloro-4-(isoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride (reference example 4.2) and 3-chloromethylisoxazole (reference example 31) in 71% yield; Mass Spectrum MH^+ 466.

Example 5.11**4-(3-Chloro-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(tetrahydropyran-4-yloxy)quinazoline (reference example 42) and 3-chloromethyl-5-methylisoxazole in 44% yield; NMR Spectrum (DMSO- d_6) 1.8 - 1.94 (m, 2H), 2.14 - 2.25 (m, 2H), 2.41 (s, 3H), 3.54 (dt, 2H), 3.91 (dt, 2H), 4.91 - 5.01 (m, 1H), 5.25 (s, 2H), 6.34 (s, 1H), 7.23 - 7.26 (m, 3H), 7.51 (dd, 1H), 7.72 (t, 1H), 8.16 (d, 1H), 8.51 (s, 1H), 10.04 (s, 1H); Mass spectrum MH^+ 467.

Example 5.12**4-(3-Chloro-4-(2-pyrazinylmethoxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(tetrahydropyran-4-yloxy)quinazoline (reference example 42) and 2-chloromethylpyrazine (obtained as described in Synthesis, 1984, 676) in 4% yield; NMR Spectrum (DMSO- d_6) 1.80 - 1.93 (m, 2H), 2.14 - 2.27 (m, 2H), 3.54 (t, 2H), 3.85 - 3.95 (m, 2H), 4.90 - 5.01 (m, 1H), 5.37 (s, 2H), 7.24 - 7.36 (m, 4H), 7.52 (dd, 1H), 7.72 (t, 1H), 8.18 (d, 1H), 8.51 (s, 1H), 8.62 - 8.69 (m, 1H), 8.84 (s, 1H), 10.05 (s, 1H); Mass spectrum MH^+ 464.

Example 5.13**4-(3-Chloro-4-(5-methylisoxazol-3-yl)anilino)-5-(tetrahydrofuran-3-yloxy)quinazoline**

Obtained by reacting 4-(3-chloro-4-hydroxyanilino)-5-(tetrahydrofuran-3-yloxy)quinazoline (reference example 42.1) and 3-chloromethyl-5-methylisoxazole in 85% yield; NMR Spectrum (DMSO- d_6) 2.12 - 2.23 (m, 1H), 2.28 - 2.41 (m, 1H), 2.41 (s, 3H), 3.79 - 3.99 (m, 3H), 4.19 (d, 1H), 5.24 (s, 2H), 5.46 (t, 1H), 6.33 (s, 1H), 7.19 (d, 1H), 7.29 (d, 1H), 7.35 (d, 1H), 7.57 (dd, 1H), 7.74 (t, 1H), 8.19 (d, 1H), 8.53 (s, 1H), 10.05 (s, 1H); Mass spectrum MH^+ 453.

30

35

Example 5.14**4-(3-Chloro-4-(2-morpholinothiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(4-chloromethylthiazol-2-yl)morpholine (obtained as described in Example 9 of US Patent US 5,455,351) and 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) in 41% yield; NMR spectrum (DMSO-d₆) 1.85 - 1.97 (m, 2H), 2.10 - 2.20 (m, 2H), 2.18 (s, 3H), 2.23 - 2.34 (m, 2H), 2.58 - 2.68 (m, 2H), 3.38 (m, 4H), 3.71 (m, 4H), 4.80 (m, 1H), 5.05 (s, 2H), 6.91 (s, 1H), 7.23 (d, 1H), 7.32 (d, 1H), 7.34 (d, 1H), 7.52 (dd, 1H), 7.73 (dd, 1H), 8.11 (d, 1H), 8.52 (s, 1H), 10.06 (s, 1H); Mass spectrum MH⁺ 567.

Example 5.15**4-(4-Benzoyloxy-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from benzyl chloride and 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) in 48% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.8 (m, 2H), 4.6 (m, 1H) 5.1 (s, 2H), 6.9 (t, 2H), 7.3 - 7.5 (m, 8H), 7.6 (t, 1H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 455.

Example 5.16**4-(4-(2-Fluorobenzoyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 2-fluorobenzyl chloride in 74% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.8 (m, 2H), 4.6 (m, 1H) 5.2 (s, 2H), 6.9 (d, 1H), 6.9 (d, 1H), 7.1 (t, 1H), 7.2 (t, 1H), 7.3 (m, 1H), 7.4 - 7.6 (m, 5H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 473.

Example 5.17**4-(4-(2,6-Difluorobenzoyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 2,6-difluorobenzyl chloride in 72% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.3 (m, 5H), 2.4 (m, 5H), 2.8 (m, 2H), 4.6 (m, 1H) 5.1 (s, 2H), 6.9 (m, 3H), 7.0 (d, 1H), 7.3 (m, 1H), 7.4 (m, 2H), 7.5 (dd, 1H), 7.6 (t, 1H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 491.

Example 5.18**4-(3-Methyl-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 3-chloromethyl-5-methylisoxazole in 70% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.4 (s, 3H), 2.8 (m, 2H), 4.6 (m, 1H) 5.1 (s, 2H), 6.1 (s, 1H), 6.9 (m, 2H), 7.5 (m, 2H), 7.6 (t, 1H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 460.

Example 5.19**5-(1-Methylpiperidin-4-yloxy)-4-(3-methyl-4-(thiazol-4-ylmethoxy)anilino)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 4-chloromethylthiazole in 46% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.8 (m, 2H), 4.6 (m, 1H) 5.3 (s, 2H), 6.9 (d, 1H), 6.9 (d, 1H), 7.4 - 7.6 (m, 5H), 8.6 (s, 1H), 8.8 (d, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 462.

Example 5.20**4-(4-(2-Cyanobenzoyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 2-chloromethyl benzonitrile in 33% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.8 (m, 2H), 4.7 (m, 1H) 5.3 (s, 2H), 6.9 (m, 2H), 7.4 (m, 2H), 7.5 (m, 2H), 7.6 (m, 2H), 7.7 (m, 2H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 480.

Example 5.21**4-(4-(3-Fluorobenzoyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-(3-methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 38) with 3-fluorobenzyl chloride in 43% yield; NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 - 2.4 (m, 10H), 2.8 (m, 2H), 4.6 (m, 1H) 5.1 (s, 2H), 6.9 (m, 2H), 7.0 (m, 1H), 7.2 (m, 2H), 7.3 - 7.5 (m, 4H), 7.6 (t, 1H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 473.

Example 5.22**4-(3-Fluoro-4-(3-fluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 3-fluorobenzyl chloride and 4-(3-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.3) in 46% yield; NMR spectrum (DMSO-d₆) 2.0 (m, 2H), 2.2 (m, 5H), 2.3 (m, 2H), 2.7 (m, 2H), 3.2 (m, 2H), 4.8 (m, 1H) 5.3 (s, 2H), 7.2 - 7.4 (m, 8H), 7.5 (m, 1H), 7.7 (m, 1H), 8.0 (m, 1H), 8.6 (s, 1H), 10.1 (s, 1H); Mass spectrum MH⁺ 477.

Example 5.23**4-(3-Chloro-4-(2-methyloxazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(3-Chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) and methanesulphonic acid 2-methyloxazol-4-ylmethyl ester (reference example 48) in 11% yield; NMR Spectrum (DMSO-d₆) 1.90 (m, 2H), 2.12 (m, 2H), 2.17 (s, 3H), 2.25 (m, 2H), 2.40 (s, 3H), 2.60 (m, 2H), 4.79 (m, 1H), 5.02 (s, 2H), 7.20 (d, 1H), 7.32 (d, 1H), 7.35 (d, 1H), 7.51 (dd, 1H), 7.70 (t, 1H), 8.05 (s, 1H), 8.10 (d, 1H), 8.50 (s, 1H), 10.04 (s, 1H); Mass spectrum M-H⁺ 480.

Example 5.24**4-(5-Chloro-2-fluoro-4-(2-pyridylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(5-chloro-2-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.7) and 2-picolyl chloride hydrochloride in 28% yield; NMR spectrum (DMSO-d₆, 373K), 1.8 - 1.95 (m, 2H), 2.1 - 2.2 (m, 2H), 2.2 (s, 3H), 2.2 - 2.3 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 - 4.85 (m, 1H), 5.3 (s, 2H), 7.25 (d, 1H), 7.3 - 7.4 (m, 3H), 7.6 (d, 1H), 7.7 - 7.8 (t, 1H), 7.85 - 7.95 (m, 1H), 8.6 (s, 1H), 8.6 - 8.7 (m, 2H), 10.0 (bs, 1H); Mass Spectrum MH⁺ 495.

Example 6**4-(3-Chloro-4-(decahydroquinolin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

A suspension of 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24) (74.8 mg) in NMP (1.7 ml) was treated with a solution of HATU (76 mg) in NMP (1.7 ml) followed by a solution of di-isopropylethylamine (43 mg) in NMP (1.7 ml) and the mixture stirred at room temperature for 30

minutes. To this was added decahydroquinoline (93 mg) and the mixture stirred at room temperature overnight. The solvent was removed *in vacuo* to give a gum that was treated with aqueous sodium hydrogen carbonate and extracted with DCM. The DCM solution was evaporated *in vacuo* to give an oil which was purified by column chromatography, eluting with DCM/methanol/880 NH₄OH (100/8/1), to give the title compound (37 mg, 42%); Mass spectrum MH⁺ 534.

The procedure described above was repeated using the appropriate amine and acid. Thus were obtained the compounds described below:

Example 6.1

10 4-(3-Chloro-4-(decahydroisoquinolin-2-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24) and decahydroisoquinoline in 35% yield; Mass spectrum MH⁺ 534.

15 Example 6.2

4-(3-Chloro-4-(3-methylpiperidin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24) and 3-methylpiperidine in 59% yield; NMR spectrum (DMSO-d₆, 373K) 0.90 (m, 3H), 1.25 (m, 1H), 1.50 (m, 1H), 1.68 (m, 2H), 1.84 (m, 1H), 2.00 (m, 2H), 2.20 (m, 2H), 2.25 (s, 3H), 2.35 (m, 2H), 2.75 (m, 3H), 2.95 (m, 3H), 4.82 (m, 1H), 7.27 (d, 1H), 7.37 (d, 1H), 7.43 (d, 1H), 7.70 (d, 1H), 7.78 (t, 1H), 8.23 (d, 1H), 8.64 (s, 1H), 10.25 (s, 1H); Mass spectrum MH⁺ 494.

Example 6.3

25 4-(3-Chloro-4-(4-methylpiperidin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24) and 4-methylpiperidine in 71% yield; NMR spectrum (DMSO-d₆, 373K) 1.00 (d, 3H), 1.16 (m, 2H), 1.70 (m, 3H), 2.00 (m, 2H), 2.20 (m, 2H), 2.28 (s, 3H), 2.36 (m, 2H), 2.75 (m, 2H), 2.95 (m, 4H), 4.82 (m, 1H), 7.28 (d, 1H), 7.39 (d, 1H), 7.42 (d, 1H), 7.70 (d, 1H), 7.78 (t, 1H), 8.21 (d, 1H), 8.65 (s, 1H), 10.24 (s, 1H); Mass spectrum MH⁺ 494.

Example 6.4**4-(3-Ethynyl-4-(decahydroquinolin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 2-ethynyl-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24.1) and decahydroquinoline in 34% yield; Mass spectrum MH^+ 524.

Example 6.5**4-(3-Ethynyl-4-(homopiperidin-1-ylcarbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 2-ethynyl-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride (reference example 24.1) and homopiperidine in 51% yield; Mass spectrum MH^+ 484.

Example 7**4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline**

Trifluoroacetic acid (0.5 ml) was added to a solution of 5-(1-*tert*-butoxycarbonylpiperidine-4-yloxy)-4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)quinazoline (0.31 g) (example 3) in DCM (2 ml) and the solution stirred for 1 hour. The reaction was concentrated *in vacuo* and the residue triturated with *conc.* aq. ammonium hydroxide and filtered to give the title compound as a white solid (0.196 g, 87%); NMR spectrum (DMSO- d_6) 1.76 - 1.92 (m, 2H), 2.24 - 2.36 (m, 2H), 2.77 - 2.90 (m, 2H), 3.15 - 3.28 (m, 2H), 4.65 - 4.77 (m, 1H), 5.16 (s, 2H), 6.90 - 7.08 (m, 3H), 7.18 - 7.29 (m, 2H), 7.32 - 7.66 (m, 4H), 7.92 (t, 1H), 8.63 (s, 1H), 10.05 (bs, 1H); Mass spectrum MH^+ 479.

Example 8**4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(1-propylpiperidin-4-yloxy)quinazoline**

4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (96 mg) (example 7), propyl bromide (27 mg) and potassium carbonate (0.11 g) in DMA (2 ml) were stirred at room temperature overnight. The reaction was filtered and concentrated *in vacuo*. The residue was purified by chromatography (using DCM-5% methanol as eluent) to give the title compound as a white solid after trituration with ether (61 mg, 59%); NMR spectrum (CDCl₃) 0.92 (t, 3H), 1.45 - 1.61 (m, 2H), 1.92 - 2.06 (m, 2H), 2.20 - 2.39 (m, 6H), 2.79 - 2.92 (m, 2H), 4.64 (m, 1H), 5.16 (s, 2H), 6.87 - 7.06 (m, 3H), 7.17 - 7.29 (m, 2H), 7.31 - 7.41 (m,

1H), 7.44 (d, 1H), 7.52 (dd, 1H) 7.61 (t, 1H), 7.90 (d, 1H), 8.62 (s, 1H), 10.00 (bs, 1H); Mass spectrum MH^+ 521.

The procedure described above was repeated using the appropriate amine and alkyl bromide or chloride. Thus were obtained the compounds described below:

5 Example 8.1

4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(1-allylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (example 7) with allyl bromide in 75% yield; NMR spectrum ($CDCl_3$) 1.94 - 2.09 (m, 2H), 2.21 - 2.43 (m, 4H), 2.80 - 2.93 (m, 2H), 3.04 (d, 2H), 4.68 (m, 1H), 5.14 - 5.25 (m, 4H), 5.89 (m, 1H), 6.88 - 7.07 (m, 3H), 7.17 - 7.27, (m, 2H), 7.30 - 7.40 (m, 1H); 7.46 (d, 1H), 7.53 (dd, 1H), 7.61 (t, 1H), 7.90 (d, 1H), 8.62 (s, 1H), 9.98 (bs, 1H); Mass spectrum MH^+ 519.

Example 8.2

4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(1-(2-propynyl)piperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (example 7) with 3-bromoprop-1-yne in 65% yield; NMR spectrum ($CDCl_3$) 1.97 - 2.12 (m, 2H), 2.18 (t, 1H), 2.25 - 2.34 (m, 2H), 2.50 - 2.62 (m, 2H), 2.86 - 2.97 (m, 2H), 3.39 (d, 2H), 4.68 (m, 1H), 5.15 (s, 2H), 6.90 - 7.07 (m, 3H), 7.19 - 7.29 (m, 2H), 7.31 - 7.40 (m, 1H), 7.46 (d, 1H), 7.51 (dd, 1H), 7.63 (t, 1H), 7.91 (d, 1H), 8.62 (s, 1H), 9.94 (bs, 1H); Mass spectrum MH^+ 517.

Example 8.3

4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(1-(2-methoxyethyl)piperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (example 7) with 2-bromoethyl methylether in 44% yield; NMR spectrum ($CDCl_3$) 1.96 - 2.10 (m, 2H), 2.23 - 2.33 (m, 2H), 2.35 - 2.47 (m, 2H), 2.64 (t, 2H), 2.87 - 2.97 (m, 2H), 3.36 (s, 3H), 3.52 (t, 2H), 4.66 (m, 1H), 5.16 (s, 2H), 6.88 - 7.08 (m, 3H), 7.18 - 7.29 (m, 2H), 7.31 - 7.40 (m, 1H), 7.44 (d, 1H), 7.52 (dd, 1H), 7.61 (t, 1H), 7.92 (d, 1H), 8.34 (s, 1H), 9.98 (bs, 1H); Mass spectrum MH^+ 537.

Example 8.4**2-(4-(4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)quinazolin-5-yloxy)piperidin-1-yl)acetone.**

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (example 7) with chloroacetone in 54 % yield; NMR spectrum (CDCl₃)
5 1.98 - 2.12 (m, 2H), 2.17, (s, 3H), 2.23 - 2.36 (m, 2H), 2.44 - 2.55 (m, 2H), 2.82 - 2.93 (m, 2H), 3.28 (s, 3H), 4.67 (m, 1H), 5.57 (s, 2H), 6.90 (d, 1H), 6.94 - 7.07 (m, 2H), 7.19 - 7.28 (m, 2H), 7.32 - 7.41 (m, 1H), 7.46 (d, 1H), 7.51 (dd, 1H), 7.61 (t, 1H), 7.92 (d, 1H), 8.63 (s, 1H), 9.96 (s, 1H); Mass spectrum MH⁺ 535.

Example 8.5**10 Methyl 2-(4-(4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)quinazolin-5-yloxy)piperidin-1-yl)acetate**

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (example 7) with methyl bromoacetate in 38% yield; NMR spectrum (CDCl₃)
15 2.00 - 2.15 (m, 2H), 2.24 - 2.35 (m, 2H), 2.54 - 2.65 (m, 2H), 2.91 - 3.02 (m, 2H), 3.31 (s, 2H), (s, 3H), 4.66 (m, 1H), 5.16 (s, 2H), 6.91 (d, 1H), 6.97 (d, 1H), 6.90 - 7.08 (m, 1H), 7.18 - 7.27 (m, 2H), 7.31 - 7.39 (m, 1H), 7.46 (d, 1H), 7.50 (dd, 1H), 7.62 (t, 1H), 7.93 (d, 1H), 8.63 (s, 1H), 9.96 (bs, 1H), Mass spectrum MH⁺ 551.

Example 9**20 4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(1-(methanesulphonyl)piperidin-4-yloxy)quinazoline**

4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (48 mg) (example 7), methanesulphonyl chloride (12.6 mg) and di-*iso*-propylethylamine (19.4 mg) in DCM (2 ml) were stirred at room temperature overnight. The reaction was concentrated *in vacuo*. The residue was purified by chromatography (using DCM to DCM-5% methanol as
25 eluent) to give the title compound as a white solid (38 mg, 68%); NMR spectrum (DMSO-d₆) 1.93 - 2.06 (m, 2H), 2.21 - 2.33 (m, 2H), 2.84 (s, 3H), 3.05 - 3.15 (m, 2H), 3.45 - 3.55 (m, 2H), 4.92 (m, 1H), 5.26 (s, 2H), 7.13 - 7.20 (m, 1H), 7.22 - 7.38 (m, 5H), 7.41 - 7.56 (m, 2H), 7.73 (t, 3H), 8.17 (s, 1H), 8.51 (s, 1H), 9.90 (bs, 1H); Mass spectrum MH⁺ 557.

The procedure described above was repeated using the appropriate amine and alkyl
30 bromide or chloride. Thus was obtained the compound described below:

Example 9.1**2-(4-(4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)quinazolin-5-yloxy)piperidin-1-yl)acetamide**

Obtained by reacting 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-(piperidin-4-yloxy)quinazoline (**example 7**) with 2-chloroacetamide in 75% yield; NMR spectrum (DMSO-d₆) 1.90 - 2.02 (m, 2H), 2.15 - 2.25 (m, 2H), 2.37 - 2.46 (m, 2H), 2.72 - 2.80 (m, 2H), 2.89 (s, 2H), 4.81 (m, 1H), 5.27 (s, 2H), 7.09 (bs, 1H), 7.16 (m, 1H), 7.22 - 7.36 (m, 6H), 7.42 - 7.52 (m, 2H), 7.72 (t, 1H), 8.17 (d, 1H), 8.52 (s, 1H), 10.10 (bs, 1H); Mass spectrum MH⁺ 536.

Example 10**4-(1-(5-Methylisoxazol-3-ylmethyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Sodium hydride (60% dispersion in mineral oil, 25 mg) was added to a stirred solution of 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**reference example 4.4**) (0.2 g) in DMF (2 ml). The solution was stirred at room temperature for 30 mins, then 5-methyl-3-isoxazol-3-ylmethyl chloride (85 mg) added. The reaction was stirred for a further 3 hours, then poured into water. The resulting solid precipitate was filtered, dried and triturated with ether to give the title compound as a pale green solid (42 mg, 17%); NMR spectrum (DMSO-d₆) 1.8 - 2.0 (m, 2H), 2.15 (s + m, 5H), 2.28 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.43 (s, 2H), 6.00 (s, 1H), 6.50 (d, 1H), 7.10 (d, 1H), 7.30 (m, 2H), 7.49 (m, 2H), 7.67 (t, 1H), 8.07 (dd, 1H), 8.42 (s, 1H), 10.10 (bs, 1H); Mass Spectrum MH⁺ 469.

The procedure described above was repeated using the appropriate indole and alkyl bromide or chloride. Thus were obtained the compounds described below:

Example 10.1**4-(1-(2,6-Difluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**reference example 4.4**) and 2,6-difluorobenzyl bromide in 89% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.12 (s + m, 5H), 2.28 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.44 (s, 2H), 6.45 (d, 1H), 7.10 - 7.20 (m, 3H), 7.23 - 7.50 (m, 5H), 7.67 (t, 1H), 8.04 (d, 1H), 8.40 (s, 1H), 10.09 (bs, 1H); Mass Spectrum MH⁺ 500.

Example 10.2**4-(1-(2-Cyanobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2-cyanobenzyl bromide in 50% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.61 (m, 2H), 4.80 (m, 1H), 5.64 (s, 2H), 6.58 (d, 1H), 6.92 (d, 1H), 7.10 (d, 1H), 7.30 (m, 2H), 7.40 - 7.50 (m, 3H), 7.60 (dt, 1H), 7.68 (t, 1H), 7.90 (d, 1H), 8.13 (d, 1H), 8.42 (s, 1H), 10.11 (bs, 1H); Mass Spectrum MH⁺ 489.

Example 10.3**5-(1-Methylpiperidin-4-yloxy)-4-(1-(2-pyridylmethyl)indol-5-ylamino)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2-picoyl chloride hydrochloride in 42% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.62 (m, 2H), 4.80 (m, 1H), 5.50 (s, 2H), 6.52 (d, 1H), 6.98 (d, 1H), 7.10 (d, 1H), 7.25 (m, 3H), 7.42 (d, 1H), 7.52 (d, 1H), 7.69 (m, 2H), 7.90 (d, 1H), 8.08 (d, 1H), 8.41 (s, 1H), 8.53 (d, 1H), 10.10 (bs, 1H); Mass Spectrum MH⁺ 465.

Example 10.4**5-(1-Methylpiperidin-4-yloxy)-4-(1-(thiazol-4-ylmethyl)indol-5-ylamino)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 4-chloromethylthiazole in 75% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.52 (s, 2H), 6.48 (d, 1H), 7.10 (d, 1H), 7.30 (m, 2H), 7.48 (m, 2H), 7.58 (d, 1H), 7.68 (t, 1H), 8.07 (d, 1H), 8.42 (s, 1H), 9.02 (d, 1H); Mass Spectrum MH⁺ 471.

Example 10.5**4-(1-(4-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 4-fluorobenzyl chloride in 78% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.40 (s, 2H), 6.50 (d, 1H), 7.10 - 7.30 (m, 7H), 7.48 (d, 1H), 7.53 (d, 1H), 7.69 (t, 1H), 8.08 (d, 1H), 8.42 (s, 1H), 10.10 (bs, 1H); Mass Spectrum MH⁺ 482.

Example 10.6**4-(1-(2-Methoxybenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2-methoxybenzyl chloride in 20% yield; NMR spectrum

5 (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.60 (m, 2H), 3.84 (3H, s), 4.80 (m, 1H), 5.37 (s, 2H), 6.48 (d, 1H), 6.80 (m, 2H), 7.02 (d, 1H), 7.18 (d, 1H), 7.22 (m, 2H), 7.28 (d, 1H), 7.42 (m, 2H), 7.67 (t, 1H), 8.07 (d, 1H), 8.42 (s, 1H), 10.10 (bs, 1H); Mass Spectrum MH⁺ 494.

Example 10.7**10 4-(1-(2-Chlorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2-chlorobenzyl chloride in 61% yield; NMR spectrum (DMSO-

d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.52 (s, 2H), 6.55 (d, 1H), 6.70 (d, 1H), 7.02 (d, 1H), 7.20 (m, 2H), 7.30 (m, 3H), 7.40 (d, 1H), 7.45
15 (m, 2H), 7.68 (t, 1H), 8.12 (d, 1H), 8.42 (s, 1H), 10.11 (bs, 1H); Mass Spectrum M⁺ 498.

Example 10.8**4-(1-(2,5-Dimethylbenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2,5-dimethylbenzyl chloride in 75% yield; NMR spectrum

20 (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.10 (s, 3H), 2.15 (s + m, 5H), 2.23 (s, 3H), 2.25 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.37 (s, 2H), 6.51 (m, 2H), 6.97 (d, 1H), 7.05 (d, 1H), 7.20 (d, 1H), 7.22 - 7.37 (m, 4H), 7.40 (d, 1H), 7.68 (t, 1H), 8.12 (d, 1H), 8.42 (s, 1H), 10.12 (bs, 1H); Mass Spectrum MH⁺ 492.

Example 10.9**25 4-(1-(3-Chlorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 3-chlorobenzyl chloride in 70% yield; NMR spectrum (DMSO-

d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.60 (m, 2H), 4.80 (m, 1H), 5.43 (s, 2H), 6.52 (d, 1H), 7.10 - 7.38 (m, 7H), 7.48 (d, 1H), 7.57 (d, 1H), 7.70 (t, 1H), 8.10 (d, 1H),
30 8.42 (s, 1H), 10.10 (bs, 1H); Mass Spectrum M⁺ 498.

Example 10.10**5-(1-Methylpiperidin-4-yloxy)-4-(1-(2-methylthiazol-4-ylmethyl)indol-5-ylamino)quinazoline**

- Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 4-chloromethyl-2-methylthiazole hydrochloride in 37% yield; NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.15 (s + m, 5H), 2.25 (m, 2H), 2.58 (s, 3H), 2.60 (m, 2H), 4.80 (m, 1H), 5.41 (s, 2H), 6.47 (d, 1H), 7.20 (d, 1H), 7.30 (m, 3H), 7.45 (d, 1H), 7.56 (d, 1H), 7.68 (t, 1H), 8.05 (d, 1H), 8.42 (s, 1H), 10.10 (bs, 1H); Mass Spectrum MH⁺ 485.

Example 10.11

- 10 **4-(1-(2-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 2-fluorobenzyl chloride in 18% yield; Mass Spectrum MH⁺ 482.

Example 10.12**4-(1-(3-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

- 15 Obtained from 4-(indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.4) and 3-fluorobenzyl chloride in 33% yield; Mass Spectrum MH⁺ 482.

Example 11**4-(4-Benzyloxy-3-ethynylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

- Sodium dithionite (1 g) was added to a solution of 4-benzyloxy-3-ethynyl-nitrobenzene (reference example 28) in ethanol (10 ml) and water (10 ml) and the solution heated at 80°C for 3 hours. The ethanol was removed *in vacuo* and the residue extracted with DCM. Combined organic extracts were dried and concentrated and the residue dissolved in IPA (0.5 ml). To this was added di-*iso*-propylethylamine (0.025 ml) and 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and the solution heated at 90°C for 3 hours. The reaction was cooled and the resulting precipitate collected, washed with IPA and ether, and dried to give the title compound as a cream-coloured solid (3.5 mg, 2%); Mass Spectrum MH⁺ 465.

Example 12**4-(3-Ethynyl 4-(2-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

- 30 **hydrochloride**

A solution of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) (70 mg) and 3-ethynyl-4-(2-fluorobenzyloxy)aniline (reference example 30) (70 mg) in

IPA (2 ml) was heated at reflux for 2 hours. The solution was cooled and the resulting precipitate filtered, washed with IPA and ether to give the title compound as a yellow solid (0.115 g, 89%); Mass Spectrum MH^+ 483.

The procedure described above was repeated using the appropriate chloroquinazoline and aniline. Thus were obtained the compounds described below:

Example 12.1

4-(3-Ethynyl 4-(3-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-ethynyl-4-(3-fluorobenzyloxy)aniline (reference example 30.1) in 80% yield; Mass Spectrum MH^+ 483.

Example 12.2

4-(3-Ethynyl 4-(2,6-difluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 4-(2,6-difluorobenzyloxy)-3-ethynylaniline (reference example 30.2) in 73% yield; Mass Spectrum MH^+ 502.

Example 12.3

4-(3-Ethynyl 4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-ethynyl-4-(5-methylisoxazol-3-ylmethoxy)aniline (reference example 30.4) in 61% yield; Mass Spectrum MH^+ 471.

Example 12.4

4-(3-Ethynyl 4-(thiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-ethynyl-4-(thiazol-4-ylmethoxy)aniline (reference example 30.3) in 20% yield; Mass Spectrum MH^+ 473.

Example 12.5**4-(3-Chloro-4-(2-pyrimidinylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) with 3-chloro-4-(2-pyrimidinylmethoxy)aniline (reference example 9.2) in 40% yield; NMR spectrum (DMSO-d₆) 1.90 (m, 2H), 2.15 (m, 2H), 2.20 (s, 3H), 2.25 (m, 2H), 2.60 (m, 2H), 4.75 (m, 1H), 5.40 (s, 2H), 7.15 (d, 1H), 7.20 (d, 2H), 7.30 (d, 1H), 7.45 (dd, 1H), 7.47 (t, 1H), 7.70 (t, 1H), 8.10 (d, 1H), 8.50 (s, 1H), 8.85 (d, 2H), 10.0 (s, 1H); Mass spectrum M-H⁺ 477.

10 Example 12.6**4-(4-(2-Aminothiazol-4-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 4-(2-aminothiazol-4-ylmethoxy)-3-chloroaniline (reference example 9.3) in 75% yield; NMR spectrum (DMSO-d₆) 1.84 - 1.95 (m, 2H), 2.07 - 2.17 (m, 2H), 2.17 (s, 3H), 2.22 - 2.32 (m, 2H), 2.57 - 2.67 (m, 2H), 4.78 (m, 1H), 4.94 (s, 2H), 6.59 (s, 1H), 6.95 (s, 2H), 7.21 (d, 1H), 7.29 (d, 1H), 7.32 (d, 1H), 7.49 (dd, 1H), 7.70 (dd, 1H), 8.08 (d, 1H), 8.49 (s, 1H), 10.03 (s, 1H); Mass spectrum M-H⁺ 497.

Example 12.7**20 4-(3-Fluoro-4-(1-methyl-1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-fluoro-4-(1-methyl-1H-imidazol-2-ylthio)aniline (reference example 6.2) in 31% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.1 - 2.2 (m, 2H), 2.2 (s, 3H), 2.25 - 2.35 (m, 2H), 2.6 - 2.7 (m, 2H), 3.7 (s, 3H), 4.7 - 4.8 (m, 1H), 7.0 (s, 1H), 7.1 - 7.15 (t, 1H), 7.2 (d, 1H), 7.3 (s, 1H), 7.35 - 7.42 (dd, 1H), 7.7 - 7.8 (t, 1H), 8.0 - 8.1 (dd, 1H), 8.55 (s, 1H), 10.2 (bs, 1H); Mass Spectrum MH⁺ 465.

Example 12.8**30 4-(3-Fluoro-4-(1-methyl-1H-1,3,4-triazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-fluoro-4-(1-methyl-1H-1,3,4-triazol-2-ylthio)aniline (reference example 6.3) in 16% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.1 - 2.2 (m, 2H), 2.2 (s, 3H), 2.25

- 2.35 (m, 2H), 2.6 - 2.7 (m, 2H), 3.6 (s, 3H), 4.7 - 4.8 (m, 1H), 7.2 (d, 1H), 7.35 - 7.5 (m, 3H), 7.7 - 7.8 (t, 1H), 8.1 - 8.2 (d, 1H), 8.5 - 8.6 (d, 2H), 10.2 (bs, 1H); Mass Spectrum MH^+ 466.

Example 12.9

4-(3-Chloro-4-(2-pyridylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

5 hydrochloride

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-Chloro-4-(2-pyridylthio)aniline (reference example 6.4) in 11% yield; NMR spectrum (DMSO- d_6 , 373K) 1.9 - 2.1 (m, 2H), 2.15 - 2.3 (m, 2H), 2.3 (s, 3H), 2.3 - 2.45 (m, 2H), 2.7 - 2.85 (m, 2H), 4.75 - 4.9 (m, 1H), 7.0 - 7.1 (d, 1H), 7.15 - 7.25 (dd, 1H), 7.25 - 7.35 (d, 1H), 7.4 - 7.5 (d, 1H), 7.6 - 7.7 (m, 1H), 7.7 - 7.85 (m, 3H), 8.3 - 8.45 (m, 2H), 8.6 (s, 1H), 10.3 (bs, 1H); Mass Spectrum MH^+ 478.

Example 12.10

4-(3-Chloro-4-(2-pyrimidinylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

hydrochloride

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-pyrimidinylthio)aniline (reference example 6.5) in 6% yield; NMR spectrum (DMSO- d_6 , 373K) 1.9 - 2.0 (m, 2H), 2.2 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.4 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 - 4.85 (m, 1H), 7.2 - 7.25 (t, 1H), 7.25 - 7.3 (d, 1H), 7.4 - 7.45 (d, 1H), 7.7 - 7.75 (m, 1H), 7.75 - 7.85 (d, 2H), 8.45 (s, 1H), 8.6 (d, 2H), 8.65 (s, 1H), 10.3 (bs, 1H); Mass Spectrum MH^+ 479.

Example 12.11

4-(3-Chloro-4-(1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

hydrochloride

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(1H-imidazol-2-ylthio)aniline (reference example 9.6) in 74% yield; NMR spectrum (DMSO- d_6 , 373K) 1.8 - 1.9 (m, 2H), 2.1 - 2.25 (m, 2H), 2.15 (s, 3H), 2.6 - 2.7 (m, 2H) 4.7 - 4.8. (m, 1H), 6.7 - 6.8 (d, 1H), 7.1 (bs, 1H), 7.2 - 7.25 (d, 1H), 7.3 - 7.4 (d, 1H), 7.4 - 7.5 (dd, 1H), 7.7 - 7.8 (t, 1H), 8.35 (d, 1H), 8.55 (s, 1H), 10 - 10.2 (bs, 1H), 12.8 - 12.9 (bs, 1H); Mass spectrum MH^+ 467.

Example 12.12**4-(3-Fluoro-4-(1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-fluoro-4-(1*H*-imidazol-2-ylthio)aniline (reference example 8.5) in 33% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.6 - 2.7 (m, 2H) 4.7 - 4.8. (m, 1H), 7.1 (s, 1H), 7.2 - 7.3 (m, 1H), 7.4 - 7.5 (d, 1H), 7.7 - 7.8 (t, 1H), 8.1 - 8.2 (d, 1H), 8.6 (s, 1H), 10.2-10.5 (bs, 1H), 12.0 - 12.8 (bs, 1H); Mass spectrum MH⁺ 451.

Example 12.13**4-(3-Chloro-4-(2-thiazolylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-(2-thiazolylthio)aniline (reference example 9.7) in 29% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.4 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 - 4.85 (m, 1H), 7.2 - 7.5 (d, 1H), 7.4 - 7.45 (d, 1H), 7.7 (d, 1H), 7.8 (s, 1H), 7.8 - 7.9 (m, 2H), 8.4 (s, 1H), 8.65 (s, 1H), 10.4 (bs, 1H); Mass spectrum MH⁺ 482.

Example 13**4-(3-Chloro-4-(2-pyrazinylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

A suspension of 2-methylpyrazine (940 mg); *N*-chlorosuccinimide (1.34 g) and benzoyl peroxide (70%, 71 mg) in carbon tetrachloride (50 ml) was heated at reflux for 24 hours. The reaction was cooled to 0°C in an ice bath and then filtered through diatomaceous earth. To the crude solution of 2-chloromethylpyrazine in carbon tetrachloride was added potassium carbonate (138 mg), 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) (90 mg) and *cis*-dicyclohexano-18-crown-6 (5 mg), and the reaction mixture heated at 80°C. After 16 hours, the reaction mixture was concentrated *in vacuo* and the resulting residue was partitioned between DCM and water. The combined organic extracts were dried and concentrated and the residue purified by chromatography (0-3% 7N methanolic ammonia in DCM) to yield the title compound as a pale pink solid (46 mg; 41%); NMR spectrum (DMSO-d₆) 1.90 (m, 2H), 2.15 (m, 2H), 2.20 (s, 3H), 2.25 (m, 2H), 2.60 (m, 2H), 4.75 (m, 1H), 5.35 (s, 2H), 7.20 (d, 1H), 7.35 (d, 2H), 7.50 (dd, 1H), 7.70 (t, 1H), 8.15 (d, 1H), 8.50 (s, 1H), 8.65 (dd, 2H), 8.85 (s, 1H); Mass spectrum M-H⁺ 477.

The procedure described above was repeated using the appropriate methylheterocycle. Thus was obtained the compound described below:

Example 13.1

4-(3-Chloro-4-(4-pyrimidinylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

5 Obtained from 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) and 4-methylpyrimidine in 12% yield; NMR spectrum (DMSO-d₆) 1.95 (m, 2H), 2.15 (m, 2H), 2.25 (s, 3H), 2.40 (m, 2H), 2.75 (m, 2H), 4.80 (m, 1H), 5.35 (s, 2H), 7.25 (t, 1H), 7.35 (d, 2H), 7.50 (dd, 1H), 7.65 (dd, 1H), 7.70 (t, 1H), 8.15 (d, 1H), 8.50 (s, 1H), 8.85 (d, 2H), 9.20 (d, 1H), 10.0 (s, 1H); Mass spectrum M-H⁺ 477.

10 **Example 14**

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Potassium carbonate (43.2 g) was added to a solution of 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (30 g) (reference example 4.2), 2-picoly chloride hydrochloride (13.8 g) and 18-crown-6 (1 g) in acetonitrile (1000 ml) and the reaction heated at
15 reflux for 3 hours. The reaction was filtered whilst hot, and allowed to cool. The resulting crystalline solid was filtered and recrystallised from acetonitrile to give the title compound as beige needles (19.36 g, 52%); NMR spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.10 - 2.20 (s + m, 5H), 2.22 (m, 2H), 2.60 (m, 2H), 4.75 (m, 1H), 5.24 (s, 2H), 7.18 (d, 1H), 7.23 (d, 1H), 7.33 (m, 2H), 7.45 (dd, 1H), 7.54 (d, 1H), 7.66 (t, 1H), 7.83 (dt, 1H), 8.12 (d, 1H), 8.47 (s, 1H), 8.57
20 (d, 1H), 10.02 (bs, 1H); Mass spectrum M⁺ 476.

The procedure described above was repeated using the appropriate halomethyl compound. Thus were obtained the compounds described below:

Example 14.1

4-(3-Chloro-4-(imidazo[1, 2-a]pyridin-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

25 Obtained from 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) and 2-chloromethylimidazo[1, 2-a]pyridine in 28% yield; NMR spectrum (DMSO-d₆) 1.85 - 1.97 (m, 2H), 2.10 - 2.20 (m, 2H), 2.17 (s, 3H), 2.23 - 2.34 (m, 2H), 2.56 - 2.69 (m, 2H), 4.80 (m, 1H), 5.32 (s, 2H), 6.89 (ddd, 1H), 7.23 (d, 1H),
30 7.24 (ddd, 1H), 7.33 (d, 1H), 7.41 (d, 1H), 7.51 (dd, 1H), 7.54 (dd, 1H), 7.72 (dd, 1H), 8.02 (s, 1H), 8.11 (d, 1H), 8.50 (s, 1H), 8.55 (dd, 1H), 10.06 (s, 1H); Mass spectrum MH⁺ 515.

Example 14.2**4-(4-(Benzo[d]isoxazol-3-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 3-bromomethylbenzo[d]isoxazole (prepared as in Chim. Ther. 1972, 5 7(2), 127-132) and 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) in 36% yield; NMR spectrum (DMSO-d₆) 1.86 - 1.97 (m, 2H), 2.08 - 2.22 (m, 2H), 2.18 (s, 3H), 2.23 - 2.33 (m, 2H), 2.57 - 2.68 (m, 2H), 4.79 (m, 1H), 5.73 (s, 2H), 7.23 (d, 1H), 7.34 (d, 1H), 7.42 (d, 1H), 7.47 (dd, 1H), 7.55 (dd, 1H), 7.71 (ddd, 1H), 7.73 (dd, 1H), 7.80 (d, 1H), 8.03 (d, 1H), 8.15 (d, 1H), 8.52 (s, 1H), 10.07 (s, 1H); Mass spectrum MH⁺ 10 516.

Example 15**4-(3-Chloro-4-(2-pyrimidinylloxy)-5-(tetrahydrofuran-3-yloxy)quinazoline**

A mixture of 4-(3-chloro-4-hydroxyanilino)-5-(tetrahydrofuran-3-yloxy)quinazoline (70 mg) (reference example 42.1), 2-chloropyrimidine (25 mg), potassium carbonate (275 mg) 15 and 1,4,7,10,13,16-hexaoxacyclooctadecane (5 mg) in acetonitrile (8 ml) was heated at reflux for 30 hours. Water (30 ml) was added and the mixture was extracted with DCM (30 ml). The extracts were dried and concentrated *in vacuo*. The residue was purified by chromatography using 0-2% methanol-DCM as eluent to afford the title compound as a solid (15 mg, 17%); NMR Spectrum (DMSO-d₆) 2.15 - 2.24 (m, 1H), 2.30 - 2.42 (m, 1H), 3.8 - 3.94 (m, 2H), 3.98 20 (q, 1H), 4.23 (d, 1H), 5.46 (t, 1H), 7.23 (d, 1H), 7.28 (t, 1H), 7.39 (d, 1H), 7.41 (d, 1H), 7.70 (dd, 1H), 7.77 (t, 1H), 8.31 (d, 1H), 8.60 (s, 1H), 8.65 (d, 2H), 10.21 (s, 1H); Mass spectrum MH⁺ 436.

Example 16**4-(3-Chloro-4-(1,2,4-oxadiazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline** 25

2-(2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy)-N-hydroxyacetamide (200 mg) (reference example 35) was dissolved in formic acid (4 ml). Trimethyl orthoformate (4 ml) was added, and the mixture stirred at ambient temperature for 1 hour. The mixture was evaporated, and the residue purified by chromatography, using 0 to 30 1.5% (7:1 methanol / conc. ammonia (aq)) in DCM as eluent to give the title compound as a white crystalline solid (157 mg, 77%); NMR spectrum (DMSO-d₆) 1.83 - 1.96 (m, 2H), 2.07 - 2.20 (m, 2H), 2.17 (s, 3H), 2.22 - 2.32 (m, 2H), 2.57 - 2.67 (m, 2H), 4.78 (m, 1H), 5.45 (s, 2H),

7.22 (d, 1H), 7.32 (d, 1H), 7.33 (d, 1H), 7.52 (dd, 1H), 7.71 (dd, 1H), 8.12 (d, 1H), 8.51 (s, 1H), 9.67 (s, 1H), 10.05 (s, 1H); Mass spectrum MH^+ 467.

The procedure described above was repeated using the appropriate hydroxyacetamidine and orthoester. Thus was obtained the compound described below:

5 **Example 16.1**

4-(3-Chloro-4-(5-methyl-1,2,4-oxadiazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained from 2-(2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy)-*N*-hydroxyacetamidine (reference example 35) and trimethyl orthoacetate
10 in 60% yield; NMR spectrum (DMSO- d_6) 1.83 - 1.97 (m, 2H), 2.08 - 2.18 (m, 2H), 2.17 (s, 3H), 2.21 - 2.31 (m, 2H), 2.57 - 2.67 (m, 2H), 2.61 (s, 3H), 4.78 (m, 1H), 5.34 (s, 2H); 7.21 (d, 1H), 7.31 (d, 1H), 7.32 (d, 1H), 7.52 (dd, 1H), 7.71 (dd, 1H), 8.11 (d, 1H), 8.50 (s, 1H), 10.05 (s, 1H); Mass spectrum MH^+ 467.

Example 17

15 **4-(4-(5-Amino-1,3,4-oxadiazol-2-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Potassium hydrogen carbonate (10 mg) was dissolved in a mixture of ethanol (2 ml) and water (4 ml) 2-[2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetic acid hydrazide (reference example 37) (40 mg) was added. The
20 suspension was briefly sonicated, then cooled to 0°C. Cyanogen bromide (3M in DCM, 31 μ l) was added dropwise. The mixture was allowed to warm to ambient temperature overnight with stirring. The resulting yellow solution was evaporated, and the residue purified by chromatography, using 0 to 4% (7:1 methanol / conc. ammonia (aq)) in DCM as eluent to give the title compound as a white crystalline solid (14 mg, 33%); NMR spectrum (DMSO- d_6) 1.87
25 - 1.97 (m, 2H), 2.08 - 2.20 (m, 2H), 2.17 (s, 3H), 2.20 - 2.34 (m, 2H), 2.58 - 2.69 (m, 2H), 4.79 (m, 1H), 5.26 (s, 2H), 7.20 (s, 2H), 7.24 (d, 1H), 7.34 (d, 1H), 7.36 (d, 1H), 7.56 (d, 1H), 7.72 (dd, 1H), 8.13 (d, 1H), 8.52 (s, 1H), 10.09 (s, 1H); Mass spectrum MH^+ 482.

Example 18

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline

30 Phosphorus oxychloride (0.21 ml) was added dropwise to a solution of 3,4-dihydro-5-(tetrahydropyran-4-yloxy)quinazolin-4-one (reference example 1.2) (57 mg) and di-*iso*-propylethylamine (0.045 ml) in anhydrous 1, 2-dichloroethane (5 ml) at 0°C. The mixture was

heated at 80°C for 3 hours and then concentrated *in vacuo*. The residue was azeotroped with toluene (5 ml), di-*iso*-propylethylamine (0.5 ml) was added and the mixture was again concentrated *in vacuo*. The residue was dissolved in IPA (1 ml) and 3-chloro-4-(2-pyridylmethoxy)aniline (obtained as described in PCT Int. Appl. WO 9615118) (100 mg) was added. The resulting mixture was heated at 80°C for 12 hours and then concentrated *in vacuo*. The residue was purified by chromatography, using 0-5% methanol - DCM as eluent, to give the title compound as a solid (48 mg, 43%); NMR Spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.13 - 2.27 (m, 2H), 3.54 (t, 2H), 3.84 - 3.98 (m, 2H), 4.89 - 5.00 (m, 1H), 5.28 (s, 2H), 7.21 - 7.39 (m, 4H), 7.49 (d, 1H), 7.56 (d, 1H), 7.71 (t, 1H), 7.85 (t, 1H), 8.15 - 8.20 (m, 1H), 8.50 (s, 1H), 8.54 - 8.61 (m, 1H), 10.03 (s, 1H); Mass spectrum MH⁺ 463.

The procedure described above was repeated using the appropriate 3,4-dihydroquinazolin-4-one and aniline. Thus were obtained the compounds described below:

Example 18.1

4-(1-(3-Fluorobenzyl)indazol-5-ylamino)-5-(tetrahydropyran-4-yloxy)quinazoline

Obtained by reacting 3,4-dihydro-5-(tetrahydropyran-4-yloxy)quinazolin-4-one (reference example 1.2) and 5-amino-1-(3-fluorobenzyl)indazole (obtained as described in PCT Int. Appl. WO 9802438) in 19% yield; NMR Spectrum (DMSO-d₆) 1.80 - 1.95 (m, 2H), 2.16 - 2.27 (m, 2H), 3.55 (dt, 2H), 3.91 (dt, 2H), 4.93 - 5.03 (m, 1H), 5.69 (s, 2H), 7.02 - 7.12 (m, 3H), 7.26 (d, 1H), 7.30 - 7.38 (m, 2H), 7.54 (dd, 1H), 7.71 (t, 2H), 8.14 (s, 1H), 8.39 (d, 1H), 8.48 (s, 1H), 10.16 (s, 1H); Mass spectrum MH⁺ 470.

Example 18.2

4-(3-Chloro-4-((1H-imidazol-2-ylthio)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline

Obtained by reacting 3,4-dihydro-5-(tetrahydropyran-4-yloxy)quinazolin-4-one (reference example 1.2) and 3-chloro-4-(1H-imidazol-2-ylthio)aniline (reference example 9.6) in 10% yield; NMR Spectrum (DMSO-d₆) 1.78 - 1.93 (m, 2H), 2.12 - 2.24 (m, 2H), 3.52 (dt, 2H), 3.89 (dt, 2H), 4.88 - 4.99 (m, 1H), 6.75 (d, 1H), 7.26 (bs, 1H), 7.28 (d, 1H), 7.35 - 7.44 (m, 3H), 7.73 (t, 1H), 8.40 (d, 1H), 8.57 (s, 1H), 10.18 (s, 1H), 12.86 (s, 1H); Mass spectrum M-H⁺ 452.

Example 18.3

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(tetrahydrofuran-3-yloxy)quinazoline

Obtained by reacting 3,4-dihydro-5-(tetrahydrofuran-3-yloxy)quinazolin-4-one (reference example 1.3) and 3-chloro-4-(2-pyridylmethoxy)aniline (obtained as described in

PCT Int. Appl. WO 9615118) in 15% yield; NMR Spectrum (CDCl₃) 2.23 - 2.45 (m, 2H), 3.94 - 4.04 (m, 2H), 4.10 (q, 1H), 4.26 (d, 1H), 5.25 (t, 1H), 5.28 (s, 2H), 6.84 (d, 1H), 6.99 (d, 1H), 7.22 (dd, 1H), 7.44 - 7.54 (m, 2H), 7.59 - 7.66 (m, 2H), 7.74 (dt, 1H), 8.11 (d, 1H), 8.60 (d, 1H), 8.64 (s, 1H), 9.90 (s, 1H); Mass spectrum MH⁺ 449.

5 Example 19

4-(3-Chloro-4-(1-cyanomethyl-1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

To a stirred solution of 4-(3-chloro-4-(1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) (460 mg) in acetonitrile (30 ml) was
10 added potassium carbonate (1.0 g), chloroacetonitrile (80 mg) and *cis*-dicyclohexano-18-crown-6 (20 mg) and the solution stirred and heated at reflux for 18 hours. The solution was cooled, filtered and evaporated, and the residue purified by chromatography using ethyl acetate and then ammonia/methanol/DCM (30ml of 2.3N ammonia in methanol, 970ml of DCM) as eluent. Evaporation of the relevant fractions and trituration with ether yielded the title
15 compound as a white solid (160 mg, 31%); NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.4 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 - 4.85 (m, 1H), 5.4 (s, 2H), 7.0 - 7.05 (d, 1H), 7.2 - 7.25 (t, 1H), 7.4 - 7.45 (d, 1H), 7.5 - 7.6 (dd, 1H), 7.6 (s, 1H), 7.7 - 7.8 (t, 1H), 8.3 (d, 1H), 8.6 ((s, 1H), 10.1 (bs, 1H); Mass spectrum MH⁺ 504.

The procedure described above was repeated using the appropriate imidazole and alkyl
20 halide. Thus were obtained the compounds described below:

Example 19.1

4-(4-(1-Carbamoylmethyl-1H-imidazol-2-ylthio)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) with 2-chloroacetamide in 34% yield;
25 NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.4 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 (s, 2H), 4.75 - 4.85 (m, 1H), 7.0 - 7.05 (d, 1H), 7.0 - 7.2 (bs, 2H), 7.2 (s, 1H), 7.2 - 7.25 (d, 1H), 7.4 - 7.45 (d, 1H), 7.5 - 7.55 (dd, 1H), 7.7 - 7.8 (t, 1H), 8.25 (d, 1H), 8.6 (s, 1H), 10.1 (bs, 1H); Mass spectrum MH⁺ 522.

30 Example 19.2

4-(3-Chloro-4-(1-(2-methoxyethyl)-1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) with 2-bromoethylmethylether in 61% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.45 (m, 2H), 2.7 - 2.8 (m, 2H), 3.2 (s, 3H), 3.6 - 3.65 (t, 2H), 4.2 - 4.25 (t, 2H), 4.75 - 4.85 (m, 1H), 6.9 - 6.95 (d, 1H), 7.1 (s, 1H), 7.2 - 7.25 (d, 1H), 7.4 - 7.45 (d, 1H), 7.45 (s, 1H), 7.5 - 7.55 (dd, 1H), 7.7 - 7.8 (t, 1H), 8.2 (s, 1H), 8.55 (s, 1H), 10.1 (bs, 1H); Mass spectrum MH⁺ 525.

Example 19.3

4-(3-Chloro-4-(1-(*N,N*-diethylcarbamoylmethyl)-1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) with 2-chloro-*N,N*-diethylacetamide in 28% yield; NMR spectrum (DMSO-d₆, 373K) 1.0 - 1.25 (bs, 6H), 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.45 (m, 2H), 2.7 - 2.8 (m, 2H), 3.3 - 3.4 (q, 4H), 4.75 - 4.85 (m, 1H), 5.0 (s, 1H), 6.95 - 7.0 (d, 1H), 7.15 (s, 1H), 7.2 - 7.25 (d, 1H), 7.3 - 7.35 (d, 1H), 7.35 (s, 1H), 7.45 - 7.5 (dd, 1H), 7.7 - 7.8 (t, 1H), 8.2 (s, 1H), 8.55 (s, 1H), 10.1 (bs, 1H); Mass spectrum MH⁺ 578.

Example 19.4

4-(4-(1-*tert*-Butoxycarbonylmethyl)-1*H*-imidazol-2-ylthio)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) with *tert*-butyl 2-bromoacetate in 44% yield; Mass spectrum MH⁺ 581.

Example 19.5

4-(3-Chloro-4-(1-difluoromethyl-1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-chloro-4-(1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (**example 12.11**) with ethyl difluorobromoacetate in 34% yield; NMR spectrum (DMSO-d₆, 373K) 1.95 - 2.05 (m, 2H), 2.15 - 2.25 (m, 2H), 2.3 (s, 3H), 2.3 - 2.45 (m, 2H), 2.7 - 2.85 (m, 2H), 4.75 - 4.85 (m, 1H), 7.15 - 7.2 (d, 1H), 7.2 - 7.3 (d, 1H),

7.3 (s, 1H), 7.4 - 7.45 (d, 1H), 7.6 - 7.65 (dd, 1H), 7.7 - 8.0 (q, 1H), 7.7 - 7.8 (t, 1H), 8.3 (s, 1H), 8.65 (s, 1H), 10.0-10.2 (bs, 1H); Mass spectrum MH^+ 517.

Example 19.6

4-(4-(1-Cyanomethyl-1H-imidazol-2-ylthio)-3-fluoroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

Obtained by reacting 4-(3-fluoro-4-(1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (example 12.12) with chloroacetonitrile in 40% yield; NMR spectrum (DMSO-d₆, 373K) 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.2 (s, 3H); 2.25 - 2.35 (m, 2H), 2.6 - 2.7 (m, 2H), 4.7 - 4.8 (m, 1H), 5.35 (s, 2H), 7.1 (s, 1H), 7.2 - 7.25 (d, 1H), 7.25 - 7.3 (t, 1H), 7.4 - 7.42 (d, 1H), 7.42 - 7.45 (dd, 1H), 7.5 (s, 1H), 7.7 - 7.8 (t, 1H), 8.0 - 8.1 (dd, 1H), 8.6 (s, 1H), 10.2 (bs, 1H); Mass spectrum MH^+ 490.

Example 20

4-(4-(1-Carboxymethyl-1H-imidazol-2-ylthio)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline dihydrochloride.

A solution of 4-(3-chloro-4-(1-tert-butoxycarbonylmethyl-1H-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (example 19.4) (100 mg) in dioxane (20 ml) was treated with an ethereal solution of hydrogen chloride (3 ml, 1M), and heated at reflux for 4 hours. The solution was cooled to give a precipitate that was filtered and washed with ether to give the title compound as a white solid (98 mg, 95%); NMR spectrum (DMSO-d₆, 373K) 2.3 - 2.5 (m, 2H), 2.5 - 2.6 (m, 2H), 2.8 (s, 3H), 3.1 - 3.4 (bs, 2H), 3.4 - 3.7 (bs, 2H), 4.9 (s, 2H), 5.0 - 5.2 (m, 1H), 7.15 - 7.2 (d, 1H), 7.3 (s, 1H), 7.45 - 7.5 (d, 1H), 7.6 (s, 1H), 7.6 - 7.7 (m, 1H), 7.9 - 8.0 (t, 1H), 8.2 - 8.3 (bs, 1H), 8.8 (s, 1H), 10.2-10.5 (bs, 1H), 10.8 - 11.5 (bs, 1H); Mass spectrum MH^+ 523.

Example 21

5-(1-Methylpiperidin-4-yloxy)-4-(4-(thiazol-2-ylthio)anilino)quinazoline

To a solution of 4-(4-iodoanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.6) (229 mg) in anhydrous and degassed toluene (3 ml) in a pressure vial was added 2-mercaptothiazole (114 mg), cuprous bromide (15 mg) and 1,8-diazabicyclo[5,4,0]undec-7-ene (152 mg). The solution was purged with nitrogen and the vial capped and then heated at 118°C for 18 hours. The contents of the vessel were purified by chromatography using ammonia/methanol/DCM (30 ml of 2.3N ammonia in methanol, 970 ml of DCM) as eluent to give the title compound as a solid (195 mg, 87%); NMR spectrum (DMSO-d₆, 373K) 1.5 - 1.7

(m, 2H), 2.1 - 2.2 (m, 1H), 2.2 (s, 3H), 2.2 - 2.4 (m, 1H), 2.6 - 2.65 (m, 1H), 2.7 - 2.8 (m, 1H), 3.4 - 3.5 (t, 1H), 3.5 - 3.6 (d, 1H), 4.7 - 4.9 (1H), 7.25 - 7.3 (d, 1H), 7.35 - 7.4 (d, 1H), 7.6 (s, 1H), 7.65 - 7.8 (m, 4H), 7.95 - 8.0 (d, 2H), 8.6 (s, 1H), 10.3 - 10.35 (bs, 1H); Mass spectrum MH^+ 450.

5 Example 22

5-(1-Methylpiperidin-4-yloxy)-4-(4-(2-thiazolylsulphonyl)anilino)quinazoline

To a solution of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) (80 mg) and 4-thiazol-2-ylsulphonylaniline (reference example 8.6) (140 mg) in anhydrous THF was added sodium hydride (100 mg, 40% dispersion in oil). The solution was stirred and heated at reflux overnight. The reaction was concentrated and the residue purified by chromatograph using ethyl acetate and then a mixture of 3% 2.3N ammonia in methanol in DCM as eluent to give the title compound as a white solid (40 mg, 28%); NMR spectrum (DMSO- d_6 , 373K). 1.9 - 2.0 (m, 2H), 2.15 - 2.25 (m, 2H), 2.25 (s, 3H), 2.3 - 2.4 (m, 2H), 2.7 - 2.8 (m, 2H), 4.75 - 4.85 (m, 1H), 7.2 - 7.3 (d, 1H), 7.4 - 7.5 (d, 1H), 7.7 - 7.8 (t, 1H), 8.0 - 8.1 (m, 3H), 8.1 - 8.2 (m, 3H), 8.65 (s, 1H), 10.4 (bs, 1H); Mass spectrum MH^+ 482.

Example 23

4-(3-Chloro-4-(3-fluorobenzoyloxy)aniline)-5-(1-(4-methylpiperazin-1-yl)cyclohex-4-yloxy)quinazoline

Sodium triacetoxyborohydride (0.43 g) was added to a solution of 4-(3-chloro-4-(3-fluorobenzoyloxy)aniline)-5-(1-oxo-cyclohex-4-yloxy)quinazoline acetate (reference example 47) (0.1 g) and 1-methylpiperazine (0.34 ml) in DCE (25 ml) and the reaction stirred for 16 hours. The solution was concentrated *in vacuo* and the residue partitioned between DCM and saturated aqueous sodium hydrogen carbonate solution. Combined organic extracts were dried and concentrated and the residue purified by chromatography using DCM - 5% ammonia (7N) in methanol as eluent to give the title compound as a white solid (24 mg, 23%); Mass Spectrum M^+ 576.

Example 24

4-(3-Chloro-4-(1,2,3-thiadiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

4-Hydroxymethyl-1,2,3-thiadiazole (obtained as described in Example 1 of European Patent Appl. EP 0326640) (81 mg) was dissolved in DCM (5 ml). Di-*iso*-propylethylamine (122 μ l) was added, and the solution cooled to 0°C. Methanesulphonyl chloride (54 μ l) was

added dropwise; the solution was allowed to warm to ambient temperature, and was stirred for a further 2 hours. The solvent was evaporated, and the residue dissolved in DMA (10 ml). Potassium carbonate (552 mg) was added, followed by 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) (150 mg). The mixture was
5 briefly sonicated, then stirred at ambient temperature for 16 hours. The solvent was evaporated, and the residue partitioned between DCM and water. Combined organic extracts were filtered through a silicone-treated filter paper and evaporated. The residue was purified by chromatography, using 0 to 2% (7:1 methanol / c. ammonia (aq)) in DCM as eluent. Evaporation of the appropriate fractions followed by crystallisation of the residue from ethyl
10 acetate / iso-hexane gave the title compound as a white crystalline solid (50 mg, 26%); NMR Spectrum (DMSO-d₆) 1.88 - 1.98 (m, 2H), 2.12 - 2.22 (m, 2H), 2.19 (s, 3H), 2.25 - 2.33 (m, 2H), 2.61 - 2.69 (m, 2H), 4.81 (m, 1H), 5.72 (s, 2H), 7.24 (d, 1H), 7.35 (d, 1H), 7.45 (d, 1H), 7.57 (dd, 1H), 7.74 (dd, 1H), 8.15 (d, 1H), 8.54 (s, 1H), 9.32 (s, 1H), 10.09 (s, 1H); Mass Spectrum MH⁺ 483.4.

15 Example 25

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(2-piperidinoethoxy)quinazoline

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) (76 mg) and 2-piperidinoethanol (30 mg) were dissolved in 1,4-dioxane (5 ml) in a 10ml pressure vessel and sodium hydride (40 mg, 60% dispersion in oil) added. The vessel was
20 sealed and heated at 150°C for 20 minutes, using the "Discover™ microwave synthesis system" (CEM Microwave technology Ltd). The reaction was cooled and pressure released and vessel uncapped. The solution purified by chromatography using 2.3 M ammonia / methanol in DCM (3:97) as eluent to give the title compound as a white solid after trituration with ether (25 mg, 26%); NMR spectrum (DMSO-d₆, 373K) 1.3 - 1.4 (m, 2H), 1.4 - 1.5 (m, 4H), 2.4 - 2.5 (m, 4H), 2.8 - 2.85 (t, 2H), 4.35 - 4.45 (t, 2H), 5.3 (s, 2H), 7.1 (d, 1H), 7.2 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.55 (d, 1H), 7.65 - 7.75 (m, 2H), 7.8 - 7.9 (m, 1H), 7.95 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 10.15 - 10.2 (bs, 1H); Mass Spectrum MH⁺ 491.

The procedure described above was repeated using the appropriate alcohol and the 5-fluoroquinazoline. Thus were obtained the compounds described below:

Example 25.1**4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(1-methylpiperidin-2-ylmethoxy)quinazoline**

Obtained by reacting (1-methylpiperidin-2-yl)methanol with 4-(3-chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 23% yield; NMR spectrum (DMSO-d₆, 373K) 1.3 - 1.5 (m, 2H), 1.5 - 1.6 (m, 1H), 1.6 - 1.7 (m, 1H), 1.7 - 1.8 (m, 2H), 2.2 - 2.3 (m, 1H), 2.35 (s, 3H), 4.2 (d, 1H), 4.5 (d, 1H), 5.3 (s, 2H), 7.1 (d, 1H), 7.3 (d, 1H), 7.3 (m, 1H), 7.35 (d, 1H), 7.5 (d, 1H), 7.7 - 7.8 (t, 1H), 7.85 - 7.95 (m, 2H), 8.0 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 10.3 - 10.4 (bs, 1H); Mass Spectrum MH⁺ 491.

Example 25.2**10 4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(2-azepan-1-ylethoxy)quinazoline**

Obtained by reacting 2-azepan-1-ylethanol with 4-(3-chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 46% yield; NMR spectrum (DMSO-d₆, 373K) 1.4 - 1.5 (m, 6H), 1.5 - 1.6 (m, 2H), 2.7 - 2.8 (m, 2H), 3.05 - 3.1 (t, 2H), 4.35 - 4.4 (t, 2H), 5.3 (s, 2H), 7.1 (d, 1H), 7.2 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.6 (d, 1H), 7.7 (dd, 1H), 7.75 (t, 1H), 7.8 - 7.9 (t, 1H), 7.95 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 10.2 (bs, 1H); Mass Spectrum MH⁺ 505.

Example 25.3**4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(2-morpholinoethoxy)quinazoline**

Obtained by reacting 2-morpholinoethanol with 4-(3-chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 42% yield; NMR spectrum (DMSO-d₆, 373K) 2.8 - 2.9 (m, 6H), 3.3 - 3.4 (m, 4H), 4.4 (t, 2H), 5.3 (s, 2H), 7.1 (d, 1H), 7.25 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.6 (d, 1H), 7.7 (dd, 1H), 7.75 (t, 1H), 7.8 - 7.9 (t, 1H), 7.95 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 10.2 (bs, 1H); Mass Spectrum MH⁺ 493.

Example 25.4**25 4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(2-pyrrolidinoethoxy)quinazoline**

Obtained by reacting 2-pyrrolidinoethanol with 4-(3-chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 44% yield; NMR spectrum (DMSO-d₆, 373K) 1.6 - 1.7 (m, 4H), 2.6 - 2.7 (m, 4H), 3.0 (t, 2H), 4.4 (t, 2H), 5.3 (s, 2H), 7.1 (d, 1H), 7.2 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.6 (d, 1H), 7.65 - 7.75 (m, 2H), 7.8 - 7.9 (t, 1H), 7.95 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 10.3 (bs, 1H); Mass Spectrum MH⁺ 475.

Example 25.5**4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(3-morpholinopropoxy)quinazoline**

Obtained by reacting 3-morpholinopropanol with 4-(3-chloro-4-(2-

- 5 pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 15% yield; NMR spectrum (DMSO-d₆, 373K) 2.1 - 2.2 (m, 2H), 2.3 - 2.4 (m, 4H), 2.5 - 2.6 (m, 2H), 3.5 - 3.6 (m, 4H), 4.4 (t, 2H), 5.3 (s, 2H), 7.1 (d, 1H), 7.25 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.5 - 7.6 (m, 2H), 7.75 (t, 1H), 7.8 - 7.9 (t, 1H), 8.05 (s, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 9.9 (bs, 1H); Mass Spectrum MH⁺ 507.

10 **Example 25.6**

4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline

Obtained by reacting 3-(4-methylpiperazin-1-yl)propanol with 4-(3-chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline (reference example 4.8) in 13% yield; NMR

- 15 spectrum (DMSO-d₆, 373K) 2.1 - 2.2 (m, 2H), 2.2 (s, 3H), 2.25 - 2.35 (m, 4H), 2.35 - 2.45 (m, 4H), 2.5 - 2.6 (m, 2H), 4.4 - 4.45 (t, 4H), 5.3 (s, 2H), 7.1 (d, 1H), 7.25 (d, 1H), 7.3 - 7.35 (m, 1H), 7.35 (d, 1H), 7.55 - 7.65 (m, 2H), 7.7 - 7.75 (t, 1H), 7.8 - 7.85 (t, 1H), 8.05 (d, 1H), 8.5 (s, 1H), 8.6 (d, 1H), 9.9 (bs, 1H); Mass Spectrum MH⁺ 520.

Example 25.7

- 20 **4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(1-methylpiperidin-2-ylmethoxy)quinazoline**

Obtained by reacting (1-methylpiperidin-2-yl)methanol with 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 17% yield; NMR spectrum (DMSO-d₆, 373K) 1.3 - 1.6 (m, 4H), 1.6 - 1.8 (m, 2H), 2.2 - 2.3 (dt, 1H), 2.35 (s, 3H), 2.4 - 2.6 (m, 2H), 4.2 (dd, 1H), 4.5 (dd, 1H), 5.25 (s, 1H), 7.1 (d, 1H), 7.1 - 7.15 (m, 1H), 25 7.3 (m, 1H), 7.25 - 7.4 (m, 3H), 7.4 - 7.5 (m, 1H), 7.7 - 7.8 (t, 1H), 7.9 - 7.95 (dd, 1H), 8.0 (d, 1H), 8.5 (s, 1H), 10.4 (bs, 1H); Mass Spectrum MH⁺ 508.

Example 25.8**4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(2-pyrrolidinoethoxy)quinazoline**

Obtained by reacting 2-pyrrolidinoethanol with 4-(3-chloro-4-(3-

- 30 fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 22% yield; NMR spectrum (DMSO-d₆, 373K) 1.6 - 1.7 (m, 4H), 2.6 - 2.7 (m, 4H), 3.0 - 3.1 (m, 2H), 4.4 - 4.5

(m, 2H), 5.3 (s, 2H), 7.1 - 7.2 (m, 2H), 7.3 (d, 1H), 7.3 - 7.4 (m, 3H), 7.45 - 7.5 (m, 1H), 7.7 - 7.8 (m, 2H), 8.0 (d, 1H), 8.5 (s, 1H) 10.3 (bs, 1H); Mass Spectrum MH^+ 493.

Example 25.9

4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(3-morpholinopropoxy)quinazoline

- 5 Obtained by reacting 3-morpholinopropanol with 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 22% yield; NMR spectrum (DMSO- d_6 , 373K) 2.1 - 2.2 (m, 2H), 2.4 - 2.45 (m, 4H), 2.5 - 2.6 (m, 2H), 3.55 - 3.6 (m, 4H), 4.4 - 4.45 (t, 2H), 5.3 (s, 2H), 7.1 - 7.2 (m, 2H), 7.3 (d, 1H), 7.3 - 7.4 (m, 3H), 7.45 - 7.5 (m, 1H), 7.6 - 7.7 (m, 1H), 7.8 - 7.9 (t, 1H), 8.0 (d, 1H), 8.55 (s, 1H) 10.0 (bs, 1H); Mass Spectrum MH^+ 524.

Example 25.10

4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-[2-(1-methylpyrrolidin-2-yl)ethoxy]quinazoline

- 15 Obtained by reacting 2-(1-methylpyrrolidin-2-yl)ethanol with 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 22% yield; Mass Spectrum MH^+ 508.

Example 25.11

4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(2-morpholinoethoxy)quinazoline

- 20 Obtained by reacting 2-morpholinoethanol with 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 25% yield; NMR spectrum (DMSO- d_6 , 373K) 2.5 - 2.6 (m, 4H), 2.9 - 3.0 (m, 2H), 3.5 - 3.6 (m, 4H), 4.4 - 4.5 (m, 2H), 5.3 (s, 2H), 7.1 - 7.2 (m, 2H), 7.3 (d, 1H), 7.3 - 7.35 (m, 3H), 7.35 - 7.4 (m, 1H), 7.7 - 7.8 (m, 2H), 8.0 (d, 1H), 8.5 (s, 1H), 10.2 (bs, 1H); Mass Spectrum MH^+ 510.

Example 25.12

- 25 **4-(3-Chloro-4-(3-fluorobenzoyloxy)anilino)-5-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline**

Obtained by reacting 3-(4-methylpiperazin-1-yl)propanol with 4-(3-chloro-4-(3-fluorobenzoyloxy)anilino)-5-fluoroquinazoline (reference example 33) in 13% yield; Mass Spectrum MH^+ 537.

30 **Example 26**

The compounds in Table 1 were prepared as follows:

A stock solution of 4-(3-chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) (335 mg) in DMF (7.92 ml) was prepared, and 100 μ l aliquots transferred to a 96 well plate. Using a resin loader, potassium carbonate (300mg) was distributed evenly across the plate. An appropriate benzyl or heteroarylmethyl chloride (0.11 mM) were dissolved in DMF (1 ml) and 100 μ l of each solution transferred to the plate. The plate was agitated at 25°C for 16 hours. To each well was added DMF (100 μ L) and trisamine scavenger resin (726 mg) by using the resin loader to distribute evenly across the wells. The plate was agitated for 3 hours at 25°C. Each well was filtered to remove resin and inorganic material and the remaining filtrates were concentrated *in vacuo*. DMSO (550 μ l) was added to each well and aliquots of 50 μ l were then taken from each well for LCMS purity determination. LCMS purity was determined on a Phenomenex Synergi column (reverse phase silica, 50 x 2 mm, flow rate 1.1 ml/minute), eluting with acetonitrile-water containing formic acid (0.05%) on a gradient from 5-95% over 4.5 minutes, with UV detection at 254 nm. There was thus obtained the compound shown in bold in Table 1.

15 **Table 1**

In Table 1 EG refers to Example, RT refers to the LCMS retention time (minutes)

EG	<u>Compound</u>	<i>M-H⁺</i>	RT
26.1	4-(3-Chloro-4-(2, 6-dichlorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2,6-dichlorobenzyl chloride	542	1.40
26.2	4-(3-Chloro-4-(4-fluorobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 4-fluorobenzyl chloride	492	1.30
26.3	4-(3-Chloro-4-(3-nitrobenzoyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 3-nitrobenzyl chloride	519	1.29
26.4	4-(3-Chloro-4-(3-pyridylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 3-picolyl chloride	475	0.81

EG	Compound	$M-H^+$	RT
26.5	4-(4-(Benzo(1,3)dioxol-5-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with benzo(1,3)dioxol-5-ylmethyl chloride	518	1.26
26.6	4-(3-Chloro-4-(2-methoxybenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-methoxybenzyl chloride	504	1.31
26.7	4-(3-Chloro-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 5-methylisoxazol-3-ylmethyl chloride	479	1.05
26.8	4-(3-Chloro-4-(2-chlorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-chlorobenzyl chloride	508	1.37
26.9	4-(3-Chloro-4-(2-chloro-6-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-chloro-6-fluorobenzyl chloride	526	1.34
26.10	4-(3-Chloro-4-(2,5-dimethylbenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2, 5-dimethylbenzyl chloride	503 (M^+)	1.45
26.11	4-(3-Chloro-4-(3-methoxybenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 3-methoxybenzyl chloride	504	1.26
26.12	4-(3-Chloro-4-(2-nitrobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-nitrobenzyl chloride	519	1.27
26.13	4-(3-Chloro-4-(4-pyridylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 4-picoly l chloride	474	0.74

EG	<u>Compound</u>	<i>M-H</i> ⁺	RT
26.14	4-(3-Chloro-4-(2,6-difluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2,6-difluorobenzyl chloride	509	1.27
26.15	4-(3-Chloro-4-(2-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-fluorobenzyl chloride	492	1.28
26.16	4-(3-Chloro-4-(3-chlorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 3-chlorobenzyl chloride	508	1.39
26.17	4-(3-Chloro-4-(3-methylbenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 3-methylbenzyl chloride	488	1.37
26.18	4-(3-Chloro-4-(5-chlorothiophen-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 5-chlorothiophen-2-ylmethyl chloride	514	1.39
26.19	4-(3-Chloro-4-(2-cyanobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-cyanobenzyl chloride	500 (<i>M</i> ⁺)	1.20
26.20	4-(3-Chloro-4-(2-methylthiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 2-methylthiazol-4-ylmethyl chloride	495	1.06
26.21	4-(3-Chloro-4-(4-methyl-2-nitrobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 4-methyl-2-nitrobenzyl chloride	533	1.37
26.22	4-(3-Chloro-4-(thiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 4-chloromethylthiazole	481	0.97

EG	Compound	M-H ⁺	RT
26.23	4-(3-Chloro-4-(6-chloropyrid-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline Obtained by reaction with 6-chloropyrid-3-ylmethyl chloride	509	1.13

Example 27**Pharmaceutical composition**

The following illustrates a representative pharmaceutical dosage forms of the invention as defined herein (the active ingredient being termed "Compound X"), for therapeutic or prophylactic use in humans:

10	(a) Tablet I	mg/tablet
	Compound X.....	100
	Lactose Ph.Eur.....	182.75
	Croscarmellose sodium.....	12.0
	Maize starch paste (5% w/v paste).....	2.25
	Magnesium stearate.....	3.0
15	(b) Injection I	(50 mg/ml)
	Compound X.....	5.0% w/v
	1M Sodium hydroxide solution.....	15.0% v/v
	0.1M Hydrochloric acid (to adjust pH to 7.6)	
	Polyethylene glycol 400.....	4.5% w/v
	Water for injection to 100%.	

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. For example the tablet may be prepared by blending the components together and compressing the mixture into a tablet.

Starting Materials

The starting materials used in the examples were prepared as follows.

5 Reference Example 1

3,4-Dihydro-5-(1-methylpiperidin-4-yloxy)quinazolin-4-one

Sodium hydride (4.1 g, 60%) was added in portions to 4-hydroxy-1-methylpiperidine (10.7 g) in DMA (125 ml). The reaction was stirred at room temperature for 15 minutes, 50°C for 15 minutes then allowed to cool to room temperature. 5-Fluoro-3,4-dihydroquinazolin-4-one (5.1 g) was added in a single portion, and the mixture heated at 80°C for 2 hours. The reaction was cooled, concentrated *in vacuo* and the residue dissolved in methanol. DOWEX H⁺ ion exchange resin (75 g) was added and the mixture was stirred at room temperature for 1 hour. The mixture was then filtered and the resin washed with methanol. The resin was then suspended in ammonia (7N solution in methanol) and this mixture was stirred at room temperature for 1 hour. The mixture was then filtered, and the filtrate was concentrated *in vacuo* to give the title compound as a white solid after trituration with ether (7.3 g, 91%); NMR spectrum (DMSO-d₆) 1.72 (m, 2H), 1.88 (m, 2H), 2.15 (s, 3H), 2.19 (m, 2H), 2.63 (m, 2H), 4.46 (m, 1H), 7.00 (d, 1H), 7.14 (d, 1H), 7.61 (t, 1H), 7.91 (s, 1H), 11.75 (bs, 1H).

The procedure described above was repeated using the appropriate alcohol. Thus was obtained the compound described below:

Reference Example 1.1

1-(tert-Butoxycarbonylpiperidin-4-yloxy)-3,4-dihydroquinazolin-4-one

Obtained from 1-(tert-butoxycarbonyl-4-hydroxypiperidine in 87% yield; NMR spectrum (DMSO-d₆) 1.39 (s, 9H), 1.6 - 1.87 (m, 4H), 3.32 - 3.43 (m, 2H), 3.47 - 3.60 (m, 2H), 4.75 (m, 1H), 7.08 (d, 1H), 7.17 (d, 1H), 7.64 (t, 1H) 8.84 (s, 1H), 11.80 (bs, 1H); Mass spectrum MH⁺ 346.

Reference Example 1.2

3,4-Dihydro-5-(tetrahydropyran-4-yloxy)quinazolin-4-one

Obtained from tetrahydropyran-4-ol in 27% yield; NMR Spectrum (DMSO-d₆) 1.61 - 1.78 (m, 2H), 1.84 - 2.00 (m, 2H), 3.42 - 3.56 (m, 2H), 3.85 - 3.97 (m, 2H), 4.65 - 4.78 (m, 1H), 7.04 (d, 1H), 7.18 (d, 1H), 7.61 (t, 1H), 7.93 (s, 1H); Mass spectrum MH⁺ 247.

Reference Example 1.3**3,4-Dihydro-5-(tetrahydrofuran-3-yloxy)quinazolin-4-one**

- 5 Obtained from tetrahydrofuran-3-ol in 71% yield; NMR Spectrum (DMSO-d₆) 1.98 - 2.08 (m, 1H), 2.16 - 2.26 (m, 1H), 3.75 - 3.98 (m, 4H), 5.02 - 5.10 (m, 1H), 6.96 (d, 1H), 7.18 (d, 1H), 7.61 (t, 1H), 7.93 (s, 1H), 11.8 (s, 1H); Mass spectrum MH⁺ 233.

Reference Example 1.4**3,4-Dihydro-5-(1,4-dioxaspiro[4.5]dec-8-yloxy)quinazolin-4-one**

- 10 Obtained from 1, 4-dioxaspiro[4.5]decan-8-ol (reference example 46) in 45 yield; NMR Spectrum (DMSO-d₆) 1.48 (m, 2H), 1.70 - 2.00 (m, 6H), 3.85 (s, 4H), 4.63 (m, 1H), 7.03 (d, 1H), 7.14 (d, 1H), 7.60 (t, 1H), 7.90 (s, 1H), 11.74 (bs, 1H); Mass Spectrum M-H⁺ 301.

Reference Example 2

- 15 **4-Chloro-5-(1-methylpiperidin-4-yloxy)quinazoline**

Phosphorus oxychloride (3.59 ml) was added to a solution of 5-(1-methylpiperidin-4-yloxy)-3,4-dihydroquinazolin-4-one (1.0 g) (Reference Example 1) and di-*iso*-propylethylamine (2.01 ml) in DCM (70 ml), and the resulting solution heated at reflux for 16 hours. The reaction was cooled and concentrated *in vacuo* and the residue dissolved in ethyl acetate and cooled to 0°C. Cold saturated aqueous sodium hydrogen carbonate solution was added and the two-phase mixture stirred at 0°C for 15 minutes. The organic layer was separated, dried and concentrated *in vacuo* to yield the title compound as a yellow solid (0.665 g, 62%); NMR spectrum (CDCl₃) 2.10 (m, 2H), 2.23 (m, 2H), 2.42 (s, 3H), 2.60 (m, 2H), 2.84 (m, 2H), 4.73 (m, 1H), 7.04 (d, 1H), 7.62 (d, 1H), 7.81 (t, 1H), 8.93 (s, 1H); Mass spectrum M⁺ 278.

The procedure described above was repeated using the appropriate 3,4-dihydroquinazolin-4-one. Thus was obtained the compound described below:

Reference Example 2.1**5-(1-*tert*-Butoxycarbonylpiperidin-4-yloxy)-4-chloroquinazoline**

- 30 Obtained from 5-(1-*tert*-butoxycarbonylpiperidin-4-yloxy)-3,4-dihydroquinazoline (reference example 1.1) in 66% yield; NMR spectrum (DMSO-d₆) 1.38 (s, 9H), 1.58 - 1.90 (m, 4H), 3.30 - 3.60 (m, 4H), 4.82 (m, 1H), 7.14 - 7.28 (m, 2H), 7.74 (t, 1H), 8.33 (s, 1H).

Reference Example 3**4-Chloro-5-fluoroquinazoline hydrochloride**

To a suspension of 5-fluoro-3,4-dihydroquinazolin-4-one (0.5 g) in thionyl chloride (5 ml) was added DMF (0.2 ml). The mixture was heated at reflux under an atmosphere of nitrogen for 3 hours. The mixture was evaporated *in vacuo*, the residue re-suspended in dry toluene, and evaporated again. The residue was dried *in vacuo* to give the title compound as a pale yellow solid (634 mg, 95%), which was used without further manipulation.

Reference Example 4**4-(4-(Azepan-1-ylcarbonyl)-3-chloroanilino)-5-fluoroquinazoline hydrochloride**

To a solution of 4-(azepan-1-ylcarbonyl)-3-chloroaniline (reference example 11.1) (126.4 mg) in IPA (3 ml) was added 4-chloro-5-fluoroquinazoline (reference example 3) (131 mg). The mixture was heated at reflux, under an atmosphere of nitrogen, for 1 hour producing a precipitate. The mixture was cooled to room temperature and the product filtered off, washed with IPA, diethyl ether and dried *in vacuo* to yield the title compound (199 mg, 100%); NMR spectrum (DMSO-d₆) 1.54 (bs, 6H), 1.72 (bs, 2H), 3.21 (bs, 2H), 3.58 (bs, 2H), 7.46 (d, 1H), 7.62 - 7.72 (m, 2H), 7.81 (d, 1H), 7.91 (s, 1H), 8.07 (m, 1H), 8.89 (s, 1H); Mass spectrum MH⁺ 399.

The procedure described above was repeated using the appropriate aniline and 4-chloroquinazoline. Thus was obtained the compound described below:

Reference Example 4.1**4-(1-(3-Fluorobenzyl)indazol-5-ylamino)-5-fluoroquinazoline hydrochloride**

Obtained from 4-chloro-5-fluoroquinazoline hydrochloride (reference example 3) and 5-amino-1-(3-fluorobenzyl)indazole (obtained as described in PCT Int. Appl. WO9802438) in 66% yield; NMR spectrum (DMSO-d₆) 5.74 (s, 2H), 7.00 - 7.15 (m, 3H), 7.36 (m, 1H), 7.54 (dd, 1H), 7.67 (dd, 1H), 7.77 (d, 1H), 7.82 (d, 1H), 7.97 (s, 1H), 8.06 (m, 1H), 8.22 (s, 1H), 8.77, (s, 1H); Mass spectrum MH⁺ 383.

Reference Example 4.2**4-(3-Chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-chloro-4-hydroxyaniline in 60% yield; NMR spectrum (DMSO-d₆) 1.9 (m, 2H), 2.1 (m, 5H), 2.3 (m, 2H), 2.6 (m, 2H), 4.8 (m, 1H), 7.0 (d, 1H), 7.2 (d, 1H), 7.3 (m, 2H), 7.7 (m, 1H), 8.0 (d, 1H), 8.4 (s, 1H), 10.0 (s, 1H); Mass spectrum MH⁺ 385.

Reference Example 4.3**4-(3-Fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline hydrochloride**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 3-fluoro-4-hydroxyaniline in 87% yield; Mass Spectrum MH^+ 369.

5 Reference Example 4.4**4-(Indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained by reacting 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and 5-aminoindole in 55% yield; NMR spectrum (DMSO- d_6) 1.80 - 2.00 (m, 4H), 2.10 - 2.20 (m + s, 5H), 2.27 (m, 2H), 2.62 (m, 2H), 4.80 (m, 1H), 6.42 (s, 1H), 7.19 (d, 1H),
10 7.28 (m, 2H), 7.35 (m, 1H), 7.40 (d, 1H), 7.67 (t, 1H), 8.06 (s, 1H), 8.42 (s, 1H); Mass Spectrum MH^+ 374.

Reference Example 4.5**2-[2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetonitrile hydrochloride**

15 Obtained from 2-(4-amino-2-chlorophenoxy)acetonitrile (reference example 9.4) and 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) in 96 % yield; NMR spectrum (DMSO- d_6) 2.10 - 2.32 (m, 2H), 2.32 - 2.50 (m, 2H), 2.76 (s, 3H), 3.03 - 3.25 (m, 2H), 3.40 - 3.60 (m, 2H), 4.98 (m, 1H), 5.28 (s, 2H), 7.28 (d, 1H), 7.33 (d, 1H), 7.38 (d, 1H), 7.65 (d, 1H), 7.76 (dd, 1H), 8.21 (d, 1H), 8.55 (s, 1H), 9.95 (bs, 1H); Mass Spectrum MH^+
20 424.

Reference Example 4.6**4-(4-Iodoanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-iodoaniline and 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) in 67% yield; NMR spectrum (DMSO- d_6) 1.8 - 2.0 (m, 2H), 2.1 - 2.3
25 (m, 4H), 2.17 (s, 3H), 2.6 - 2.7 (m, 2H), 4.7 - 4.85 (m, 1H), 7.2 - 7.25 (d, 1H), 7.35 - 7.4 (d, 1H), 7.6 - 7.8 (m, 5H), 8.5 (s, 1H), 10.15 - 10.2 (bs, 1H); Mass spectrum MH^+ 461.

Reference Example 4.7**4-(5-Chloro-2-fluoro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline**

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example
30 2) and 4-amino-2-chloro-5-fluorophenol (reference example 8.7) in 83% yield; Mass Spectrum MH^+ 495.

Reference Example 4.8**4-(3-Chloro-4-(2-pyridylmethoxy)anilino)-5-fluoroquinazoline**

Obtained from 4-chloro-5-fluoroquinazoline hydrochloride (reference example 3) and 3-chloro-4-(2-pyridylmethoxy)aniline (obtained as described in PCT Int. Appl. (1996) WO 9615118) in 78% yield; Mass spectrum MH^+ 381.

Reference Example 5**3-Chloro-4-(3-pyridyloxy)aniline**

Titanium (III) chloride (10 wt. % in 20-30% aq. HCl, 6.5 ml) was added to a solution of 3-chloro-1-nitro-4-(3-pyridyloxy)benzene (0.25 g) (reference example 14) in acetone (18 ml) and water (3.6 ml) containing ammonium acetate (1.6 g) and the solution stirred overnight at room temperature. The reaction was concentrated *in vacuo* and the residue partitioned between ethyl acetate and 880 ammonia solution. Combined organic extracts were dried and concentrated to give the title compound (0.15 g, 68%); Mass spectrum MH^+ 221.

The procedure described above was repeated using the appropriate nitro compound.

Thus were obtained the compounds described below:

Reference Example 5.1**3-Chloro-4-(3-fluorophenoxy)aniline**

Obtained from 3-chloro-4-(3-fluorophenoxy)-1-nitrobenzene (reference example 14.1) in 65% yield; Mass spectrum MH^+ 238.

Reference Example 5.2**3-Chloro-4-(2,3-difluorophenoxy)aniline**

Obtained from 3-chloro-4-(2,3-difluorophenoxy)-1-nitrobenzene (reference example 14.2) in 62% yield; Mass spectrum MH^+ 256.

Reference Example 5.3**3-Methyl-4-(N-(2-pyridylmethyl)amino)aniline**

Obtained from 3-methyl-4-(N-(2-pyridylmethyl)amino)nitrobenzene (reference example 15.1) in 71% yield; NMR Spectrum (DMSO- d_6) 2.07 (s, 3H), 4.13 (s, 2H), 4.17 (d, 2H), 4.88 (t, 1H), 6.14 (d, 1H), 6.21 (dd, 1H), 6.36 (dd, 1H), 7.20 (dd, 1H), 7.32 (d, 1H), 7.68 (ddd, 1H), 8.50 (d, 1H); Mass spectrum MH^+ 214.

Reference Example 5.4**3-Chloro-4-[N-methyl-N-(2-pyridyl)amino]aniline**

Obtained from 3-chloro-4-[N-methyl-N-(2-pyridyl)amino]nitrobenzene (reference example 16) in 95% yield; NMR Spectrum (DMSO-d₆) 3.23 (s, 3H), 5.50 (s, 2H), 6.04 (d, 1H), 6.57 (dd, 1H), 6.59 (dd, 1H), 6.74 (d, 1H), 7.01 (d, 1H), 7.35 (ddd, 1H), 8.08 (d, 1H); Mass spectrum MH⁺ 236.

5 **Reference Example 5.5**

3-Chloro-4-(2-pyridyl)amino)aniline

Obtained from 3-chloro-4-(2-pyridylamino)nitrobenzene (reference example 16.1) in 60% yield; NMR Spectrum (DMSO-d₆) 5.23 (s, 2H), 6.44 (d, 1H), 6.51 (dd, 1H), 6.58 (dd, 1H), 6.68 (dd, 1H), 7.16 (d, 1H), 7.42 (ddd, 1H), 7.92 (s, 1H), 7.97 (d, 1H); Mass spectrum MH⁺ 222.

Reference Example 5.6

3-Methyl-4-(2-pyridylamino)aniline

Obtained from 3-methyl-4-(2-pyridylamino)nitrobenzene (reference example 16.2) in 96% yield; NMR Spectrum (DMSO-d₆) 2.01 (s, 3H), 4.88 (s, 2H), 6.28 (d, 1H), 6.39 (dd, 1H), 6.46 (d, 1H), 6.51 (dd, 1H), 6.87 (d, 1H), 7.37 (ddd, 1H), 7.73 (s, 1H), 7.95 (d, 1H); Mass spectrum MH⁺ 200.

Reference Example 5.7

3-Methyl-4-[N-methyl-N-(2-pyridyl)amino]aniline

Obtained from 3-methyl-4-[N-methyl-N-(2-pyridyl)amino]nitrobenzene (reference example 16.3) in 89% yield; NMR Spectrum (DMSO-d₆) 1.90 (s, 3H), 3.22 (s, 3H), 5.09 (s, 2H), 5.96 (d, 1H), 6.46 (dd, 1H), 6.48 (dd, 1H), 6.50 (d, 1H), 6.80 (d, 1H), 7.30 (ddd, 1H), 8.08 (d, 1H); Mass spectrum MH⁺ 214.

Reference Example 6

5-Amino-3-chloro-1-(2-pyridylmethyl)indole

25 A solution of 3-chloro-5-nitro-1-(2-pyridylmethyl)indole (reference example 13) (2.5 g) in ethanol (130 ml) was stirred at room temperature. Sodium dithionite (7.6 g) in water (18 ml) was added, and the mixture was heated to 50°C for 5 hours, then cooled to room temperature. The ethanol was removed *in vacuo*, and the residue was partitioned between DCM and water. The DCM layer was separated, dried over sodium sulphate, then concentrated
30 *in vacuo* to give the crude material, which was purified by column chromatography using 50% DCM in hexane then neat DCM to give the title compound as an orange solid (488 mg, 23%); NMR Spectrum (CDCl₃) 3.53 (s, 2H), 5.27 (s, 2H), 6.61 (dd, 1H), 6.68 (d, 1H), 6.86 (d, 1H),

7.01 (d, 1H), 7.04 (s, 1H), 7.11 (dd, 1H), 7.47 (dt, 1H), 8.55 (m, 1H); Mass Spectrum MH^+ 258.

The procedure described above was repeated using the appropriate aryl nitro compound. Thus were obtained the compounds described below:

5 **Reference Example 6.1**

5-Amino-3-chloro-1-(2-pyridylmethyl)indazole

Obtained from 3-chloro-5-nitro-1-(2-pyridylmethyl)indazole (reference example 13.1) in 24% yield; NMR Spectrum ($CDCl_3$) 3.3 (bs, 2H), 6.65 (dd, 1H), 6.77 (m, 1H), 6.84 - 7.02 (m, 5H), 7.24 (m, 1H).

10 **Reference Example 6.2**

3-Fluoro-4-(1-methyl-1H-imidazol-2-ylthio)aniline

Obtained from 3-fluoro-4-(1-methyl-1H-imidazol-2-ylthio)nitrobenzene (reference example 43) in 86% yield; Mass spectrum MH^+ 224.

Reference Example 6.3

15 **3-Fluoro-4-(1-methyl-1H-1,3,4-triazol-2-ylthio)aniline**

Obtained from 3-fluoro-4-(1-methyl-1H-1,3,4-triazol-2-ylthio)nitrobenzene (reference example 43.1) in 14% yield; Mass spectrum MH^+ 225.

Reference Example 6.4

3-Chloro-4-(2-pyridylthio)aniline

20 Obtained from 3-chloro-4-(2-pyridylthio)nitrobenzene (reference example 44) in 21% yield; Mass spectrum MH^+ 237.

Reference Example 6.5

3-Chloro-4-(2-pyrimidinylthio)aniline

25 Obtained from 3-chloro-4-(2-pyrimidinylthio)nitrobenzene (reference example 44.1) in 27%; Mass spectrum MH^+ 238.

Reference Example 7

5-Amino-1-benzenesulphonylindole

1-Benzenesulphonyl-5-nitroindole (reference example 26) (0.3 g) and 10% Pd/C (30 mg) in ethyl acetate / ethanol (9:1, 20 ml) were stirred under a hydrogen atmosphere for 2
30 hours. The reaction was filtered and concentrated *in vacuo* to give the title compound as a beige solid (0.18 g, 67%); NMR spectrum ($DMSO-d_6$) 4.9 (bs, 2H), 6.6 (m, 3H), 7.6 (m, 5H), 7.8 (d, 2H); Mass spectrum MH^+ 273.

The procedure described above was repeated using the appropriate nitro aryl compound. Thus was obtained the compound described below:

Reference Example 7.1

5-Amino-1-benzylindole

- 5 Obtained from 1-benzyl-5-nitroindole (reference example 26.1) in 61% yield; NMR spectrum (DMSO-d₆) 4.5 (bs, 2H), 5.2 (s, 2H), 6.2 (d, 1H), 6.4 (dd, 1H), 6.6 (s, 1H), 7.1 (d, 1H), 7.2 (d, 2H), 7.3 (m, 4H); Mass spectrum MH⁺ 223.

Reference Example 8

3-Chloro-4-phenoxyaniline

- 10 To a solution of 2-chloro-4-nitro-1-phenoxybenzene (reference example 14.3) (1.76 g) in ethyl acetate / ethanol (105 ml, 9.5:1) was added 10% Pt/C (135 mg). The resulting solution was subjected to a hydrogen atmosphere for 18 hours at room temperature. The catalyst was then filtered off and the solvent concentrated *in vacuo* to give a yellow oil which was purified by flash chromatography eluting with 10 - 15% ethyl acetate / *iso*-hexane to give the title
15 compound as a clear oil (1.15 g, 74%); NMR Spectrum (DMSO-d₆) 5.28 (bs, 2H), 6.54 (dd, 1H), 6.71 (d, 1H), 6.78 (d, 2H), 6.88 (d, 1H), 6.98 (tt, 1H), 7.28 (t, 2H); Mass Spectrum MH⁺ 220.

The procedure described above was repeated using the appropriate nitro compound. Thus were obtained the compounds described below:

- 20 **Reference Example 8.1**

3-Chloro-4-(3-fluorobenzoyloxy)aniline

Obtained from 2-chloro-1-(3-fluorobenzoyloxy)-4-nitrobenzene (reference example 18) in 75% yield; NMR Spectrum (DMSO-d₆) 4.91 (s, 2H), 5.01 (s, 2H), 6.45 (dd, 1H), 6.63 (s, 1H), 6.89 (d, 1H), 7.12 (t, 1H), 7.24 (t, 2H), 7.40 (m, 1H); Mass Spectrum MH⁺ 252.

- 25 **Reference Example 8.2**

tert-Butyl 4-amino-2-chlorobenzoate

Obtained from *tert*-butyl 2-chloro-4-nitrobenzoate (reference example 20.1) in 75% yield; NMR Spectrum (CDCl₃) 1.58 (s, 9H), 3.99 (s, 2H), 6.51 (dd, 1H), 6.66 (d, 1H), 7.68 (d, 1H); Mass Spectrum M⁺ 227.

- 30 **Reference Example 8.3**

3-(2-Chloro-4-aminophenoxyethyl)-1,5-dimethylpyrazole

Obtained from 3-(2-chloro-4-nitrophenoxymethyl)-1,5-dimethylpyrazole (reference example 40) in 99% yield; NMR data (DMSO-d₆) 2.3 (s, 3H), 3.7 (s, 3H), 4.9 (s, 2H), 5.2 (bs, 2H), 6.1 (s, 1H), 6.5 (m, 1H), 6.7 (d, 1H), 7.0 (d, 1H), 8.6 (bs, 1H); Mass spectrum MH⁺ 252.

Reference Example 8.4

5 **3-(2-Chloro-4-aminophenoxymethyl)-1-methylpyrazole**

Obtained from 3-(2-chloro-4-nitrophenoxymethyl)-1-methylpyrazole (reference example 40.1) in 87% yield; NMR data (DMSO-d₆) 3.8 (s, 3H), 4.9 (s, 4H), 6.3 (d, 1H), 6.4 (m, 1H), 6.6 (d, 1H), 6.9 (d, 1H); Mass spectrum MH⁺ 252.

Reference Example 8.5

10 **3-Fluoro-4-(1H-imidazol-2-ylthio)aniline**

Obtained from 3-fluoro-4-(1H-imidazol-2-ylthio)nitrobenzene (reference example 44.3) in 98% yield.

Reference Example 8.6

4-(Thiazol-2-ylsulphonyl)aniline

15 Obtained from 4-(thiazol-2-ylsulphonyl)nitrobenzene (reference example 45) in 79% yield; Mass spectrum MH⁺ 240.

Reference Example 8.7

4-Amino-2-chloro-5-fluorophenol

Obtained from 2-chloro-5-fluoro-4-nitrophenol (reference example 49) in 100% yield;
20 NMR spectrum (CDCl₃) 4.2 - 4.8 (bs, 2H), 6.6 - 6.7 (d, 1H), 6.75 - 6.8 (d, 1H), 9.3 (s, 1H).

Reference Example 9

tert-Butyl 4-amino-2-ethynylbenzoate

To a stirred suspension of *tert*-butyl 2-ethynyl-4-nitrobenzoate (reference example 22) (8.76 g) in a mixture of acetic acid (110 ml) and water (11 ml) was added iron (reduced by
25 hydrogen, 8.63 g) in portions, with cooling in a cold water bath. The mixture was stirred at room temperature for 3 hours, diluted with ethyl acetate and filtered. The filtrate was evaporated *in vacuo*, the residue treated with aqueous sodium hydrogen carbonate and extracted with DCM. The organic extract was dried and evaporated *in vacuo* to give a brown solid. This was purified by column chromatography using DCM / ethyl acetate (97/3 to 95/5)
30 as eluent to give the title compound (7.30 g, 95%); NMR spectrum (CDCl₃) 1.59 (s, 9H), 3.30 (s, 1H), 3.97 (s, 2H), 6.60 (dd, 1H), 6.83 (d, 1H), 7.77 (d, 1H).

The procedure described above was repeated using the appropriate nitro compound. Thus were obtained the compounds described below:

Reference Example 9.1

3-Chloro-4-(2-pyridylmethyl)amino)aniline

- 5 Obtained from 3-chloro-4-(2-pyridylmethyl)amino)nitrobenzene (reference example 15) in 58% yield; NMR Spectrum (DMSO-d₆) 4.34 (d, 2H), 4.54 (s, 2H), 5.36 (t, 1H), 6.37 (m, 2H), 6.60 (s, 1H), 7.26 (dd, 1H), 7.33 (d, 1H), 7.73 (dd, 1H), 8.54 (d, 1H); Mass spectrum MH⁺ 236.

Reference Example 9.2

10 **3-Chloro-4-(2-pyrimidinylmethoxy)aniline**

Obtained from 3-chloro-4-(2-pyrimidinylmethoxy)nitrobenzene (reference example 14.4) in 93% yield; Mass spectrum M-H⁺ 266.

Reference Example 9.3

4-(2-Aminothiazol-4-ylmethoxy)-3-chloroaniline

- 15 Obtained from 4-(2-aminothiazol-4-ylmethoxy)-3-chloronitrobenzene (reference example 34) (80 mg) in 85% yield; NMR spectrum (DMSO-d₆) 4.74 (s, 2H), 4.89 (bs, 2H), 6.46 (dd, 1H), 6.51 (s, 1H), 6.63 (d, 1H), 6.91 (d, 1H), 6.92 (s, 2H); Mass spectrum MH⁺ 256.

Reference Example 9.4

2-(4-Amino-2-chlorophenoxy)acetonitrile

- 20 Obtained from 2-(2-chloro-4-nitrophenoxy)acetonitrile (reference example 18.6) in 96% yield; NMR spectrum (DMSO-d₆) 4.99 (s, 2H), 5.11 (bs, 2H), 6.49 (dd, 1H), 6.64 (d, 1H), 6.98 (d, 1H).

Reference Example 9.5

5-(2-Chloro-4-aminophenoxy)methyl)-3-methylisoxazole

- 25 Obtained from 5-(2-chloro-4-nitrophenoxy)methyl)-3-methylisoxazole (reference example 40.2) in 8% yield; NMR spectrum (DMSO-d₆) 2.2 (s, 3H), 5.0 (s, 2H), 5.1 (s, 2H), 6.4 (s, 1H), 6.5 (m, 1H), 6.6 (d, 1H), 6.9 (d, 1H); Mass spectrum MH⁺ 239.

Reference Example 9.6

3-Chloro-4-(1H-imidazol-2-ylthio)aniline

- 30 Obtained from 3-chloro-4-(1H-imidazol-2-ylthio)nitrobenzene (reference example 44.2) in 76% yield.

Reference Example 9.7**3-Chloro-4-(2-thiazolylthio)aniline**

Obtained from 3-chloro-4-(2-thiazolylthio)nitrobenzene (reference example 44.5) in
5 56% yield; Mass spectrum MH^+ 243.

Reference Example 10**3-Chloro-4-((3-fluorophenylamino)methyl)aniline**

3-Chloro-4-((3-fluorophenylamino)methyl)nitrobenzene (reference example 19) (50
mg) was dissolved in ethanol (2 ml). Saturated ammonium chloride (0.2 ml) was added, and
10 the mixture heated to reflux. Indium metal powder (100 mesh, 143 mg) was added, and the
mixture stirred at reflux for 1 hour. The mixture was cooled to room temperature, made basic
with concentrated aqueous ammonia solution, and filtered through a bed of diatomaceous earth.
The diatomaceous earth was washed with a 1:1 mixture of DCM and methanol containing 1%
concentrated aqueous ammonia solution (20 ml). The combined filtrates were evaporated, and
15 the residue partitioned between water and DCM. The organic layer was evaporated, and the
residue purified by chromatography, using 15% ethyl acetate in *iso*-hexane as eluent to give the
title compound as an orange oil (20 mg, 45%); NMR spectrum (DMSO- d_6) 4.10 (d, 2H), 5.15
(s, 2H), 6.20 - 6.30 (m, 3H), 6.37 (d, 1H), 6.45 (dd, 1H), 6.60 (d, 1H), 6.95 - 7.05 (m, 2H).

Reference Example 11**3-Chloro-4-(8-quinolylthio)aniline**

To 3-chloro-1-nitro-4-(8-quinolylthio)benzene (reference example 17) (26 g) was
added ethanol (5.0 ml) and tin (II) chloride (2.20 g). This was heated to 80°C for 15 minutes,
then allowed to cool overnight. The mixture was made basic with 880 ammonia solution, and
the product extracted into ethyl acetate (3 x 15 ml), after centrifuging. The solvent was
25 removed *in vacuo* to yield the title compound (0.203 mg, 86%); Mass spectrum MH^+ 286.

The procedure described above was repeated using the appropriate nitro compound. Thus
was obtained the compound described below:

Reference Example 11.1**4-(Azepan-1-ylcarbonyl)-3-chloroaniline**

Obtained from 1-(2-chloro-4-nitrobenzoyl)azepane (reference example 25) in 73%
30 yield; NMR spectrum (CDCl₃) 1.48 - 1.74 (m, 6H), 1.74 - 1.92 (m, 2H), 3.23 - 3.33 (m, 2H),

3.35 - 3.84 (m, 2H), 3.85 (s, 2H), 6.54 (dd, 1H), 6.68 (d, 1H), 7.02 (d, 1H); Mass spectrum MH^+ 253.

Reference Example 12

3-Chloro-5-nitroindole

- 5 *N*-Chlorosuccinimide (1.65 g) was added in portions to a solution of 5-nitroindole (2.00 g) in DMF (20 ml). The resulting solution was stirred at room temperature for 18 hours. The pale brown solution was poured into water (200 ml) to give a yellow precipitate which was filtered, washed with water and dried *in vacuo* to give the title compound as a yellow solid (2.40 g, 99%). Mass spectrum $M-H^+$ 195.

10 Reference Example 13

3-Chloro-5-nitro-1-(2-pyridylmethyl)indole

- Picolyl chloride hydrochloride (3.26 g) was added to a stirred mixture of 3-chloro-5-nitroindole (reference example 12) (1.97 g) and potassium carbonate (13.8 g) in DMF (50 ml). The mixture was heated to 50°C and stirred for 2 hours, after which time the solvent was
15 removed *in vacuo*. The residue was dissolved in DCM, then washed with water, and dried over sodium sulphate to give the product as a yellow solid (2.53 g, 88%); NMR Spectrum ($CDCl_3$) 5.43 (s, 2H), 6.90 (d, 1H), 7.23 (dd, 1H), 7.35 (s, 1H), 7.37 (d, 1H), 7.62 (dt, 1H), 8.12 (dd, 1H), 8.60 (m, 2H).

- The procedure described above was repeated using the appropriate indazole. Thus was
20 obtained the compound described below:

Reference Example 13.1

3-Chloro-5-nitro-1-(2-pyridylmethyl)indazole

Obtained from 3-chloro-5-nitroindazole in 74% yield; NMR Spectrum ($CDCl_3$) 5.32 (s, 2H), 6.80 (d, 1H), 6.89 (d, 1H), 7.01 (dt, 1H), 7.28 (m, 3H), 8.12 (dd, 1H), 8.61 (d, 1H).

25 Reference Example 14

3-Chloro-1-nitro-4-(3-pyridyloxy)benzene

- To a stirred solution of 3-hydroxypyridine (1.05 g) in DMA (25 ml) under a nitrogen atmosphere was added sodium hydride (0.44 g, 60% suspension in oil). The solution was stirred at ambient temperature for two hours. To this solution was added 3-chloro-4-fluoronitrobenzene (1.75 g) and the solution was stirred and heated at 90°C for four hours. The
30 solution was cooled and poured into water (120 ml) and extracted with ethyl acetate. The combined organic extracts were washed with water, brine, and dried with anhydrous

magnesium sulphate. Evaporation of the solvent gave a solid, which was recrystallised from a mixture of ethyl acetate / *iso*-hexane to give the title compound as a yellow solid (2.15 g, 86%); NMR spectrum (CDCl₃) 6.95-6.98 (dd, 1H), 7.4 (d, 2H), 8.1 (dd, 1H), 8.4 (d, 1H), 8.5 - 8.7 (d, 2H); Mass spectrum MH⁺ 251.

- 5 The procedure described above was repeated using 3-chloro-4-fluoronitrobenzene and the appropriate phenol or alcohol. Thus were obtained the compounds described below:

Reference Example 14.1

3-Chloro-4-(3-fluorophenoxy)nitrobenzene

- Obtained from 3-chloro-4-fluoronitrobenzene and 3-fluorophenol in 43% yield; NMR
10 spectrum (CDCl₃) 6.8-6.9 (m, 2H), 6.95-7.0 (m, 2H), 7.3-7.45 (m, 1H), 8.1 (dd, 1H), 8.4 (d, 1H); Mass spectrum M-H⁺ 266.

Reference Example 14.2

3-Chloro-4-(2,3-difluorophenoxy)nitrobenzene

- Obtained from 3-chloro-4-fluoronitrobenzene and 2,3-difluorophenol in 55% yield; NMR
15 spectrum (CDCl₃); 6.8 - 6.9 (dd, 1H), 6.9 - 7.0 (m, 1H), 7.1 - 7.2 (m, 2H), 8.0 - 8.1 (dd, 1H), 8.4 (t, 1H); Mass spectrum M-H⁺ 284.

Reference Example 14.3

3-Chloro-4-phenoxy nitrobenzene

- Obtained from 3-chloro-4-fluoronitrobenzene and phenol in 42% yield; NMR Spectrum
20 (DMSO-d₆) 7.02 (d, 1H), 7.18 (dt, 2H), 7.30 (tt, 1H), 7.49 (t, 2H), 8.16 (dd, 1H), 8.44 (d, 1H); Mass Spectrum M-H⁺ 248.

Reference Example 14.4

3-Chloro-4-(pyrimidin-2-ylmethoxy)nitrobenzene

- Obtained from 3-chloro-4-fluoronitrobenzene and pyrimidine-2-methanol (**reference**
25 **example 27**) in 36% yield; NMR spectrum (DMSO-d₆) 5.60 (s, 2H), 7.30 (d, 1H), 7.47 (t, 1H), 8.14 (dd, 1H), 8.31 (d, 1H), 8.82 (d, 1H); Mass spectrum M-H⁺ 266.

Reference Example 15

3-Chloro-4-(2-pyridyl)amino)nitrobenzene

- A mixture of 3-chloro-4-fluoronitrobenzene (1.76 g), 2-(aminomethyl)pyridine (1.03
30 ml) and di-*iso*-propylethylamine (1.74 ml) was dissolved in acetonitrile (60 ml) and heated overnight at reflux. The mixture was cooled to room temperature and the resulting solid was collected by filtration and washed with diethyl ether, giving the title compound as a yellow

solid (2.21 g, 84%); NMR spectrum (DMSO-d₆) 4.66 (d, 2H), 6.73 (d, 1H), 7.30 (dd, 1H), 7.33 (dd, 1H), 7.59 (t, 1H), 7.78 (ddd, 1H), 8.00 (dd, 1H), 8.19 (d, 1H), 8.56 (d, 1H); Mass spectrum MH⁺ 266.

The procedure described above was repeated using the appropriate amine and fluoronitro
5 compound. Thus was obtained the compound described below:

Reference Example 15.1

3-Methyl-4-(2-pyridylmethyl)amino)nitrobenzene

Obtained from 4-fluoro-3-methylnitrobenzene and 2-(aminomethyl)pyridine in 6%
yield; NMR spectrum (DMSO-d₆) 2.26 (s, 3H), 4.58 (d, 2H), 6.48 (d, 1H), 7.17 (t, 1H), 7.28
10 (dd, 1H), 7.31 (d, 1H), 7.76 (ddd, 1H), 7.88 (dd, 1H), 7.92 (d, 1H), 8.55 (d, 1H); Mass
spectrum MH⁺ 244.

Reference Example 16

3-Chloro-4-[N-methyl-N-(2-pyridyl)amino]nitrobenzene

Sodium hydride (60% dispersion, 200 mg) was suspended in DMA (25 ml), and 2-
15 (methylamino)pyridine (513 µl) was added under a nitrogen atmosphere at room temperature.
The mixture was stirred for 20 minutes at room temperature. 3-Chloro-4-fluoronitrobenzene
(878 mg) was added as a solution in DMA (5 ml), causing a dark red solution to form. The
mixture was stirred overnight at room temperature, quenched by the careful addition of water,
acidified with acetic acid, and the pH adjusted to pH 8 with concentrated aqueous ammonia
20 solution. The mixture was poured into water (200 ml), and extracted with DCM (3 x 100ml).
The combined extractions were filtered through a phase separating filter paper, and evaporated
to give a brown oil, which was purified by chromatography, using 15% ethyl acetate in *iso*-
hexane as eluent to give the title compound as a yellow solid (226mg, 17%); NMR spectrum
(DMSO-d₆) 3.17 (s, 3H), 6.57 (d, 1H), 6.78 (dd, 1H), 7.57 (dd, 1H), 7.70 (d, 1H), 8.11 (d, 1H),
25 8.24 (dd, 1H), 8.37 (d, 1H); Mass spectrum MH⁺ 266.

The procedure described above was repeated using the appropriate fluoronitrobenzene
and aminopyridine. Thus were obtained the compounds described below:

Reference Example 16.1

3-Chloro-4-(2-pyridylamino)nitrobenzene

30 Obtained from 3-chloro-4-fluoronitrobenzene and 2-aminopyridine in 16% yield; NMR
spectrum (DMSO-d₆) 7.04 (dd, 1H), 7.36 (d, 1H), 7.76 (ddd, 1H), 8.16 (dd, 1H), 8.30 (d, 1H),
8.31 (d, 1H), 8.61 (d, 1H), 9.01 (s, 1H); Mass spectrum MH⁺ 252.

Reference Example 16.2**3-Methyl-4 4-(2-pyridylamino)nitrobenzene**

Obtained from 4-fluoro-3-methylnitrobenzene and 2-aminopyridine in 36% yield;

NMR spectrum (DMSO-d₆) 2.39 (s, 3H), 6.92 (dd, 1H), 7.19 (d, 1H), 7.68 (ddd, 1H), 8.00 (dd, 1H), 8.06 (d, 1H), 8.21 (d, 1H), 8.31 (d, 1H), 8.51 (s, 1H); Mass spectrum MH⁺ 230.

Reference Example 16.3**3-Methyl-4-[N-methyl-N-(2-pyridyl)amino]nitrobenzene**

Obtained from 4-fluoro-3-methylnitrobenzene and 2-(methylamino)pyridine in 39% yield; NMR spectrum (DMSO-d₆) 2.12 (s, 3H), 3.32 (s, 3H), 6.36 (d, 1H), 6.71 (dd, 1H), 7.48 (d, 1H), 7.49 (ddd, 1H), 8.10 (m, 2H), 8.20 (d, 1H); Mass spectrum MH⁺ 244.

Reference Example 17**3-Chloro-1-nitro-4-(8-quinolylthio)benzene**

8-Quinolinethiol (0.32 g) was dissolved in DMF (4 ml) under argon. To this was added sodium bis(trimethylsilyl)amine (2.2 ml, 1.0M in THF) and the reaction stirred at room temperature for 1 hour. To this was added a solution of 3-chloro-4-fluoronitrobenzene (0.35 g) in DMF (0.5 ml), and stirring continued at room temperature overnight. The solution was then poured into water (20 ml) and stirred for 30 minutes. The resulting solid was filtered off, washed with water (5 ml) then with a 1:1 mix of ethyl acetate / *iso*-hexane (6 ml) and dried *in vacuo* to yield the title compound (0.26 g, 41%); Mass spectrum MH⁺ 317.

Reference Example 18**2-Chloro-1-(3-fluorobenzyloxy)-4-nitrobenzene**

To a solution of 2-chloro-4-nitrophenol (20.0 g) in acetone (400 ml) was added potassium carbonate (47.76 g) followed by the dropwise addition of 3-fluorobenzylbromide (32.67 g) over 15 minutes. The reaction mixture was then stirred at room temperature for 16 hours and filtered to remove insoluble material. The solvent was then concentrated *in vacuo* and the solid remaining was purified by flash chromatography, using 30 - 80% DCM / *iso*-hexane as eluent to give the title compound (30.96 g, 95%); NMR Spectrum (DMSO-d₆) 5.39 (s, 2H), 7.18 (t, 1H), 7.30 (m, 2H), 7.45 (m, 2H), 8.23 (dd, 1H), 8.32 (d, 1H); Mass Spectrum M-H⁺ 280.

The procedure described above was repeated using the appropriate phenol and alkyl chloride or bromide. Thus were obtained the compounds described below:

Reference Example 18.1**4-(2-Fluorobenzoyloxy)-3-iodo-nitrobenzene**

Obtained from 4-hydroxy-3-iodo-nitrobenzene (obtained as described in *J. Org. Chem.* 1990, 55, 5287-5291) and 2-fluorobenzyl bromide (0.39 g) in 85% yield; Mass Spectrum M-H⁺ 372.

Reference Example 18.2**4-(3-Fluorobenzoyloxy)-3-iodo-nitrobenzene**

Obtained by reacting 4-hydroxy-3-iodo-nitrobenzene and 3-fluorobenzyl bromide in 99% yield; Mass Spectrum M-H⁺ 372.

Reference Example 18.3**4-(2, 6-Difluorobenzoyloxy)-3-iodo-nitrobenzene**

Obtained by reacting 4-hydroxy-3-iodo-nitrobenzene and 2, 6-difluorobenzyl bromide in 100% yield; Mass Spectrum M-H⁺ 390.

Reference Example 18.4**3-Iodo-4-(4-thiazolylmethoxy)-nitrobenzene**

Obtained by reacting 4-hydroxy-3-iodo-nitrobenzene and 4-chloromethylthiazole hydrochloride in 100% yield; Mass Spectrum MH⁺ 363.

Reference Example 18.5**3-Iodo-4-(5-methylisoxazol-3-ylmethoxy)-nitrobenzene**

Obtained by reacting 4-hydroxy-3-iodo-nitrobenzene and 3-chloromethyl-5-methylisoxazole in 74% yield; Mass Spectrum M-H⁺ 359.

Reference Example 18.6**2-(2-Chloro-4-nitrophenoxy)acetonitrile**

Obtained by reacting 2-chloro-4-nitrophenol and chloroacetonitrile in 39% yield; NMR spectrum (CDCl₃) 4.96 (s, 2H), 7.15 (d, 1H), 8.22 (dd, 1H), 8.36 (d, 1H); Mass Spectrum M-H⁺ 211.

Reference Example 19**3-Chloro-4-((3-fluorophenylamino)methyl)nitrobenzene**

2-Chloro-4-nitrobenzyl bromide (501 mg) was dissolved in acetonitrile (30 ml). Di-*iso*-propylethylamine (348 µl) and 3-fluoroaniline (192 µl) were added, and the mixture heated at reflux overnight. The solvent was evaporated, and the residue purified by chromatography, using 15 to 25% ethyl acetate in *iso*-hexane as eluent to give the title compound as a brown oil

(470 mg, 84%); NMR spectrum (DMSO-d₆) 4.42 (d, 2H), 6.25 - 6.38 (m, 3H), 6.74 (t, 1H), 7.05 (dd, 1H), 7.59 (d, 1H), 8.17 (dd, 1H), 8.38 (d, 1H); Mass spectrum MH⁺ 283.

Reference Example 20

tert-Butyl 2-bromo-4-nitrobenzoate

5 A solution of 2-bromo-4-nitrobenzoic acid (1.66 g) in THF (8 ml) was stirred in a nitrogen atmosphere, carbonyl diimidazole (1.37 g) added and the mixture stirred at 60°C for 2 hours. After cooling to 40°C, *tert*-butanol (0.98 g) and DBU (1.03 g) were added and stirring at 40°C continued overnight. Ether was added to the cooled solution and the solution washed with 10% aqueous HCl, water, aqueous sodium hydrogen carbonate and brine. The ether solution
10 was dried over MgSO₄, filtered and evaporated *in vacuo* to give an oil, which was purified by chromatography using DCM as eluent to give the title compound (1.41 g, 69%); NMR spectrum (DMSO-d₆) 1.56 (s, 9H), 7.87 (d, 1H), 8.27 (dd, 1H), 8.45 (d, 1H).

The procedure described above was repeated using the appropriate benzoic acid. Thus was obtained the compound described below:

15 Reference Example 20.1

tert-Butyl 2-chloro-4-nitrobenzoate

Obtained in 81% yield; NMR spectrum (CDCl₃) 1.63 (s, 9H), 7.84 (d, 1H), 8.13 (dd, 1H), 8.29 (d, 1H); Mass spectrum M⁺ 257.

Reference Example 21

20 tert-Butyl 4-nitro-2-(trimethylsilylethynyl)benzoate

A solution of *tert*-butyl 2-bromo-4-nitrobenzoate (reference example 20) (10.8 g) in triethylamine (85 ml) was stirred under a nitrogen atmosphere, (trimethylsilyl)acetylene (4.27 g) was added followed by *bis*(triphenylphosphine)palladium (II) chloride (0.34 g) and the mixture stirred at 80°C for 3 hours. The mixture was filtered, the residue washed with ethyl
25 acetate and the combined washings and filtrate evaporated *in vacuo*. The resulting oil was dissolved in ethyl acetate and heated briefly at reflux with decolourising charcoal. The charcoal was removed by filtration and the solvent removed *in vacuo* to give a brown oil. This was purified by column chromatography, eluting with *iso*-hexane / ethyl acetate (95/5) to give the title compound (10.70 g, 94%); NMR spectrum (CDCl₃) 0.21 (s, 9H), 1.57 (s, 9H), 7.80 (d,
30 1H), 8.07 (dd, 1H), 8.29 (d, 1H).

Reference Example 22

tert-Butyl 2-ethynyl-4-nitrobenzoate

A solution of *tert*-butyl 4-nitro-2-(trimethylsilylethynyl)benzoate (reference example 21) (12.54 g) in methanol (250 ml) was treated with potassium carbonate (25.0 g) and stirred for 15 minutes. The mixture was poured into water, extracted with ethyl acetate, the extracts dried, filtered and evaporated *in vacuo* to give the crude product. This was purified by chromatography, using *iso*-hexane / ethyl acetate (9/1) to give the title compound (8.77 g, 90%); NMR spectrum (CDCl₃) 1.63 (s, 9H), 3.49 (s, 1H), 7.99 (d, 1H), 8.18 (dd, 1H), 8.40 (d, 1H).

Reference Example 23

tert-Butyl 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoate

To a solution of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) (1.2 g) and *tert*-butyl 4-amino-2-chlorobenzoate (reference example 8.2) (0.98 g) in DMA (60 ml) was added HCl in dioxane (4.0M, 3.0 ml) and the mixture heated at 80°C for 1 hour. The reaction was cooled, diluted with acetone and the resulting precipitate filtered off and dried. The crude product was purified by chromatography, using DCM / methanol / 880 NH₄OH (100/8/1) as eluent, to give the title compound (1.04 g, 52%); NMR spectrum (DMSO-d₆) 1.54 (s, 9H), 1.94 (m, 2H), 2.18 (s, 3H), 2.25 (m, 4H), 2.65 (m, 2H), 4.78 (m, 1H), 7.27 (d, 1H), 7.38 (d, 1H), 7.65 (dd, 1H), 7.78 (m, 2H), 8.34 (d, 1H), 8.64 (s, 1H), 10.36 (s, 1H); Mass spectrum MH⁺ 469.

The procedure described above was repeated using the appropriate aniline and chloroquinazoline. Thus was obtained the compound described below:

Reference Example 23.1

tert-Butyl 2-ethynyl-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoate hydrochloride

Obtained from 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 2) and *tert*-butyl 4-amino-2-ethynylbenzoate (reference example 9) in 57% yield; NMR spectrum (CDCl₃) 1.62 (s, 9H), 2.07 (m, 2H), 2.31 (m, 4H), 2.35 (s, 3H), 2.84 (m, 2H), 3.39 (s, 1H), 4.65 (m, 1H), 6.95 (d, 1H), 7.49 (d, 1H), 7.64 (t, 1H), 7.91 (dd, 1H), 7.96 (d, 1H), 8.08 (d, 1H), 8.71 (s, 1H), 10.29 (1, 1H); Mass spectrum MH⁺ 459.

Reference Example 24

2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid hydrochloride

A solution of *tert*-butyl 2-chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoate (reference example 23) (1.04 g) in 2N HCl (50 ml) was stirred at 50°C for 2 hours and allowed to cool to room temperature. The mixture was diluted with acetone and the resulting precipitate filtered and dried to give the title compound (1.00g, 100%); Mass spectrum MH⁺ 413.

The procedure described above was repeated using the appropriate *tert*-butyl ester. Thus was obtained the compound described below:

Reference Example 24.1

2-Ethynyl-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoic acid

10 hydrochloride

Obtained from *tert*-butyl 2-ethynyl-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)benzoate hydrochloride (reference example 23.1) in 91% yield; Mass spectrum MH⁺ 403.

Reference Example 25

15 1-(2-Chloro-4-nitrobenzoyl)azepane

To a solution of 2-chloro-4-nitrobenzoic acid (4.02 g) and triethylamine (2.20 g) in DCM (150 ml) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (8.36 g). This mixture was stirred at room temperature under an atmosphere of nitrogen for 3 hours. Hexamethyleneimine (2.18 g) was added and this mixture stirred at room temperature for 2 hours. The solvent was removed *in vacuo* and the residue purified by chromatography using 0 - 10% ethyl acetate in DCM as eluent to give the title compound as a colourless oil which crystallised upon standing (5.09 g, 90%); NMR spectrum (CDCl₃) 1.53 - 1.77 (m, 6H), 1.79-1.93 (m, 2H), 3.15 - 3.27 (m, 2H), 3.60 - 3.84 (m, 2H), 7.48 (d, 1H), 8.18 (dd, 1H), 8.30 (d, 1H); Mass spectrum MH⁺ 283.

25 Reference Example 26

1-Benzenesulphonyl-5-nitroindole

5-Nitroindole (2 g) was added to sodium hydride (1.98 g) in THF at 0°C over 15 minutes to give a red solution. Once effervescence had subsided, benzene sulphonyl chloride (1.73 ml) was added dropwise over 5 minutes to give an orange suspension. After 30 minutes a further amount of benzene sulphonyl chloride (0.47 ml) was added to give a yellow suspension. The THF was removed *in vacuo* and water was added to give a yellow solid. This was filtered and dried *in vacuo* to afford the title compound as a yellow solid (4.33g, > 100%);

NMR spectrum (DMSO-d₆) 7.0 (d, 1H), 7.6 (m, 2H), 7.7 (m, 1H), 8.0 (m, 3H), 8.2 (m, 2H), 8.6 (d, 1H).

The procedure described above was repeated using the appropriate alkylating agent. Thus was obtained the compound described below:

5 **Reference Example 26.1**

1-Benzyl-5-nitroindole

Obtained in > 100% yield; NMR spectrum (DMSO-d₆) 5.5 (s, 2H), 6.8 (d, 1H), 7.2 (m, 5H), 7.7 (d, 1H), 7.8 (d, 1H), 8.0 (dd, 1H), 8.6 (d, 1H); Mass spectrum M-H⁺ 251.

Reference Example 27

10 **Pyrimidine-2-methanol**

Sodium borohydride (0.28 g) was added in a single portion to a stirred solution of methyl pyrimidine-2-carboxylate (1 g) in ethanol (50 ml). The reaction was stirred for 4 hours, then water (4 ml) added and the reaction stirred overnight. The resulting solid was filtered and the filtrate concentrated. The residue was purified by chromatography using DCM - 10%
15 methanol as eluent to give the title compound as a colourless liquid (0.28 g, 35%); NMR spectrum (CDCl₃) 3.74 (bs, 1H), 4.83 (d, 1H), 7.22 (dt, 1H), (8.75 (d, 2H); Mass Spectrum MH⁺ 111.

The procedure described above was repeated using the appropriate ester. Thus was obtained the compound described below:

20 **Reference Example 27.1**

3-Hydroxymethylisoxazole

Obtained from 3-ethoxycarbonylisoxazole (**reference example 41**) in 93% yield; NMR spectrum (CDCl₃) 2.50 (bs, 1H), 4.80 (s, 2H), 6.42 (d, 1H), 8.37 (d, 1H).

Reference Example 27.2

25 **5-(Hydroxymethyl)-3-methylisoxazole**

Obtained from 3-methylisoxazole-5-carboxylic acid methyl ester (**reference example 41**) in 73% yield; NMR spectrum (DMSO-d₆) 2.2 (s, 3H), 4.9 (s, 2H), 5.5 (t, 1H), 6.2 (s, 1H).

Reference Example 28

4-Benzoyloxy-3-ethynyl-nitrobenzene

30 Benzyl bromide (0.22 ml) was added to 4-hydroxy-3-iodo-nitrobenzene (0.5 g) and potassium carbonate (0.86 g) in DMF (10 ml) and the reaction stirred overnight. To the resulting suspension was added trimethylsilylacetylene (0.77 ml), copper(I)iodide (7 mg),

bis(triphenylphosphine)-dichloropalladium (26 mg) and triethylamine (2 ml) under nitrogen and the mixture stirred at room temperature for 1 hour. Methanol was added to the reaction, and stirring was continued for a further hour, before concentration *in vacuo*. The residue was purified by chromatography using DCM - 25% *iso*-hexane as eluent to give the title compound as a yellow solid (0.1 g, 21%); NMR spectrum (DMSO-d₆) 4.47 (s, 1H), 5.35 (s, 2H), 7.30 - 7.50 (m, 6H), 8.23 (m, 2H); Mass Spectrum M-H⁺ 252.

Reference Example 29

4-(2-Fluorobenzyloxy)-3-(trimethylsilylethynyl)-nitrobenzene

To a solution of 4-(2-fluorobenzyloxy)-3-iodo-nitrobenzene (0.49 g) (reference example 18.1) in acetonitrile (10 ml) was added trimethylsilylacetylene (0.54 ml), copper(I)iodide (5 mg), bis(triphenylphosphine)-dichloropalladium (18 mg) and triethylamine (10 ml) under nitrogen and the mixture stirred at room temperature for 4 hours. The reaction was concentrated *in vacuo* and the residue purified by chromatography using DCM as eluent to give the title compound as a yellow solid (0.33 g, 73%); Mass Spectrum M-H⁺ 342.

The procedure described above was repeated using the appropriate iodobenzene. Thus were obtained the compounds described below:

Reference Example 29.1

4-(3-Fluorobenzyloxy)-3-(trimethylsilylethynyl)-nitrobenzene

Obtained from 4-(3-fluorobenzyloxy)-3-iodo-nitrobenzene (reference example 18.2); Mass Spectrum M-H⁺ 342.

Reference Example 29.2

4-(2, 6-Difluorobenzyloxy)-3-(trimethylsilylethynyl)-nitrobenzene

Obtained from 4-(2, 6-difluorobenzyloxy)-3-iodo-nitrobenzene (reference example 18.3); Mass Spectrum M-H⁺ 360.

Reference Example 29.3

4-(4-Thiazolylmethoxy)-3-(trimethylsilylethynyl)-nitrobenzene

Obtained from 3-iodo-4-(thiazol-4-ylmethoxy)-nitrobenzene (reference example 18.4) in 60% yield; Mass Spectrum MH⁺ 333.

Reference Example 29.4

4-(5-methylisoxazol-3-ylmethoxy)-3-(trimethylsilylethynyl)-nitrobenzene

Obtained from 3-iodo-4-(5-methylisoxazol-3-ylmethoxy)-nitrobenzene (reference example 18.5) in 86% yield; Mass Spectrum MH⁺ 333.

Reference Example 30**3-Ethynyl 4-(2-fluorobenzyloxy)aniline**

A solution of 4-(2-fluorobenzyloxy)-3-(trimethylsilylethynyl)nitrobenzene (310 mg) (reference example 29) and 10% Pt/C in ethyl acetate / ethanol (9:1, 10 ml) was stirred under an atmosphere of hydrogen for 20 mins. The catalyst was removed by filtration and the solution concentrated *in vacuo* to give a green solid. This was dissolved in methanol (100 ml) and DCM (50 ml), potassium carbonate (0.375 g) added and the solution stirred for 30 mins. The reaction was filtered and concentrated *in vacuo*. The residue was purified by chromatography using DCM as eluent to give the title compound as a yellow oil (0.13 g, 62%, 2 steps); Mass Spectrum MH^+ 283.

The procedure described above was repeated using the appropriate nitrobenzene. Thus were obtained the compounds described below:

Reference Example 30.1**3-Ethynyl 4-(3-fluorobenzyloxy)aniline**

Obtained from 4-(3-fluorobenzyloxy)-3-(trimethylsilylethynyl)nitrobenzene (reference example 29.1); Mass Spectrum MH^+ 283.

Reference Example 30.2**4-(2,6-Difluorobenzyloxy)-3-ethynylaniline**

Obtained from 4-(2, 6-difluorobenzyloxy)-3-(trimethylsilylethynyl)nitrobenzene (reference example 29.2); Mass Spectrum MH^+ 301.

Reference Example 30.3**3-Ethynyl 4-(4-thiazolylmethoxy)aniline**

Obtained from 4-(thiazol-4-ylmethoxy)-3-(trimethylsilylethynyl)nitrobenzene (reference example 29.3); Mass Spectrum MH^+ 231.

Reference Example 30.4**3-Ethynyl 4-(5-methylisoxazol-3-ylmethoxy)aniline**

Obtained from 4-(5-methylisoxazol-3-ylmethoxy)-3-(trimethylsilylethynyl)nitrobenzene (reference example 29.4); Mass Spectrum MH^+ 229.

Reference Example 31**3-Chloromethylisoxazole**

Thionyl chloride (5 ml) was added to 3-hydroxymethylisoxazole (reference example 27.1) (0.5 g), followed by DMF (0.1 ml). The reaction was heated at 100°C for 3 hours and

then concentrated *in vacuo*. The residue was partitioned between DCM and saturated aqueous sodium hydrogen carbonate solution ($T < 5^{\circ}\text{C}$). Combined organic extracts were washed with water, dried and concentrated to give the title compound as a brown oil (0.163 g, 28%); NMR spectrum (CDCl_3) 4.63 (s, 2H), 6.45 (d, 1H), 8.40 (m, 1H).

5 **Reference Example 32**

3-Ethoxycarbonylisoxazole

Triethylamine (2.21 ml) in ether (13 ml) was added dropwise to a solution of carboethoxychloraldoxime (2 g) and vinyl acetate (11.4 g) in ether (27 ml) heated at reflux. The solution was heated for a further hour then cooled and extracted with water. Organic
10 extracts were dried and concentrated and the resulting oil was heated at 180°C for 30 mins. The reaction was partitioned between DCM and water. Combined organic extracts were dried and concentrated to give the title compound as a pale brown oil (1.64 g, 88%); NMR spectrum (CDCl_3) 1.43 (t, 3H), 4.46 (q, 2H), 6.79 (d, 1H), 8.52 (d, 1H).

Reference Example 33

15 **4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-fluoroquinazoline**

3-Chloro-4-(3-fluorobenzyloxy)aniline (reference example 8.1) (1.65 g), 4-chloro-5-fluoroquinazoline (reference example 3) (1.20 g) and di-*iso*-propylethylamine (1.15 ml) in IPA (50 ml) were heated at reflux for 1 hr. The resulting precipitate was isolated by filtration, and washed with IPA and ether to give the title compound as a grey solid (2.02 g, 77%); NMR
20 spectrum ($\text{DMSO}-d_6$) 5.24 (s, 2H), 7.15 (dt, 1H), 7.23 (d, 1H), 7.30 (m, 1H), 7.36 - 7.50 (m, 2H), 7.55 - 7.63 (m, 2H), 7.78 - 7.92 (m, 2H), 8.55 (s, 1H), 9.10 (d, 1H); Mass Spectrum MH^+ 398.

Reference Example 34

4-(2-Aminothiazol-4-ylmethoxy)-3-chloronitrobenzene

25 Sodium hydride (60% dispersion in mineral oil, 200mg) was suspended in DMA (20 ml), and 2-chloro-4-nitrophenol (1.74 g) was added. The mixture was stirred at ambient temperature for 1 hour. Tetra-*n*-butylammonium iodide (20 mg) was added. 2-Amino-4-(chloromethyl)thiazole hydrochloride (185 mg) was added dropwise over 1 hour as a solution in DMA (20 ml). The solution was stirred overnight at ambient temperature. The solvent was
30 evaporated, and the residue partitioned between saturated sodium carbonate solution and DCM. Combined organic extracts were dried and concentrated and the residue purified by chromatography using ethyl acetate as eluent to give the title compound as a yellow solid (85

mg, 30%); NMR spectrum (DMSO-d₆) 5.11 (s, 2H), 6.66 (s, 1H), 6.99 (b s, 2H), 7.49 (d, 1H), 8.19 (dd, 1H), 8.29 (d, 1H); Mass Spectrum MH⁺ 286.

Reference Example 35

2-(2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy)-N-

5 hydroxyacetamide.

Hydroxylamine hydrochloride (121 mg) was dissolved in a mixture of ethanol (4 ml) and water (16 ml). Potassium carbonate (131 mg) was added. 2-[2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetonitrile hydrochloride (reference example 4.5) (200 mg) was added, the mixture briefly sonicated, then heated to 90°C for 10 hour. The colourless solution was cooled to ambient temperature and the resulting solid filtered, washed with water and diethyl ether and dried *in vacuo* to give the title compound as a white crystalline solid (163 mg, 82%); NMR spectrum (DMSO-d₆) 1.83 - 1.96 (m, 2H), 2.08 - 2.18 (m, 2H), 2.17 (s, 3H), 2.22 - 2.32 (m, 2H), 2.56 - 2.66 (m, 2H), 4.48 (s, 2H), 4.78 (m, 1H), 5.58 (s, 2H), 7.21 (d, 1H), 7.27 (d, 1H), 7.31 (d, 1H), 7.49 (dd, 1H), 7.70 (dd, 1H), 8.09 (d, 15 1H), 8.50 (s, 1H), 9.31 (s, 1H), 10.03 (s, 1H); Mass Spectrum MH⁺ 457.

Reference Example 36

Ethyl 2-[2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetate

4-(3-Chloro-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline (reference example 4.2) (384 mg) and potassium carbonate (552mg) were suspended in acetonitrile. The 20 suspension was briefly sonicated, and ethyl bromoacetate (128 µl) was added. The mixture was heated at 80°C for 90 minutes, then cooled to ambient temperature. The solid was removed by filtration, and the filtrate evaporated. The residue was purified by chromatography, using 0 to 1.5% (7:1 methanol / conc. ammonia (aq)) in DCM as eluent to give the title compound as a white crystalline solid (375mg, 80%); NMR spectrum (DMSO-d₆) 25 1.21 (t, 3H), 1.83 - 1.97 (m, 2H), 2.07 - 2.17 (m, 2H), 2.17 (s, 3H), 2.21 - 2.32 (m, 2H), 2.56 - 2.66 (m, 2H), 4.17 (q, 2H), 4.79 (m, 1H), 4.90 (s, 2H), 7.10 (d, 1H), 7.21 (d, 1H), 7.32 (d, 1H), 7.47 (dd, 1H), 7.71 (dd, 1H), 8.11 (d, 1H), 8.50 (s, 1H), 10.04 (s, 1H); Mass Spectrum MH⁺ 471.

Reference Example 37

30 2-[2-Chloro-4-(5-(1-methylpiperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetic acid hydrazide

Ethyl 2-[2-chloro-4-(5-(1-methyl-piperidin-4-yloxy)quinazolin-4-ylamino)phenoxy]acetate (**reference example 36**) (150 mg) was dissolved in ethanol (15 ml). Pyridine (2 drops) and hydrazine monohydrate (150 μ l) were added, and the mixture heated at reflux for 16 hours. The solvent was evaporated and the residue purified by chromatography, using 0 to 3% (7:1 methanol / conc. ammonia (aq)) in DCM as eluent to give the title compound as a white crystalline solid (77 mg, 53%); NMR spectrum (DMSO-d₆) 1.83 - 1.97 (m, 2H), 2.08 - 2.18 (m, 2H), 2.17 (s, 3H), 2.22 - 2.32 (m, 2H), 2.56 - 2.67 (m, 2H), 4.34 (bs, 2H), 4.58 (s, 2H), 4.78 (s, 1H), 7.11 (d, 1H), 7.21 (d, 1H), 7.32 (d, 1H), 7.48 (dd, 1H), 7.71 (dd, 1H), 8.10 (d, 1H), 8.50 (s, 1H), 9.20 (bs, 1H), 10.04 (s, 1H); Mass Spectrum MH⁺ 471.

10 Reference Example 38

4-(3-Methyl-4-hydroxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline

A solution of hydrochloric acid in dioxane (4M, 2 ml) was added to a mixture of 4-chloro-5-(1-methylpiperidin-4-yloxy)quinazoline (**reference example 2**) (2.78 g) and 4-amino-2-methylphenol (1.35 g) in THF (150 ml). The resulting suspension was heated at reflux for 3 hours, then allowed to cool to room temperature. Filtration gave the crude product as a yellow solid, which was partitioned between aqueous sodium hydrogen carbonate solution and DCM. Combined organic extracts were dried and concentrated to give the crude product, which was purified by trituration under cold methanol to give the title compound as a white solid (2.88 g, 79%); NMR spectrum (CDCl₃) 2.0 (m, 2H), 2.2 (m, 2H), 2.3 (s, 3H), 2.3 (s, 3H), 2.4 (m, 2H), 2.8 (m, 2H), 4.7 (m, 1H), 6.7 (d, 1H), 6.9 (d, 1H), 7.3 (dd, 1H), 7.4 (d, 1H), 7.4 (d, 1H), 7.6 (t, 1H), 8.6 (s, 1H), 9.9 (s, 1H); Mass spectrum MH⁺ 365.

Reference Example 39

1,5-Dimethyl-3-hydroxymethylpyrazole

A solution of lithium aluminium hydride (1M solution in THF, 17.8 ml) was added to a solution of 1,5-dimethylpyrazole-3-carboxylic acid (500 mg) in THF (20 ml) at room temperature. The mixture was heated at reflux for 4 hours, then cooled in an ice bath. Ethyl acetate (25 ml) was added dropwise to destroy the excess lithium aluminium hydride (heat evolved), and resultant mixture was then poured into aqueous hydrochloric acid (1M, 50 ml). The product was extracted into ethyl acetate, dried and concentrated *in vacuo* to give the title compound as a pale brown oil (230 mg, 51%); NMR spectrum (DMSO-d₆) 2.2 (s, 3H), 3.6 (s, 3H), 4.3 (s, 2H), 5.9 (s, 1H).

The procedure described above was repeated using the appropriate acid or ester. Thus was obtained the compound described below:

Reference Example 39.1

1-Methyl-3-hydroxymethylpyrazole

5 Obtained from 1-methylpyrazole-3-carboxylic acid in 38% yield; NMR spectrum (DMSO-d₆) 3.8 (s, 3H), 4.4 (s, 2H), 6.1 (d, 1H), 7.5 (d, 1H).

Reference Example 40

3-(2-Chloro-4-nitrophenoxyethyl)-1,5-dimethylpyrazole

 A solution di-*tert*-butylazodicarboxylate (419 mg) in DCM (2 ml) was added to a
10 solution of 2-chloro-4-nitrophenol (211 mg), triphenylphosphine (478 mg), and 1,5-dimethyl-
3-hydroxymethylpyrazole (**reference example 39**) (230mg) in DCM (8 ml), which was stirred
and cooled in an ice bath, under a nitrogen atmosphere. The mixture was then allowed to warm
to room temperature, and stirred for 3 hours. The reaction mixture was concentrated *in vacuo*,
and purified by chromatography, using 0 - 5% methanol in DCM as eluent. This gave the title
15 compound (230 mg, 67%); Mass spectrum MH⁺ 282.

 The procedure described above was repeated using the appropriate phenol and alcohol.
Thus was obtained the compound described below:

Reference Example 40.1

3-(2-Chloro-4-nitrophenoxyethyl)-1-methylpyrazole

20 Obtained by reacting 2-chloro-4-nitrophenol and 1-methyl-3-hydroxymethylpyrazole
(**reference example 39.1**) in 49% yield; NMR spectrum (DMSO-d₆) 3.8 (s, 3H), 5.3 (s, 2H),
6.3 (d, 1H), 7.5 (d, 1H), 7.7 (d, 1H), 8.2 (m, 1H), 8.3 (m, 1H); Mass spectrum MH⁺ 268.

Reference Example 40.2

5-(2-Chloro-4-nitrophenoxyethyl)-3-methylisoxazole

25 Obtained by reacting 2-chloro-4-nitrophenol (251 mg) and 5-(hydroxymethyl)-3-
methylisoxazole (**reference example 27.2**) in 100% yield; NMR spectrum (DMSO-d₆) 2.24 (s,
3H), 5.5 (s, 2H), 6.5 (s, 1H), 7.5 (d, 1H), 8.2 (m, 1H), 8.3 (m, 1H); Mass spectrum MH⁺ 267.

Reference Example 41

3-Methylisoxazole-5-carboxylic acid methyl ester

30 Potassium carbonate (1.19 g) was added portionwise to a cooled solution (0°C) of 3-
methylisoxazole-5-carboxylic acid (500 mg) in DMF (25 ml). After 5 minutes at 0°C, dimethyl
sulphate (0.37 ml) was added dropwise. The reaction mixture was stirred at room temperature

for 1 hour, then concentrated *in vacuo*. Water was added, and the product was extracted into DCM. The combined organics were dried, then concentrated *in vacuo* to give the title compound as a white solid (424 mg, 76%); NMR spectrum (DMSO-d₆) 2.3 (s, 3H), 3.9 (s, 3H), 7.1 (s, 1H).

5 Reference Example 42

4-(3-Chloro-4-hydroxyanilino)-5-(tetrahydropyran-4-yloxy)quinazoline

Phosphorus oxychloride (2.2 ml) was added dropwise to a solution of 3,4-dihydro-5-(tetrahydropyran-4-yloxy)quinazolin-4-one (reference example 1.2) (623 mg) and di-*iso*-propylethylamine (1.36 ml) in anhydrous 1, 2-dichloroethane (25 ml) at 0°C. The mixture was
10 heated at 80°C for 3 hours and then concentrated *in vacuo*. The residue was azeotroped with toluene (20 ml), di-*iso*-propylethylamine (5 ml) was added and the mixture was again concentrated *in vacuo*. The residue was dissolved in IPA (10 ml) and 4-amino-2-chlorophenol (475 mg) was added. The resulting mixture was heated at 80°C for 12 hours and then concentrated *in vacuo*. The residue was purified by chromatography, using 0-5% methanol –
15 DCM as eluent, to give the title compound as a solid (338 mg, 36%); NMR Spectrum (DMSO-d₆) 1.79 - 1.95 (m, 2H), 2.15 - 2.26 (m, 2H), 3.51 - 3.60 (m, 2H), 3.84 - 3.96 (m, 2H), 4.92 - 5.01 (m, 1H), 7.00 (d, 1H), 7.23 (d, 1H), 7.29 - 7.38 (m, 2H), 7.68 (t, 1H), 8.02 (d, 1H), 8.46 (s, 1H), 9.98 (s, 1H); Mass spectrum MH⁺ 372.

The procedure described above was repeated using the appropriate 3,4-
20 dihydroquinazolin-4-one and aniline. Thus were obtained the compounds described below:

Reference Example 42.1

4-(3-Chloro-4-hydroxyanilino)-5-(tetrahydrofuran-3-yloxy)quinazoline

Obtained from 3,4-dihydro-5-(tetrahydrofuran-3-yloxy)quinazolin-4-one (reference example 1.3) and 4-amino-2-chlorophenol in 48% yield; NMR Spectrum (DMSO-d₆) 2.12 -
25 2.21 (m, 1H), 2.17 - 2.41 (m, 1H), 3.78 - 4.00 (m, 3H), 4.18 (d, 1H), 5.40 - 5.44 (m, 1H), 6.98 (d, 1H), 7.18 (d, 1H), 7.36 (d, 1H), 7.40 (dd, 1H), 7.72 (t, 1H), 8.03 (d, 1H), 8.48 (s, 1H), 9.96 (s, 1H); Mass spectrum MH⁺ 358.

Reference Example 42.2

4-(3-Chloro-4-(3-fluorobenzoyloxy)aniline)-5-(1, 4-dioxaspiro[4.5]dec-8-yloxy)quinazoline

30 Obtained from 3,4-dihydro-5-(1, 4-dioxaspiro[4.5]dec-8-yloxy)quinazolin-4-one (reference example 1.4) and 3-chloro-4-(3-fluorobenzoyloxy)aniline (reference example 8.1) in 89% yield; NMR Spectrum (DMSO-d₆) 1.70 (m, 4H), 1.96 (m, 2H), 2.12 (m, 2H), 3.88 (s,

4H), 4.87 (m, 1H), 5.24 (s, 2H), 7.17 (dt, 1H), 7.20 – 7.35 (m, 6H), 7.45 (m, 2H), 7.73 (t, 1H), 8.10 (d, 1H), 8.50 (s, 1H), 10.00 (bs, 1H); Mass Spectrum M^+ 536.

Reference Example 43

3-Fluoro-4-(1-methyl-1H-imidazol-2-ylthio)nitrobenzene

5 To a stirred solution of 2-mercapto-1-methyl-1H-imidazole (1.14 g), in DMF (20 ml), was added sodium hydride (0.44 g) in small portions and the reaction stirred at ambient temperature until effervescence ceased. To this was added a solution of 3,4-difluoronitrobenzene (1.59 g) in DMF (10 ml), and the solution stirred at 80°C for 4 hours. The reaction was poured into water (150 ml) and organic material extracted into ethyl acetate
10 (150 ml). The organic layer was washed successively with water (3 x 150ml), brine (150 ml) and dried. Evaporation of the solvent gave an oil which was purified by chromatography using ethyl acetate and then 10% methanol / ethyl acetate as eluent to give title compound as a solid (1.8 g, 70%); NMR spectrum (DMSO-d₆) 2.5 (s, 3H), 6.7 - 6.9 (t, 1H), 7.2 (t, 1H), 7.6 (s, 1H), 7.95 - 8.05 (dd, 1H), 8.15 - 8.25 (dd, 1H); Mass spectrum MH^+ 254.

15 The procedure described above was repeated using the appropriate thiol. Thus was obtained the compound described below:

Reference Example 43.1

3-Fluoro-4-(1-methyl-1H-1,3,4-triazol-2-ylthio)nitrobenzene

Obtained by reacting 3,4-difluoronitrobenzene with 1-methyl-2-mercaptotriazole in
20 39% yield; NMR spectrum (CDCl₃) 3.7 (s, 3H), 7.3 - 7.4 (t, 1H), 7.9 - 8.0 (m, 1H), 8.36 (s, 1H); Mass spectrum MH^+ 255.

Reference Example 44

3-Chloro-4-(2-pyridylthio)nitrobenzene

To a stirred solution of 2-mercaptopyridine (1.11 g) in acetonitrile (25 ml) was added
25 potassium carbonate (2.76 g), 3-chloro-4-fluoronitrobenzene (1.75 g) and *cis*-dicyclohexano-18-crown-6 (10 mg) and the solution was stirred and heated at reflux for 6 hours. The solution was filtered and evaporated and the residue purified by chromatography using *iso*-hexane / ethyl acetate (2:1) to ethyl acetate as eluent to give a solid which was recrystallised from ethyl acetate / *iso*-hexane to give the title compound as a yellow solid (1.5 g, 58%); NMR spectrum
30 (CDCl₃) 7.2 - 7.3 (m, 2H), 7.4 - 7.48 (d, 1H), 7.5 - 7.55 (d, 1H), 7.65 - 7.75 (dt, 1H), 8.0 - 8.1 (dd, 1H), 8.3 (d, 1H), 8.6 (m, 1H); Mass spectrum MH^+ 267.

The procedure described above was repeated using the appropriate thiol and fluoronitrobenzene. Thus was obtained the compound described below:

Reference Example 44.1

3-Chloro-4-(2-pyrimidinylthio)nitrobenzene

- 5 Obtained by reacting 3-chloro-4-fluoro-nitrobenzene with 2-mercaptopyrimidine in 75% yield; NMR spectrum (CDCl₃) 7.02 - 7.1 (t, 1H), 7.94 - 8.0 (d, 1H), 8.1 - 8.2 (dd, 1H), 8.4 (d, 1H), 8.5 (d, 2H); Mass spectrum MH⁺ 268.

Reference Example 44.2

3-Chloro-4-(1H-imidazol-2-ylthio)nitrobenzene

- 10 Obtained by reacting 3-chloro-4-fluoro-nitrobenzene with 2-mercaptoimidazole in 58% yield; Mass spectrum MH⁺ 255.

Reference Example 44.3

3-Fluoro-4-(1H-imidazol-2-ylthio)nitrobenzene

- 15 Obtained by reacting 3,4-difluoronitrobenzene and 2-mercaptoimidazole in 38% yield; Mass spectrum MH⁺ 240.

Reference Example 44.4

4-(2-Thiazolylthio)nitrobenzene

Obtained by reacting 4-fluoronitrobenzene and 2-mercaptothiazole in 77% yield; NMR spectrum (CDCl₃) 7.5 (d, 1H), 7.5 - 7.6 (d, 2H), 7.9 (d, 1H), 8.1 - 8.2 (d, 2H).

- 20 **Reference Example 44.5**

3-Chloro-4-(2-thiazolylthio)nitrobenzene

Obtained by reacting 3-chloro-4-fluoronitrobenzene and 2-mercaptothiazole in 65% yield; NMR spectrum (CDCl₃) 7.25 (d, 1H), 7.6 (d, 1H), 8.0 (d, 1H), 8.0 - 8.05 (d, 1H), 8.15 (d, 1H); Mass spectrum MH⁺ 273.

- 25 **Reference Example 45**

4-(2-Thiazolylsulphonyl)nitrobenzene

- To a solution of 4-(2-thiazolylthio)nitrobenzene (reference example 44.4) (238 mg) in DCM (50 ml) was added 3-chloroperbenzoic acid (500 mg) and the reaction stirred for 18 hours. The reaction was evaporated and the residue dissolved in ethyl acetate (100 ml) and
30 washed with sodium hydroxide solution (2 x 50ml, 2N), water (2 x 50ml), brine (50 ml) and dried. Concentration of the solvent gave the title compound as a white solid (200 mg, 69%); NMR spectrum (CDCl₃) 7.75 (d, 1H), 8.0 (d, 1H), 8.3 - 8.4 (d, 2H), 8.4 - 8.5 (d, 2H).

Reference Example 46**1, 4-Dioxaspiro[4.5]decan-8-ol**

Sodium borohydride (1.61 g) was added to a solution of 1, 4-dioxaspiro[4.5]decan-8-one (6.65 g) in ethanol (100 ml) and the reaction stirred for 4 hours. Water (10 ml) was added and the reaction stirred for 16 hours. The resulting solid was filtered and the filtrate concentrated *in vacuo*. The residue was purified by chromatography using *iso*-hexane - 30% ethyl acetate as eluent to give the title compound as a colourless oil (5.49 g, 82%); NMR Spectrum (CDCl₃) 1.50 - 1.95 (m, 8H), 3.80 (m, 1H), 3.95 (s, 4H); Mass Spectrum MH⁺ 159.

Reference Example 47**4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(1-oxo-cyclohex-4-yloxy)quinazoline acetate**

A solution of 4-(3-chloro-4-(3-fluorobenzyloxy)anilino)-5-(1, 4-dioxaspiro[4.5]dec-8-yloxy)quinazoline (reference example 42.2) (2.25 g) in acetic acid (40 ml) and water (10 ml) was heated at 90°C for 18 hours, then concentrated *in vacuo*. The residue was triturated with ether to give the title compound as a white solid (1.98 g, 85%); Mass Spectrum M⁺ 492.

Reference Example 48**Methanesulphonic acid 2-methyloxazol-4-ylmethyl ester**


To a solution of (2-methyloxazol-4-yl)methanol (obtained as described in *J. Chem. Soc., Perkin Trans. 1*, 2000, 2415 – 2428) (112 mg) in DCM (10 ml) was added triethylamine (275 µl). The solution was cooled to 0°C and methane sulphonyl chloride (85 µl) was added dropwise. The reaction mixture was allowed to warm up to ambient temperature and left stirring for nine hours. The mixture was concentrated *in vacuo* to give the title compound which was used without further purification (103mg; 40%); Mass Spectrum M-H⁺ 192.

Reference Example 49**2-Chloro-5-fluoro-4-nitrophenol**

2-Chloro-5-fluorophenol (1.46 g) was stirred in ethanol and ferric nitrate monohydrate (4.04 g) added slowly. The solution turned blue and was heated at 50°C for 5 hours. The solution was cooled and evaporated to ¼ volume and 1N hydrochloric acid (100 ml) and water (100 ml) added and the solution extracted with ethyl acetate, washed with water, brine, dried and evaporated to give an oil which was purified by chromatography, using *iso*-hexane / ethyl acetate (3:1) as eluent to give the title compound as a solid; NMR spectrum (CDCl₃) 6.2 - 6.4 (bs, 1H), 6.9 - 6.95 (d, 1H), 8.2 (d, 1H).

CLAIMS

1. A quinazoline derivative of the Formula I



5

wherein:

R^1 is hydrogen or (1-6C)alkyl;

Z is a direct bond or is selected from O, S and $N(R^2)$, wherein R^2 is hydrogen or (1-6C)alkyl;


Q^1 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q^1 -Z- group are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, $N(R^3)$, CO, CH(OR³), CON(R³), $N(R^3)$ CO, SO₂N(R³), $N(R^3)$ SO₂, CH=CH and C≡C wherein R^3 is hydrogen or (1-6C)alkyl,

and wherein any CH₂ or CH₃ group within the Q^1 -Z- group optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,


and wherein any heterocyclyl, (3-7C)cycloalkyl or (3-7C)cycloalkenyl group within the Q^1 -Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different,

selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, 5 N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, amino(2-6C)alkanoyl, N-(1-6C)alkylamino(2-6C)alkanoyl, N,N-di-[(1-6C)alkyl]amino(2-6C)alkanoyl, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylaminol, N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, from a group of the formula :



wherein X^1 is a direct bond or is selected from O and N(R^5), wherein R^5 is hydrogen or (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,

15 N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, and from a group of the formula:




wherein X^5 is a direct bond or is selected from O, CO and N(R^{10}), wherein R^{10} is hydrogen or 20 (1-6C)alkyl, and Q^6 is heterocyclyl, heterocyclyl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl or (3-7C)cycloalkenyl-(1-6C)alkyl, and wherein any heterocyclyl, (3-7C)cycloalkyl or (3-7C)cycloalkenyl group in Q^6 optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, hydroxy, cyano, formyl, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, amino, 25 (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (1-6C)alkoxycarbonyl, (1-6C)alkylsulphonyl, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,


30 N-(1-6C)alkylcarbamoyl-(1-6C)alkyl and N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,

and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo or thioxo substituents; and


Q^2 is selected from a group of formula Ia, Ib, Ic, Id and Ie




Ia




Ib



Ic



Id



Ie

5 wherein G^1 , G^2 , G^3 , G^4 and G^5 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

G^6 and G^7 are each, independently, selected from hydrogen and halogeno,


10 X^2 is a direct bond or is selected from O, S, SO, SO₂, N(R⁶), CH(OR⁶), CON(R⁶), N(R⁶)CO, SO₂N(R⁶), N(R⁶)SO₂, OC(R⁶)₂, C(R⁶)₂O, SC(R⁶)₂, C(R⁶)₂S, CO, C(R⁶)₂N(R⁶) and N(R⁶)C(R⁶)₂ wherein each R⁶ is, independently, hydrogen or (1-6C)alkyl, and Q^3 is aryl, or heteroaryl,

or X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a
15 nitrogen atom,

X^3 is a direct bond or is selected from SO₂, CO, SO₂N(R⁷) and C(R⁷)₂, wherein each R⁷ is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,

- 193 -

- and any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and
- 10 N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula:



- wherein X^4 is a direct bond or is selected from O and N(R^9), wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl, or from a group of the formula
- 15



- 20 wherein Q^5 is selected from aryl, heteroaryl or heterocyclyl which is optionally substituted by 1 or 2 substituents selected from halogeno, hydroxy, (1-6C)alkyl, (1-6C)alkoxy, amino, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

- and wherein any CH_2 or CH_3 group within a substituent on any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents,
- 25 and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or thioxo substituents, or a pharmaceutically acceptable salt thereof.

- 30 2. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

R^1 is hydrogen or (1-6C)alkyl;

Z is a direct bond or is selected from O, S and N(R²), wherein R² is hydrogen or (1-6C)alkyl;

Q¹ is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

5 and wherein adjacent carbon atoms in any (2-6C)alkylene chain within the Q¹-Z- group are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, N(R³), CO, CH(OR³), CON(R³), N(R³)CO, SO₂N(R³), N(R³)SO₂, CH=CH and C≡C wherein R³ is hydrogen or (1-6C)alkyl,

and wherein any CH₂ or CH₃ group within the Q¹-Z- group optionally bears on each
 10 said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphanyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
 15 N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl,
 20 cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphanyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,
 25 N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :




wherein X¹ is a direct bond or is selected from O and N(R⁵), wherein R⁵ is hydrogen or (1-6C)alkyl, and R⁴ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl,
 30 cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,

- 195 -


N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl,

and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo or thioxo substituents;


5 Q^2 is selected from a group of formula Ia, Ib, Ic, Id and Ie




Ia




Ib



Ic



Id



Ie

wherein G^1 , G^2 , G^3 , G^4 and G^5 are each, independently, selected from hydrogen, halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

X^2 is a direct bond or is selected from O, S, $N(R^6)$, $CH(OR^6)$, $CON(R^6)$, $OC(R^6)_2$, $C(R^6)_2O$, $SC(R^6)_2$, $C(R^6)_2S$, CO, $C(R^6)_2N(R^6)$ and $N(R^6)C(R^6)_2$ wherein each R^6 is, independently, hydrogen or (1-6C)alkyl, and Q^3 is aryl, or heteroaryl,

or X^2 is CO and Q^3 is a nitrogen containing heterocyclyl group linked to X^2 by a nitrogen atom,

X^3 is a direct link or is selected from SO_2 , CO and $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or (1-6C)alkyl, and Q^4 is aryl or heteroaryl,

and any aryl, heteroaryl or heterocyclyl group in the group $-X^2-Q^3$ and $-X^3-Q^4$ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from
 5 halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and any heterocyclyl group represented by Q^3 or Q^4 optionally bears 1 or 2 oxo or
 10 thioxo substituents,
 or a pharmaceutically acceptable salt thereof.

3. A quinazoline derivative according to claim 1 or claim 2, or a pharmaceutically acceptable salt thereof, wherein each of R^1 and Q^2 is as defined in claim 1 or claim 2, and

15 Z is O; and

Q^1 is (3-7C)cycloalkyl or heterocyclyl,

and wherein any CH_2 or CH_3 group within the Q^1-Z group optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio,
 20 (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and
 25 N-(1-6C)alkyl-(1-6C)alkanesulphonylamino,

and wherein any heterocyclyl group within the Q^1-Z group optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl,
 30 (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,

- 197 -

N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and
N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein X^1 is a direct bond or is selected from O and N(R^5), wherein R^5 is hydrogen or
 5 (1-6C)alkyl, and R^4 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl,
 cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl,
 di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl,
N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl,
 (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl,
 10 and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo
 or thioxo substituents.

4. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt
 thereof, wherein each of R^1 and Q^2 is as defined in claim 1, and

15 Z is O; and

Q^1 is selected from a 4, 5 or 6 membered heterocyclyl ring containing at least 1
 nitrogen atom, said ring being linked to Z by a carbon atom, and wherein any NH group within
 a heterocyclyl group in Q^1 optionally bears a substituent selected from formyl, cyano,
 (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (2-4C)alkanoyl, carbamoyl,
 20 N-(1-4C)alkylcarbamoyl and N,N-di-(1-4C)alkylcarbamoyl, or from a group of the formula:



wherein X^1 is a direct bond, and R^4 is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl,
 (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-6C)alkyl, amino-(1-4C)alkyl,
 (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl,
 25 N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl or
 (2-4C)alkanoyl-(1-4C)alkyl,
 and wherein any heterocyclyl group within the Q^1 -Z- group optionally bears 1 or 2 oxo
 substituents.

30 5. A quinazoline derivative according to claim 1 or claim 2, or a pharmaceutically
 acceptable salt thereof, wherein each of R^1 and Q^2 is as defined in claim 1 or claim 2, and

the Q¹-Z- group is selected from cyclopentyloxy, tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, tetrahydrothiopyran-4-yloxy, 1,1-dioxotetrahydrothiopyran-4-yloxy, 1-oxotetrahydrothiopyran-4-yloxy, tetrahydrothien-3-yloxy, 1,1-dioxodotetrahydrothien-3-yloxy, 1-oxotetrahydrothien-3-yloxy, pyrrolidin-3-yloxy, 5 pyrrolidin-2-yloxy, piperidin-3-yloxy, piperidin-4-yloxy, homopiperidin-3-yloxy, homopiperidin-4-yloxy and azetidin-3-yloxy,

and wherein the azetidiny, pyrrolidinyl, piperidinyl or homopiperidinyl group within the Q¹-Z- group is optionally N-substituted by a substituent selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, allyl, 2-propynyl, acetyl, propionyl, 10 methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, methylsulphonyl, ethylsulphonyl, 2-methoxyethyl, carbamoylmethyl, N-methylcarbamoylmethyl, N,N-di-methylcarbamoylmethyl, 2-carbamoylethyl, 2-(N-methylcarbamoyl)ethyl, 2-(N,N-di-methylcarbamoyl)ethyl, acetylmethyl, 2-acetylethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl,

15 and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 or 2 oxo substituents.

6. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R¹ and Q² is as defined in claim 1, and

20 Z is O; and

Q¹ is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl, and wherein any NH group within a pyrrolidinyl or piperidinyl group in Q¹ optionally bears a substituent selected from methyl, allyl, acetyl, carbamoyl, methoxymethyl, 2-methoxyethyl, carbamoylmethyl, N-methylcarbamoylmethyl, N,N-di-methylcarbamoylmethyl, 25 2-carbamoylethyl, 2-(N-methylcarbamoyl)ethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, 2-acetylethyl, cyanomethyl, 2-cyanoethyl, fluoromethyl, chloromethyl, 2-fluoroethyl and 2-chloroethyl, and wherein any heterocyclyl group within the Q¹-Z- group optionally bears 1 oxo substituent.

30 7. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R¹, Q¹ and Z is as defined in claim 1, and


Q^2 is a group of the formula Ia as defined in claim 1 wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from hydrogen, halogeno, (1-6C)alkyl, (1-6C)alkoxy, (2-8C)alkenyl and (2-8C)alkynyl,

5 X^2 is selected from S and $OC(R^6)_2$, wherein R^6 is, independently, hydrogen or methyl, and

Q^3 is selected from phenyl, 2- or 3-furyl, 2- or 3-thienyl, 2-,4- or 5-(1,3-oxazolyl), 3-,4- or 5-isoxazolyl, 2-,4- or 5-1H-imidazolyl, 2-,4- or 5-thiazolyl, 1H-1,2,4-triazol-3-yl, 1H-1,2,4-triazol-5-yl, 1H-1,3,4-triazol-2-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-oxadiazol-5-yl, 3-, 4- or 5-1H-pyrazolyl, 2- 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, 1,2-benzisoxazol-3-yl, 1,3-benzodioxol-4-yl, 1,3-benzodioxol-5-yl, 2-imidazo[1,2-a]pyridyl, 3-benzo[d]isoxazolyl and 8-quinoliny,

and wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different,
 15 selected from halogeno, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, sulphamoyl, mercapto, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, 20 (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or 25 (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, carboxy-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, carbamoyl-(1-6C)alkyl, N-(1-6C)alkylcarbamoyl-(1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl-(1-6C)alkyl, (2-6C)alkanoyl-(1-6C)alkyl or (1-6C)alkoxycarbonyl-(1-6C)alkyl.

30

8. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R^1 , Q^1 and Z is as defined in claim 1, and

Q^2 is a group of the formula Ia as defined in claim 1 wherein G^1 , G^6 and G^7 are hydrogen,


G^2 is selected from hydrogen, fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

5 X^2 is $OC(R^6)_2$ wherein R^6 is, independently, hydrogen or methyl, and

Q^3 is phenyl, and

wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl, N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and

15 N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula:



wherein X^4 is a direct bond or is selected from O and $N(R^9)$, wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl.

20

9. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R^1 , Q^1 and Z is as defined in claim 1, and

25

Q^2 is a group of the formula Ia as defined in claim 1 wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from fluoro, chloro, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl,

X^2 is $OC(R^6)_2$ wherein R^6 is hydrogen, and

30 Q^3 is phenyl, and wherein Q^3 is optionally substituted by 1 or 2 substituents selected from fluoro and cyano.

- 201 -

10. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R^1 , Q^1 and Z is as defined in claim 1, and

Q^2 is a group of the formula Ia as defined in claim 1 wherein G^1 , G^6 and G^7 are hydrogen;

5 G^2 is selected from hydrogen, halogeno, (1-4C)alkyl, (1-4C)alkoxy and (2-4C)alkynyl, X^2 is $OC(R^6)_2$ wherein R^6 is, independently, hydrogen or methyl, and

Q^3 is selected from 2-thienyl, 3-thienyl, 1,3-oxazol-4-yl, 3-isoxazolyl, 4-isoxazolyl, 2-1H-imidazolyl, 5-1H-imidazolyl, 2-thiazolyl, 4-thiazolyl, 3-1H-pyrazolyl, 1H-1,2,4-triazol-3-yl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 2-pyridyl, 3-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 2-pyrazine and 1H-1,3,4-triazolyl-2-yl, and wherein Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, difluoromethyl, cyano, nitro, hydroxy, amino, carboxy, formyl, carbamoyl, mercapto, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-[(1-4C)alkyl]amino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, (2-4C)alkanoyl, (2-4C)alkanoyloxy, (2-4C)alkanoylamino, N-(1-4C)alkyl-(2-4C)alkanoylamino, N-(1-4C)alkylsulphamoyl; N,N-di-[(1-4C)alkyl]sulphamoyl, (1-4C)alkanesulphonylamino and N-(1-4C)alkyl-(1-4C)alkanesulphonylamino, or from a group of the formula :

20 $-X^4-R^8$

wherein X^4 is a direct bond or is selected from O and N(R^9), wherein R^9 is hydrogen or methyl, and R^8 is fluoro-(1-4C)alkyl, chloro-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, carboxy-(1-4C)alkyl, amino-(1-4C)alkyl, (1-4C)alkylamino-(1-4C)alkyl, di-[(1-4C)alkyl]amino-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl, N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl, (2-4C)alkanoyl-(1-4C)alkyl or (1-4C)alkoxycarbonyl-(1-4C)alkyl.

11. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R^1 , Q^1 and Z is as defined in claim 1, and

30 Q^2 is a group of the formula Ia as hereinbefore defined wherein G^1 , G^6 and G^7 are hydrogen,

G^2 is selected from fluoro, chloro, methyl, ethyl, methoxy and ethynyl,

X^2 is OCH_2 , and

Q^3 is selected from 3-isoxazolyl, 1,3-oxazol-4-yl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 2-pyridyl, 3-pyridyl and 2-pyrazinyl, and Q^3 optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, cyano, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methylamino)ethyl, di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-(ethyl)carbamoylmethyl 2-(N-(methyl)carbamoyl)ethyl, N,N-di-methylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl.

15

12. A quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein each of R^1 , Q^1 and Z is as defined in claim 1, and

Q^2 is a group of the formula Ib as defined in claim 1 wherein G^1 , G^3 , G^4 , G^6 and G^7 are hydrogen,

20


X^3 is $C(R^7)_2$, wherein each R^7 is, independently, hydrogen or methyl, and

Q^4 is selected from phenyl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 3-isoxazolyl, 2-pyridyl, 3-pyridyl and 4-thiazolyl and wherein Q^4 optionally bears 1 or 2 substituents selected from fluoro, chloro, cyano, carboxy, amino, hydroxy, methyl, ethyl, methoxy, ethoxy, ethynyl, acetyl, formyl, mercapto, methoxycarbonyl, ethoxycarbonyl, carbamoyl, N-methylcarbamoyl, N-ethylcarbamoyl, N,N-di-methylcarbamoyl, N,N-di-ethylcarbamoyl, methylsulphonyl, hydroxymethyl, 2-hydroxyethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, 2-cyanoethyl, carboxymethyl, 2-carboxyethyl, aminomethyl, 2-aminoethyl, methyaminomethyl, 2-(methyaminomethyl), di-methylaminomethyl, 2-(di-methylamino)ethyl, 2-(di-ethylamino)ethyl, carbamoylmethyl, 2-carbamoylethyl, N-(methyl)carbamoylmethyl, N-ethylcarbamoylmethyl 2-(N-methylcarbamoyl)ethyl, N,N-dimethylcarbamoylmethyl, N,N-diethylcarbamoylmethyl, 2-(N,N-dimethylcarbamoyl)ethyl, acetylmethyl, methoxycarbonylmethyl and 2-methoxycarbonylethyl.

30

13. A quinazoline derivative of the Formula I according to claim 1 or claim 2 wherein:

- the Q^1 -Z- group is selected from cyclopentyloxy, tetrahydrofuran-3-yloxy,
 5 1-methylpyrrolidin-3-yloxy, pyrrolidin-3-yloxy, tetrahydropyran-4-yloxy, piperidin-4-yloxy, 1-methylpiperidin-4-yloxy, 1-ethylpiperidin-4-yloxy, 1-(2-methoxyethyl)piperidin-4-yloxy, 1-acetylpiperidin-4-yloxy, 1-acetylmethylpiperidin-4-yloxy, 1-allylpiperidin-4-yloxy, 1-(2-propynyl)piperidin-4-yloxy, 1-methoxycarbonylmethylpiperidin-4-yloxy, 1-carbamoylmethylpiperidin-4-yloxy and 1-methanesulphonylpiperidin-4-yloxy;
 10 R^1 is hydrogen; and
 Q^2 is a group of the formula Ia



Ia

- 15 wherein G^1 and G^2 are each independently selected from hydrogen, fluoro and chloro, X^2 is selected from O, S, OCH_2 , NH, $N(CH_3)$, CO and $NHCH_2$, and Q^3 is selected from phenyl, 2-thienyl, 2-1H-imidazolyl, 3-isoxazolyl, 4-thiazolyl, 3-(1,2,5-thiadiazolyl), 2-pyridyl, 3-pyridyl, 4-pyridyl and 8-quinoliny, which is optionally substituted
 20 by 1 or 2 substituents selected from fluoro, chloro, methyl, methoxy, nitro and cyano; or a pharmaceutically acceptable salt thereof.

14. A quinazoline derivative of the Formula I according to claim 1 wherein:
 Z is O;
 25 Q^1 is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl, and wherein any NH group within a heterocyclyl group in Q^1 optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl,

N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears an oxo substituent;

R¹ is hydrogen; and

Q² is a group of the formula Ia as defined in claim 1 wherein G¹, G⁶ and G⁷ are hydrogen,

G² is chloro,

X² is OCH₂, and

Q³ is selected from phenyl, 2-fluorophenyl, 3-fluorophenyl, 2-cyanophenyl, 3-cyanophenyl, 2,5-difluorophenyl and 2,6-difluorophenyl;

or a pharmaceutically acceptable salt thereof.

15 15. A quinazoline derivative of the Formula I according to claim 1 wherein:

Z is O;

Q¹ is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,

and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl,

N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl,

and wherein any heterocyclyl group within the Q¹-Z- group optionally bears an oxo substituent;

25 R¹ is hydrogen; and

Q² is a group of the formula Ia as defined in claim 1 wherein G¹, G⁶ and G⁷ are hydrogen,

G² is chloro,

X² is OCH₂, and

30 Q³ is selected from 2-pyridyl, 3-pyridyl, 2-pyrazinyl, 4-thiazolyl, 1,2,5-thiadiazol-3-yl, 1,2,3-thiadiazol-4-yl, 5-methylisoxazol-3-yl, 1-methyl-1H-imidazol-5-yl and 1,3-oxazol-4-yl; or a pharmaceutically acceptable salt thereof.

16. A quinazoline derivative of the Formula I according to claim 1 wherein:
Z is O;
Q¹ is selected from pyrrolidin-3-yl, piperidin-3-yl and piperidin-4-yl,
5 and wherein any NH group within a heterocyclyl group in Q¹ optionally bears a substituent selected from methyl, acetyl, allyl, carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, fluoromethyl, chloromethyl, methoxymethyl, 2-methoxyethyl, cyanomethyl, acetylmethyl, carbamoylmethyl, N-methylcarbamoylmethyl and N,N-dimethylcarbamoylmethyl,
10 and wherein any heterocyclyl group within the Q¹-Z- group optionally bears an oxo substituent;
R¹ is hydrogen; and
Q² is a group of the formula Ib as defined in claim 1 wherein G¹, G³, G⁴, G⁶ and G⁷ are hydrogen,
15 X³ is CH₂, and
Q⁴ is selected from phenyl, 2-fluorophenyl, 2-chlorophenyl, 2-methoxyphenyl, 2-cyanophenyl, 3-fluorophenyl, 2,5-dimethylphenyl, 2,6-difluorophenyl, 2-pyridyl, 3-pyridyl and 4-thiazolyl;
or a pharmaceutically acceptable salt thereof.
20
17. A quinazoline derivative according to claim 1 selected from:
4-(3-Chloro-4-phenoxyanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
4-(3-Chloro-4-((3-fluorobenzyloxy)anilino))-5-(1-methylpiperidin-4-yloxy)quinazoline;
4-(1-(3-Fluorobenzyl)indazol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
25 4-(3-Chloro-4-((decahydroquinolin-1-yl)carbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
4-(3-Chloro-4-((decahydroisoquinolin-2-yl)carbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
4-(3-Chloro-4-((3-methylpiperidin-1-yl)carbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
30 yloxy)quinazoline;
4-(3-Ethynyl-4-((decahydroquinolin-1-yl)carbonyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(3-Chloro-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(3-Chloro-4-(2, 6-difluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(3-Chloro-4-(2-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

5 4-(3-Chloro-4-(2-cyanobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and

4-(3-Chloro-4-(thiazol-4-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

or a pharmaceutically acceptable acid addition salt thereof.

18. A quinazoline derivative according to claim 1 selected from:

10 4-(1-Benzylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(1-Benzenesulphonylindol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(1-(2-Cyanobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(1-(2-Methoxybenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(1-(2-Chlorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

15 4-(1-(2, 5-Dimethylbenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(1-(2-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and

4-(1-(3-Fluorobenzyl)indol-5-ylamino)-5-(1-methylpiperidin-4-yloxy)quinazoline

or a pharmaceutically acceptable acid addition salt thereof.

20 19. A quinazoline derivative according to claim 1 selected from:

4-(3-Chloro-4-(3-fluorophenoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(3-Chloro-4-(2-thienoyl)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

4-(3-Chloro-4-((1-methyl-1*H*-imidazol-2-yl)thio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and

25 4-(3-Chloro-4-(1-cyanomethyl-1*H*-imidazol-2-ylthio)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

or a pharmaceutically acceptable acid addition salt thereof.

20. A quinazoline derivative according to claim 1 selected from:


30 4-(3-Chloro-4-(3-fluorobenzyloxy)anilino)-5-(tetrahydropyran-4-yloxy)quinazoline;

4-(4-Benzyl-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;

- 4-(3-Methyl-4-(5-methylisoxazol-3-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(4-(3-Fluorobenzyloxy)-3-methylanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Ethynyl 4-(3-fluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 5 4-(3-Ethynyl 4-(2, 6-difluorobenzyloxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(4-(2-Aminothiazol-4-ylmethoxy)-3-chloroanilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 4-(3-Chloro-4-(pyrazin-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline; and
 4-(3-Chloro-4-(pyridin-2-ylmethoxy)anilino)-5-(1-methylpiperidin-4-yloxy)quinazoline;
 10 or a pharmaceutically acceptable acid addition salt thereof.

21. A process for the preparation of a quinazoline derivative according to claim 1, or a pharmaceutically acceptable salt thereof, which comprises:

- (a) the reaction of a quinazoline of the Formula II




II

- 15 wherein L^1 is a displaceable group and Q^1 and Z have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the Formula




- wherein Q^2 and R^1 have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means; or

- (b) for the production of those compounds of the Formula I wherein Z is an oxygen atom, the coupling of an alcohol of the Formula



- 25 wherein Q^1 has any of the meanings defined in claim 1 except that any functional group is protected if necessary with a quinazoline of the Formula VI




VI

wherein R^1 and Q^2 have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means; or

- 5 (c) for the production of those compounds of the Formula I wherein Z is an oxygen atom, the reaction of an alcohol of the Formula



wherein Q^1 has any of the meanings defined in claim 1 except that any functional group is protected if necessary with a quinazoline of the Formula VIII



VIII


10

wherein R^1 and Q^2 have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means; or

- (d) for the production of those compounds of the Formula I wherein Q^1 or Q^2 contains a
 15 primary or secondary amino group, the cleavage of the corresponding compound of Formula I wherein Q^1 or Q^2 contains a protected primary or secondary amino group; or
 (e) for the production of those compounds of the Formula I wherein Q^1 or Q^2 contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, the alkylation of a quinazoline derivative of the formula I wherein Q^1
 20 or Q^2 contains a hydroxy group or a primary or secondary amino group as appropriate; or
 (f) for the production of those compounds of the Formula I wherein Q^1 contains an amino-hydroxy-disubstituted (1-6C)alkoxy group, the reaction of a compound of the Formula I

wherein Q^1 contains an epoxy-substituted (1-6C)alkoxy group with a heterocycle or an appropriate amine; or

(g) the reaction of a quinazoline of the Formula X



5 wherein L^1 is a displaceable group and R^1 and Q^2 have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the Formula



wherein Q^1 and Z have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by:

10 conventional means; or

(h) for the production of those compounds of the Formula I wherein Q^1 contains an amino-substituted (1-6C)alkoxy group, the reaction of a compound of the Formula I wherein Q^1 contains a halogeno-substituted (1-6C)alkoxy group with an appropriate amine or a heterocycle containing a ring NH group; or

15 (i) for the production of those compounds of the Formula I wherein a heterocyclyl group in Q^1 or Q^3 contains an S- or N-oxide the oxidation of a ring N or S atom in a compound of the formula (I); or


(j) for the production of those compounds of the Formula I wherein:

(i) the group X^2-Q^3 is $CON(R^6)Q^3$ wherein R^6 and Q^3 are as defined in claim 1, or

20 (ii) the group X^2-Q^3 is COQ^3 and Q^3 is a nitrogen linked heterocyclyl group,

the coupling of the carboxy substituted quinazoline of the formula XI:

- 210 -



XI

or a reactive derivative thereof, with an amine of the formula $\text{NH}(\text{R}^6)\text{Q}^3$ or Q^3H as appropriate, wherein R^1 , R^6 , Q^1 , Q^3 , Z , G^1 , G^2 , G^6 and G^7 are as hereinbefore defined except that any functional group is protected if necessary, whereafter any protecting group that is present is

5 removed by conventional means; or

(k) for the production of those compounds wherein Q^2 is a group of the Formula Ia wherein:

(i) X^2Q^3 is OCH_2Q^3 and Q^3 is aryl or heteroaryl, or

(ii) X^2Q^3 is OQ^3 and Q^3 is heteroaryl,

10 the reaction, optionally in the presence of a base, of a compound of formula I as defined in claim 1 wherein Q^2 is a group of the formula Ia as defined in claim 1 except that the group X^2Q^3 in formula Ia is OH , with a compound of the formula $\text{L}^1\text{CH}_2\text{Q}^3$ or L^1Q^3 ,

wherein L^1 is a displaceable group, and Q^3 is as defined in claim 1 except any functional group is protected if necessary, and whereafter any protecting group that is present is removed by


15 conventional means; or

(l) for the production of those compounds of the formula I wherein Q^1 contains an (2-6C)alkanoylamino or substituted (2-6C)alkanoylamino group, the acylation of a quinazoline derivative of the formula I wherein Q^1 contains an amino group; or

(m) for the production of those compounds of the Formula I wherein Q^2 is an indole or


20 indazole derivative of the formula Ib, Ic, Id or Ie as defined in claim 1, the reaction of a compound of the formula $\text{Q}^4\text{X}^3\text{L}^1$, wherein L^1 is a displaceable group and Q^4 and X^3 are as defined in claim 1, except that any functional group is protected if necessary, with a compound of the formula XII,

- 211 -




XII


wherein Q^1 and R^1 are as defined in claim 1, except that any functional group is protected if necessary, and Q^{2a} is selected from a compound of the formula Ib', Ic', Id' and Ie'




Ib'



Ic'




Id'



Ie'

- 5 wherein G^1 , G^4 , G^5 , G^6 and G^7 have any of the meanings defined in claim 1, except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means; or
- (n) for the production of those compounds of the Formula I wherein Q^2 is a group of the formula Ia wherein Q^3 is 1,2,4-oxadiazol-4-yl, or 5-alkyl substituted 1,2,4-oxadiazol-4-yl, the
- 10 ring closure of the hydroxyamidine of the formula XIII

- 212 -




XIII

wherein Q^1 , Z , R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined in claim 1, except that any functional group is protected if necessary, with an orthoester of the formula $RCO(OCH_3)$ or carboxylic acid of the formula $RCOOH$, wherein R is hydrogen or (1-6C)alkyl,

5 whereafter any protecting group that is present is removed by conventional means; or

(o) for the production of those compounds of the Formula I wherein Q^2 is a group of the formula Ia wherein Q^3 is 5-amino-1,3,4-oxadiazol-5-yl, the cyclisation of the compound of the formula XVI




XVI

10

wherein Q^1 , Z , R^1 , X^3 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore, except that any functional group is protected if necessary, with a cyanogen halide, whereafter any protecting group that is present is removed by conventional means; or

(p) for the production of those compounds of the Formula I wherein Q^2 is a group of the
15 formula Ia wherein Q^3 is aryl or heteroaryl and X^3 is S, the coupling of the compound of formula XVIII

- 213 -



XVIII

wherein Q^1 , Z, R^1 , G^1 , G^2 , G^6 and G^7 have any of the meanings defined hereinbefore; except that any functional group is protected if necessary, with a compound of the formula Q^3SH , wherein Q^3 is aryl or heteroaryl as defined in claim 1, except that any functional group is
 5 protected if necessary, in the presence of cuprous bromide, and a base, whereafter any protecting group that is present is removed by conventional means;

(q) for the production of those compounds of the Formula I wherein Q^1 or Q^2 contains an (1-6C)alkylamino, substituted (1-6C)alkylamino group or a nitrogen linked heterocyclyl group, the reductive amination of an aldehyde or ketone group in a compound of formula 1, with a
 10 (1-6C)alkylamino, substituted (1-6C)alkylamino group or a heterocyclyl group containing an NH group in the presence of a suitable reducing agent; or

(r) the conversion of one compound of the Formula I into another compound of the Formula I;

and when a pharmaceutically acceptable salt of a quinazoline derivative of formula I is
 15 required it may be obtained using a conventional procedure.

22. A pharmaceutical composition which comprises a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable thereof, as defined in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.

20

23. A quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 for use in a method of treatment of the human or animal body by therapy.

- 214 -

24. The use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined in claim 1 in the manufacture of a medicament in the prevention or treatment of tumours which are sensitive to the inhibition of one or more erbB receptor tyrosine kinases.

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/GB 02/04931

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07D239/94 C07D401/14 C07D401/12 C07D409/12 C07D403/12
 C07D417/14 C07D413/14 C07D409/14 A61K31/505 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 09294 A (WELLCOME FOUND ;HUDSON ALAN THOMAS (GB); VILE SADIE (GB); BARRACLO) 28 March 1996 (1996-03-28) cited in the application page 4 -page 8; claims 1,26	1-24
X	WO 96 15118 A (THOMAS ANDREW PETER ;ZENECA LTD (GB); BROWN DEARG SUTHERLAND (GB);) 23 May 1996 (1996-05-23) page 3, line 23 - line 25; claims 1,12	1-24
Y	WO 97 30034 A (ZENECA LTD) 21 August 1997 (1997-08-21) page 4, line 24 -page 9, line 2; claims 1,15	1-24
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

14 January 2003

Date of mailing of the international search report

24/01/2003

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 551 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Schuemacher, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/04931

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 02434 A (COCKERILL GEORGE STUART ;GUNTRIP STEPHEN BARRY (GB); GLAXO GROUP L) 22 January 1998 (1998-01-22) claims 1,31,32 ----	1-24
X	WO 97 03069 A (GLAXO GROUP LTD ;COCKERILL GEORGE STUART (GB); CARTER MALCOLM CLIV) 30 January 1997 (1997-01-30) page 5 -page 9; claims 1,17 -----	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/04931

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9609294	A	28-03-1996	AU 3482495 A EP 0782570 A1 NO 9609294 A1 JP 10505600 T TR 960233 A2 ZA 9507853 A	09-04-1996 09-07-1997 28-03-1996 02-06-1998 21-06-1996 18-03-1997
WO 9615118	A	23-05-1996	AT 175962 T AU 703328 B2 AU 3813095 A CA 2200871 A1 DE 69507505 D1 DE 69507505 T2 DK 790986 T3 EP 0790986 A1 ES 2128092 T3 FI 971970 A WO 9615118 A1 JP 10508616 T NO 972152 A NZ 294917 A US 5821246 A ZA 9509572 A	15-02-1999 25-03-1999 06-06-1996 23-05-1996 04-03-1999 02-06-1999 13-09-1999 27-08-1997 01-05-1999 07-05-1997 23-05-1996 25-08-1998 12-05-1997 25-02-1999 13-10-1998 13-05-1996
WO 9730034	A	21-08-1997	AU 707339 B2 AU 1612697 A CA 2242102 A1 CN 1211240 A EP 0880507 A1 WO 9730034 A1 JP 2000504713 T NO 983707 A NZ 330816 A US 6399602 B1 US 5866572 A ZA 9701231 A	08-07-1999 02-09-1997 21-08-1997 17-03-1999 02-12-1998 21-08-1997 18-04-2000 13-10-1998 26-05-2000 04-06-2002 02-02-1999 14-08-1997
WO 9802434	A	22-01-1998	AT 227283 T AU 3766897 A DE 69716916 D1 WO 9802434 A1 EP 0912559 A1 JP 2000514806 T US 2002147214 A1 US 6391874 B1 ZA 9706147 A	15-11-2002 09-02-1998 12-12-2002 22-01-1998 06-05-1999 07-11-2000 10-10-2002 21-05-2002 11-01-1999
WO 9703069	A	30-01-1997	AU 6613996 A WO 9703069 A1 EP 0843671 A1 HR 960316 A1 JP 11508906 T ZA 9605935 A	10-02-1997 30-01-1997 27-05-1998 28-02-1998 03-08-1999 12-02-1998